

A Rauzy fractal unbounded in all directions of the plane

Mérodie Andrieu

Advisors : Pierre Arnoux & Julien Cassaigne
Institut de Mathématiques de Marseille

OWCWS (day of short talks)
online, February 2021

I - MOTIVATIONS : understand the **dynamics** of multiD continued fraction algorithms.

II - RESULTS : **construction** of an Arnoux Rauzy word whose Rauzy fractal is unbounded in **all directions**

III - MAIN IDEAS : study the set of **differences of abelianized factors** of all Arnoux-Rauzy words.

I - Motivations

Regular continued fraction algorithm & Sturmian words

Subtractive continued fraction algorithm = iteration of the Farey map :

$$\begin{array}{lll}
 (\mathbb{R}^+)^2 & \rightarrow & (\mathbb{R}^+)^2 \\
 (x, y) & \mapsto & \begin{array}{l} (x - y, y) \\ (x, y - x) \end{array} \quad \begin{array}{l} \text{if } x \geq y, \\ \text{otherwise.} \end{array}
 \end{array}$$

The **symbolic trajectories** under this dynamical systems give rise to the class of **Sturmian words**.

Sturmian words enjoy multiple [combinatorial, geometrical, dynamical] characterizations.

Balance characterization :

Sturmian words are exactly the aperiodic binary words for which any two factors of same length contain, with +/- 1, the same number of 0s.

Ex

A word starting with $w = 001000100100010001001\dots$ is possibly Sturmian.

A word starting with $w = 0\underline{1}1011\underline{1}00\dots$ is not.

Regular continued fraction algorithm & Sturmian words

Consequences :

1. The letters 0 and 1 are uniformly distributed with respect to a probability measure ν on $\{0, 1\}$.
2. Stronger : the difference between the observed frequency of 0s among the N first letters of w and its expected value $\nu(0)$ is bounded above by $1/N$.

Geometrically, the "broken line" made of the points $P_N := \sum_{n=0}^N e_{w[n]}$, where (e_0, e_1) is the usual basis of \mathbb{R}^2 , remains at bounded distance from its average direction.




Figure – The broken line of 01000100100...

→ Sturmian words are used to approximate lines with irrational slopes.

MultiD continued fraction algorithms

Since Jacobi, several algorithms have been proposed to generalize continued fractions to triplets of nonnegative real numbers.

Such algorithms should make it possible to **simultaneously** and **efficiently** approach two real numbers with a sequence of pairs of rational numbers.

The Arnoux-Rauzy algorithm

$$F_{AR} : \begin{array}{ll} (\mathbb{R}^+)^3 & \rightarrow (\mathbb{R}^+)^3 \\ (x, y, z) & \mapsto \begin{array}{ll} (x - y - z, y, z) & \text{if } x \geq y + z, \\ (x, y - x - z, z) & \text{if } y \geq x + z, \\ (x, y, z - x - y) & \text{if } z \geq x + y. \end{array} \end{array}$$

This algorithm gives rise to the class of Arnoux-Rauzy words, which are, from the combinatorial view point, the generalization of Sturmian word.

In particular, all Arnoux-Rauzy words admit a letters frequencies vector $(\nu(1), \nu(2), \nu(3))$.

→ What can we say of the 3D broken line ?

Properties of the Arnoux-Rauzy broken line ?

Old belief : "the broken line of any Arnoux-Rauzy word remains at bounded distance from its average direction ; or equivalently, all Arnoux-Rauzy *Rauzy fractals* are bounded."

→ **disproved** in 2000 by Cassaigne, Ferenczi and Zamboni.

Today, we barely know nothing about the geometry or the topology of these unbounded Rauzy fractals.

Modern belief : "The broken line of any Arnoux-Rauzy word remains at bounded distance from an hyperplane containing its average direction ; or equivalently, all Arnoux-Rauzy *Rauzy fractals* are trapped between two parallel lines of the plane."

→ This is suggested by the **Oseledets theorem**. Indeed, if the Lyapunov exponents of the product of matrices associated with w exist, one of these exponents at least is nonpositive since their sum is equal to zero.

This belief is wrong.

II - Main results

Arnoux-Rauzy words ["S-adic definition"]

The **Arnoux-Rauzy substitutions** are :

$$\begin{array}{lll}
 \sigma_1 : & 1 \mapsto 1 & \sigma_2 : & 1 \mapsto 21 & \sigma_3 : & 1 \mapsto 31 \\
 & 2 \mapsto 12 & & 2 \mapsto 2 & & 2 \mapsto 32 \\
 & 3 \mapsto 13 & & 2 \mapsto 23 & & 3 \mapsto 3.
 \end{array}$$

Fact : if $(s_n)_{n \in \mathbb{N}}$ is a sequence of Arnoux-Rauzy substitutions containing **infinitely many occurrences of σ_1, σ_2 and σ_3** , then the sequence of finite words $(s_0 \circ \dots \circ s_{n-1}(\alpha))$, with $\alpha \in A$, converges to an infinite word w_0 which does not depend on α .

These infinite words w_0 are called **standard Arnoux-Rauzy words**.

Ex : $w_{Trib} = (\sigma_1 \circ \sigma_2 \circ \sigma_3)^\omega(1) = 12131211121312121312111213121312\dots$

An infinite word w is an **Arnoux-Rauzy word** if it has the same language than a standard Arnoux-Rauzy word w_0 .

Letters frequencies

Let $w \in A^{\mathbb{N}}$ and $i \in A$.

The **frequency** of i in w is the limit, if it exists, of the proportion of i in the sequence of growing prefixes of w : $f_w(i) = \lim_{n \rightarrow \infty} \frac{|p_n(w)|_i}{n}$.

We denote by $f_w = (f_w(i))_{i \in \mathfrak{A}}$ the **vector of letters frequencies** of w , if it exists.

Fact : All Arnoux-Rauzy words admit a vector of letters frequencies.

Theorem (A. 20 ; Dynniov, Hubert & Skripchenko 20)

The vector of letters frequencies of an Arnoux-Rauzy word has rationally independent entries.

→ This result was conjectured by Arnoux and Starosta in 2013.

Discrepancy & Rauzy fractal

A natural question is to study the **difference** between the **predicted frequencies** of letters and their **observed occurrences**.

Definition : The **broken line** of w is $\mathcal{B}_w := \{\text{ab}(p_k(w)) \mid k \in \mathbb{N}\} \subset \mathbb{N}^3$, where :

- $p_n(w)$ is the prefix of length n of w ,
- $\text{ab}(u)$ is the **abelianized vector** of u , which counts the occurrences of each letter.

$$\text{Ex : } \text{ab}(p_6(w_{\text{trib}})) = \text{ab}(121312) = (3, 2, 1).$$

Denote by π_w the (oblique) projection parallel to $\mathbb{R}f_w$, onto Δ_0 (the plane of \mathbb{R}^3 with equation $x + y + z = 0$).

Definition - The Rauzy fractal of w is :

$$\mathcal{R}_w := \overline{\pi_w(\mathcal{B}_w)} \subset \Delta_0.$$



Figure – Rauzy fractal of $w_{trib} = (\sigma_1 \circ \sigma_2 \circ \sigma_3)^\omega(1)$.

Results

Theorem (A. 20)

There exists an Arnoux-Rauzy word whose Rauzy fractal is unbounded in all directions of the plane.

A similar result holds for **Cassaigne-Selmer words** and for **strict episturmian words** over a d -letter alphabet, for $d \geq 3$.

III - Main ideas for the construction

1. Reduce the problem to a combinatorial question

Lemma (1)

Let $w \in \{1, 2, 3\}^{\mathbb{N}}$. If for all $\vec{d} \in \mathbb{Z}^3$, there exist u and $v \in \mathcal{F}(w)$ such that $ab(u) - ab(v) = \vec{d}$, then, for any plane Π and for any $D \in \mathbb{R}^+$, there exists $k \in \mathbb{N}$ such that the euclidean distance between the point P_k , whose coordinates are $ab(p_k(w))$, and the plane Π satisfies $\text{dist}_{\mathbb{R}^3}(P_k, \Pi) \geq D$.

→ Can we construct an Arnoux-Rauzy word w_∞ such that :

$$\{ab(u) - ab(v) \mid u, v \in \mathcal{F}(w_\infty)\} = \mathbb{Z}^3 \quad ?$$

If so, the Rauzy fractal of w_∞ would be unbounded in all directions of the plane !

2. Deal with the combinatorial problem : construct w_∞ .

$$AR = \{\sigma_1, \sigma_2, \sigma_3\}.$$

Lemma (2)

For any $(a, b, c) \in \mathbb{Z}^3$, there exists $s \in AR^$ and there exist $u, v \in \mathcal{F}(s(1))$ that satisfy $ab(u) - ab(v) = (a, b, c)$.*

→ In particular, all Arnoux-Rauzy words whose directive sequence starts with s contain these two factors u and v .

Theorem (3)

There exists an Arnoux-Rauzy word w_∞ such that for all $(a, b, c) \in \mathbb{Z}^3$, there exist u and $v \in \mathcal{F}(w_\infty)$ satisfying $ab(u) - ab(v) = (a, b, c)$.

→ Constructed S-adically from Lemma 2 by a pumping strategy.

Technical part is Lemma 2.

Construction of w_∞ : technical part (glimpse)

Lemma (2)

For any $(a, b, c) \in \mathbb{Z}^3$, there exists $s \in AR^*$ and there exist $u, v \in \mathcal{F}(s(1))$ that satisfy $ab(u) - ab(v) = (a, b, c)$.

Let \mathcal{G} the infinite oriented graph \mathcal{G} whose set of vertices is \mathbb{Z}^3 and whose edges maps triplets to their images by one the 15 following applications.

For $\delta \in \{-2, -1, 0, 1, 2\}$ and $i \in \{1, 2, 3\}$:

$$\begin{aligned} \tau_{i,\delta} : \mathbb{Z}^3 &\rightarrow \mathbb{Z}^3 \\ (x_j)_{j \in \{1,2,3\}} &\mapsto (y_j)_{j \in \{1,2,3\}} \quad \text{where } \begin{cases} y_i = x_1 + x_2 + x_3 + \delta \\ y_j = x_j \text{ for } j \neq i. \end{cases} \end{aligned}$$

Ex for $i = 1$ and $\delta = -2$:

$$\begin{aligned} \tau_{1,-2} : \mathbb{Z}^3 &\rightarrow \mathbb{Z}^3 \\ (a, b, c) &\mapsto (a + b + c - 2, b, c) \end{aligned}$$

Construction of w_∞ : technical part (glimpse)

Why this graph ?

Fact : If $\tau_{i,\delta}(a, b, c) = (d, e, f)$ and if (a, b, c) is the difference of two abelianized factors of an Arnoux-Rauzy word w , then (d, e, f) is the difference of two abelianized factors of [the Arnoux-Rauzy word] $\sigma_i(w)$.

A generic example.

For $i = 1$ and $\delta = -2$.

If $u, v \in F(w)$ are **nonempty** and satisfy $\text{ab}(u) - \text{ab}(v) = (a, b, c)$, then the word $\sigma_1(u)$ starts with the letter 1, and each occurrence of $\sigma_1(v)$ in $\sigma_1(w)$ is immediately followed by the letter 1.

Therefore, the words $\tilde{u} = 1^{-1}\sigma_1(u)$ (the word u without its initial 1) and $\tilde{v} = \sigma_1(v)1$, are factors of $\sigma_1(w)$ and satisfy

$$\text{ab}(\tilde{u}) - \text{ab}(\tilde{v}) = (a + b + c - 2, b, c) = \tau_{1,-2}(\text{ab}(u) - \text{ab}(v)).$$

→ We are going to study the paths in this graph...

Construction of w_∞ : technical part (glimpse)

A careful study of \mathcal{G} then shows :

Lemma

All triplets in \mathbb{Z}^3 can be reached from the vertex $(0, 0, 0) \in \mathbb{Z}^3$, moving through a finite number of edges.

We conclude the proof by observing that $(0, 0, 0)$ is the difference between $\text{ab}(u)$ and itself - where u is an Arnoux-Rauzy factor that can be chosen **as long as we need**.

Remark. \mathcal{G} is not strongly connected.

Thank you !