

DERIVATION OF A GYROKINETIC MODEL. EXISTENCE AND UNIQUENESS OF SPECIFIC STATIONARY SOLUTIONS.

PHILIPPE GHENDRIH

CEA, DRFC
F-13108 Saint-Paul lez Durance, France

MAXIME HAURAY

LJLL, Paris 6, 7 & CNRS
175 rue du Chevaleret, 75013 Paris, France

ANNE NOURI

LATP, Université de Provence
39 rue F.Joliot Curie, 13453 Marseille Cedex 13, France

Abstract.

A finite Larmor radius approximation is derived from the classical Vlasov equation, in the limit of large (and uniform) external magnetic field. We also provide an heuristic derivation of the electroneutrality equation in the finite Larmor radius setting. Existence and uniqueness of a solution is proven in the stationary frame for solutions depending only on the direction parallel to the magnetic field and factorizing in the velocity variables.

Introduction. The ITER project is a challenge to the growing need of new sources of energy. It aims at producing energy by nuclear fusion. Nuclear reactions take place in a tokamak, where a high temperature plasma is confined. So far, confined plasmas are performed with relatively short energy confinement times due to microscale instabilities that generate turbulent transport [III99]. It is observed that the characteristic frequencies of these instabilities is several orders of magnitude smaller than the ion Larmor gyration frequency governed by the strong magnetic field. Studies of nuclear fusion in tokamaks are in full expansion, both experimentally and theoretically. Kinetic models are appropriate for studying the core of the plasma since the collisions have a very weak effect in these hot and low density plasmas. Physicists use gyrokinetic models and especially the finite Larmor radius approximation to model the core of the plasma [GIVT09]. Taking into account the fast Larmor gyration of the charged particles that characterizes magnetic confinement, these models allow one to average over that fast gyration and reduce the 6D kinetic problem to a 5D gyrokinetic one. That property is especially interesting for numerical simulation. In this paper the finite Larmor radius approximation is derived from the Vlasov equation, in the limit of large uniform magnetic field and with an external electric field. Because of the homogenization on the fast Larmor gyration, the limit equation (1.10) is written in 5D gyro-coordinates $(x_g, v_{\parallel}, |v_{\perp}|)$ defined in (1.11). These coordinates are the position of the so-called particle guiding center, x_g in the 3D space together with the parallel velocity v_{\parallel} and the amplitude of transverse velocity $|v_{\perp}|$ that is proportional to the magnetic moment, an adiabatic invariant of the particle

¹2000 Mathematics Subject Classification. 41A60, 76P05, 82A70, 78A35.

²Key words. Plasmas, gyrokinetic model, steady state solutions.

motion in the strong magnetic field limit (given a constant magnetic field) . Its mathematical structure is a combination of the Vlasov equation in the direction parallel to the magnetic field and of the Euler equation in the perpendicular direction, where the original fields are replaced by the corresponding gyro-average fields.

To close the system, physicists use the electroneutrality equation $n_e = Zn_i$ where n_e stands for the electron density and n_i the ion density, each ion having a charge Z . In the following $Z = 1$ will be considered with no loss of generality. It can be shown that the electroneutrality equation is in fact the the Poisson equation solved on scales that are significantly larger than the Debye length. Since the latter governs the Laplacian term of the Poisson equation, the electroneutrality equation is an appropriate approximation on scales of the order or larger than the Debye length. Moreover, taking into account the difference between the density of particles and that of guiding centers leads one to introduce a polarization correction due to the non uniform distribution of particles on gyro-circles. In the second part of this paper, the electroneutrality equation (1.25) is carefully written. The coupling of the 5D gyrokinetic Vlasov equation (1.10) to the electroneutrality equation (1.25) is the model used for instance in the GYSELA code, a project that aims at modeling the turbulent transport in fusion plasmas [GSG⁺06b, GSG⁺06a]. While in the GYSELA code there is the possibility to use a collision operator we concentrate here on the actual Vlasov equation with no collisions.

A difficulty raised by the electroneutrality equation taken as such is the lack of an explicit regularization term for the electric potential. Consequently, the regularity of the latter is not sufficient to ensure a mathematical solution of the Vlasov transport equation. The analysis of the well-posedness of limit model based on the electroneutrality equation thus seems impossible today. For that reason we restrict ourselves to the 1D Vlasov equation (without gyro-kinetic effects), coupled to the electroneutrality condition. That model is a particular case of the 3D model, provided the solutions do not depend on the direction perpendicular to the magnetic field. Even for that simple kinetic model, not much is known about solutions. The difficulty lies in the fact that the force term is proportional to the derivative of the density in the field direction. Indeed, while the Vlasov equation ensures that all L^p -norms of the distribution function can be bounded, there is no control on the norms of its derivatives. In this paper, we prove the existence and uniqueness of a steady state solution in a slab geometry, therefore between two boundary conditions, provided one fulfills some conditions, in particular that they are no particles trapped between the two boundaries.

1 Derivation of the finite Larmor radius approximation

The ion distribution in low density plasmas submitted to a magnetic field is well described by the Vlasov equation. The latter is valid provided one can neglect the two-particle distribution function altogether [Nic83] so that the evolution of the standard distribution function is governed by the Liouville equation for the conservation of particles

$$\frac{\partial f}{\partial t} + v \cdot \nabla_x f + \frac{Ze}{m_i} (E(t, x) + v \times B) \cdot \nabla_v f = 0 , \quad (1.1)$$

where f is the phase space density depending on time, position (in the domain D of the plasma) and velocity (in \mathbb{R}^3) of the ions, and where Ze and m_i are the ion charge and mass respectively. As stated in the introduction, we shall consider $Z = 1$ in the following with no loss of generality. A priori, the electric and magnetic fields are governed by the Maxwell equations, but a scale analysis allows one to approximate, and thus simplify, these laws. This will be made clear in the following.

For a strong external magnetic field, the charged particles exhibit a fast rotation motion around the magnetic field lines. The frequency of that gyration, the Larmor frequency, is several orders of magnitude larger than the observed frequency range of turbulence. Furthermore, in the quasineutral limit this frequency is larger than all other frequencies of the particle dynamics, thus providing the means for an efficient scale separation. It is noteworthy that this is the basis of magnetic confinement that is presently realized in devices such as the tokamaks. In this framework, one separates two parts in the particle motion, on the one hand the slow motion of the center of the Larmor gyration, and, on the other hand, the fast gyro-motion. The phase space reduction achieved by only considering the slow motion, thus ignoring the fast gyro-motion,

leads one to the 5D gyrokinetic model. These are usually derived by physicists either by averaging the single particle dynamics ([Cat78] and [CTB81]) or by a Lie-transform perturbative approach ([Hah88], [BH07] and [GLV08]). Mathematically, this is achieved by looking at the limit of equation (1.1) when the modulus of the external magnetic field $|B_{ext}|$ tends to infinity. However, different models can be obtained, depending on the way the other control parameters vary when $|B_{ext}| \rightarrow \infty$. For a magnetized plasma, two relevant scales characterize the limit regimes that have been discussed. These are the Debye length λ_D and the thermal Larmor radius of the ions ρ_{th} . The Debye length weights the Laplacian in the Poisson equation and thus defines the transition from the non-neutral description of the plasma required on the sub-Debye scales from the quasineutral plasma description for scales larger than the Debye length. The thermal Larmor radius is the radius of the fast Larmor gyration for ions with average speed v_{th} . The averaging over the small time scales governed by the ion Larmor frequency then tends to translate into an averaging over the ion Larmor scale. The finite Larmor radius terms are therefore introduced to take into account this rather weak cut-off effect. The radius of the electrons gyration is generally much smaller for comparable ion and electron temperatures and will be neglected. The two scales discussed above are defined by

$$\lambda_D = \sqrt{\frac{\varepsilon_0 T_e}{n e^2}}, \quad \rho_{th} = \frac{m_i v_{th}}{e B}, \quad (1.2)$$

where the electron thermal energy T_e is introduced rather than that of the ions since it is more appropriate to characterize the large electron mobility and therefore strong response to the electric field. Let us introduce the characteristic scale of the system L , for instance introduced by the boundary conditions. The large magnetic field limit then corresponds to the vanishing ion Larmor radius limit, namely the gyro-center approximation where $\rho_{th}/L \rightarrow 0$. Within this framework, E. Grenier uses pseudo-differential calculus to prove [Gre97] the convergence of a 2D fluid model towards an incompressible fluid model when ε goes to zero with $\rho_{th}/L = \varepsilon$ and $(\lambda_D/L)^2 = \varepsilon$. For the same scaling and cold initial distributions, *i.e.* the approximation of a Dirac measure at velocity zero for the distribution function, Y. Brenier proved in [Bre00] with modulated energy technics the convergence of solutions of the Vlasov-Poisson system towards dissipative solutions (introduced by P.-L. Lions) of the Euler equation, for weel-prepared initial data. F. Golse and L. St-Raymond used the same technics in [GSR03] to prove the convergence of a Vlasov-Poisson model on the torus towards the 2D1/2 Euler equation. This is done for a vanishing ε with $\rho_{th}/L = \varepsilon$ and using another scaling for the Debye length, namely $\lambda_D/L = \varepsilon$. This property is restricted to a torus size such that there is no resonant oscillation. This particular case appears to be relevant for electrons in a region close to the tokamak boundary.

Derivations have also been performed in the gyrocenter approximation, ρ_{th}/L_\perp small but finite. Here L_\perp is a carcteristic length in the perpendicular direction, which is usually chosen smaller than the one in the parallel direction in finite Larmor radius approximation. In [FS00a] (see also [FS00b] for a short version) E. Frénod and E. Sonnendrücker studied the convergence of a linear Vlasov equation (external magnetic field), in this finite Larmor radius limit (large B but finite Larmor radius), using two scale convergence methods. M. Bostan studied the 2D strong magnetic field limit, using Hilbert expansion technics, in a setting of finite Debye length and Larmor radius, $\lambda_D/(\varepsilon L) = \rho_{th}/(\varepsilon L) = 1$, where ε is small but finite [Bos09] and obtained strong convergence results for regular initial conditions. Here, we further examine the linear case studied by Frénod and Sonnendrücker, and obtain a simpler and more complete derivation. The finite Larmor radius approximation is correct for fusion plasmas, including the core and most of the edge plasma. However, the ion Larmor radius is much larger than the Debye length throughout the plasma. An interesting derivation should be done in the framework of present gyrokinetic calculations, namely $\rho_{th}/(\varepsilon L) = 1$, $\lambda_D/(\varepsilon L) \rightarrow 0$, yet nothing has been done in this very demanding limit (see the beginning of 1.2 for more details).

The resolution of equation (1.1) is known in the case where the fields E and B are external and C^1 or at least BV . In the BV -case, one can use the DiPerna Lions theory of transport equation [DL89b] with its latest developments ([Amb04]). In the case where the electric potential is given by self-interaction with the usual Poisson law, and the magnetic field B is still considered as external, results of existence and uniqueness obtained for the Vlasov equation without magnetic field may be used with the appropriate modifications. We refer to [LP91], [Hor93]. In the case where the self-induced magnetic field is not negligible any more, we refer to the work [DL89a] of DiPerna and Lions about the Vlasov-Maxwell equation. In the case of an electric field given by an electroneutrality equation, nothing is known from the mathematical point of view.

1.1 The finite Larmor radius gyro-kinetic approximation. Rigorous approach.

For the sake of simplicity, we neglect the variation of the external magnetic field B_{ext} . In this cylindrical approximation, the curvature of the magnetic field is not considered, and one neglects the exploration of the magnetic field variation by the particles during the fast cyclotron motion. With respect to the relevant scales this effect is of the order of the Larmor radius ρ_{th} divided by the characteristic radius of the magnetic field curvature, namely the tokamak major radius R . Although ρ_{th}/R is a small parameter, this approximation is too strong with respect to many aspects of the tokamak physics, in particular since curvature effects are considered by physicists as the cause of a large class of micro-instabilities. The strong impact of such a small parameter can be traced back to a symmetry breaking of the system since the curvature governs the leading term that produces a charge separation. The present simplification must be regarded as a first step that is only valid when one addresses issues that do not lead one to symmetry breaking. Moreover, in order to avoid problems with boundary conditions, which are complex to handle, a periodic setting is used. In other words, we work on the torus \mathbb{T}^3 in space, and \mathbb{R}^3 in speed, with a constant magnetic field $B = |B|(0, 0, 1)$.

Next, we define dimensionless variables

$$t' = \frac{t}{\tau}, \quad x'_{\parallel} = \frac{x_{\parallel}}{L_{\parallel}}, \quad x'_{\perp} = \frac{x_{\perp}}{L_{\perp}}, \quad v' = \frac{v}{v_{th}}, \quad E' = \frac{E}{E_0},$$

with characteristic time τ , parallel length L_{\parallel} , perpendicular length L_{\perp} , velocity v_{th} and electric field E_0 . In this new set of dimensionless variables the Vlasov equation (1.1) may be rewritten as:

$$\frac{\partial f}{\partial t'} + \frac{\tau v_{th}}{L_{\parallel}} \partial_{x'_{\parallel}} f + \frac{\tau v_{th}}{L_{\perp}} \cdot \nabla_{x'_{\perp}} f - \frac{\tau e E_0}{m_i v_{th}} E \cdot \nabla_{v'} f - \frac{\tau e B}{m_i} v^{\perp} \cdot \nabla_{v'} f = 0, \quad (1.3)$$

where the subscript \perp (resp. \parallel) stands for the projection on the perpendicular (resp. parallel) direction, and the superscript \perp stands for the projection on the plane orthogonal to the field line and the rotation by $-\pi/2$ around the field line direction: $v^{\perp} = (v_2, -v_1, 0)$ if $v = (v_1, v_2, v_3)$. The time scale τ can conveniently be defined to reduce the number of control parameters in Eq.(1.3) by setting:

$$\frac{\tau v_{th}}{L_{\parallel}} = 1$$

In a similar fashion, the normalization of the electric field can be used to define the third control parameter, hence:

$$\frac{\tau e E_0}{m_i v_{th}} = 1$$

The two remaining control parameters are then L_{\parallel}/L_{\perp} for the third term in Eq.(1.3) and L_{\parallel}/ρ_{th} for the last term. It will be assumed here that both parameters L_{\perp} and ρ_{th} exhibit the same asymptotic behavior $L_{\perp} \propto \rho_{th}$. Let us define the perpendicular scale as $L_{\perp} = \rho_{th} 2\pi / (k_{\perp} \rho_{th})$. In this expression, k_{\perp} is the typical wave vector of the cross-field fluctuations. For a sufficiently small Larmor radius, one can then assume that the turbulence follows the so-called gyro-Bohm scaling such that $k_{\perp} \rho_{th}$ is a constant [LEHT02]. In particular it does not depend on the magnitude of the magnetic field. One then finds the following relationship between the two control parameters

$$\frac{L_{\parallel}}{\rho_{th}} = \frac{L_{\parallel}}{L_{\perp}} \frac{1}{k_{\perp} \rho_{th}} = \frac{1}{\varepsilon},$$

that can be grasped by the following relation whenever one drops the proportionality constant

$$\frac{L_{\parallel}}{\rho_{th}} = \frac{L_{\parallel}}{L_{\perp}} = \frac{1}{\varepsilon}.$$

In practice, one could also define the normalization scale as $L_{\perp} = \rho_{th}$ so that the two control parameters would be exactly the same. In this discussion, we have considered a transverse scale related to the fluctuation properties that allows one to characterize one of the terms in the Vlasov equation. However, other control parameters must be considered to account for the boundary conditions. The so-called ρ_* parameter used in magnetic fusion is such a parameter since $\rho_* = \rho_{th}/a$ where a is the plasma minor radius, the scale related to

the boundary conditions while ρ_{th} is the fluctuation scale. Relating the parameter ε to ρ_* introduces aspect ratios that stem from the periodic boundary conditions.

$$\varepsilon = \rho_* \frac{a}{L_{\parallel}}$$

The ratio a/L_{\parallel} depends of the safety factor and the tokamak aspect ratio in the actual tokamak geometry but represents the ratio of the sizes of the domain in the radial and parallel direction with the present setting. The parameter ε can also be expressed in terms of the slow and fast characteristic times of the particle motion. The slow time τ introduced in the normalization of Eq.(1.3), is the characteristic time to explore the tokamak geometry along the parallel direction and the fast time is due to the gyration motion, hence:

$$\varepsilon = \frac{1}{\tau\Omega}$$

In this last step one thus finds that the ion Larmor frequency $\Omega = eB/m_i$ is of order $(\tau\varepsilon)^{-1}$. In other words, one thus assumes that the ion Larmor frequency is much larger than the parallel connection frequency τ^{-1} . That is a valid assumption in all regions of a tokamak. Indeed, for ITER conditions, the ion Larmor frequency is of the order of $2 \cdot 10^8 Hz$ and the connection time τ ranges from $10^{-3}s$ to $10^{-4}s$. These parameters mainly depend on the magnitude of the magnetic field and on the size of the device so that ε characterizes a given fusion device, $\varepsilon \approx 10^{-5}$ for ITER parameters.

Dropping the primes, we obtain the following rescaled version of the Vlasov equation (1.1),

$$\frac{\partial f}{\partial t} + v_{\parallel} \cdot \partial_{x_{\parallel}} f + E \cdot \nabla_v f + \frac{1}{\varepsilon} (v_{\perp} \cdot \nabla_{x_{\perp}} f + v^{\perp} \cdot \nabla_{v_{\perp}} f) = 0, \quad (1.4)$$

When $\varepsilon \rightarrow 0$, the largest terms are those proportional to the factor $1/\varepsilon$. Retaining the latter, we obtain a transport equation associated to the following system of ODE in the plane transverse to the magnetic field,

$$\dot{x} = \frac{1}{\varepsilon} v_{\perp}, \quad \dot{v} = \frac{1}{\varepsilon} v_{\perp}^{\perp}. \quad (1.5)$$

As B is homogeneous, the trajectories are circles of center $x_g = x_{\perp} + v_{\perp}^{\perp}$, and radius $|v_{\perp}|$ covered with frequency ε^{-1} . The global motion is the sum of this very quick motion of gyration and a slower and more complicated motion (with velocity of order one). In the limit of large $|B|$, particles are assumed to be evenly distributed on the gyro-circles, and their motion is the sum of a drift in the perpendicular direction, and a classical acceleration in the parallel direction. Heuristically, the electric drift v_E may be obtained from the Newton law in normalized variables,

$$\dot{v} = E + \frac{1}{\varepsilon} v^{\perp}, \quad (1.6)$$

Given the assumption that the particles have a fast motion of gyration, one can integrate the previous equation over a period of gyration, at lowest order, v_E is given by the averaged velocity such that:

$$0 \approx \langle E \rangle + \frac{1}{\varepsilon} v_E^{\perp} \quad \text{so that } v_E = \varepsilon \langle E^{\perp} \rangle. \quad (1.7)$$

In the latter expression one readily recognizes the usual form of the electric drift velocity $v_E = E \times B/B^2$, however where the electric field is averaged over a period $\langle E \rangle$. One can show that this average of the field E corresponds to an average over a circle of radius $|v_{\perp}|$ in the perpendicular plane. Provided the only dependence on the gyrophase stems from the particle motion, this average translates into a Bessel operator defined as:

$$J_{\rho_L}^0 h(x_g) = \frac{1}{2\pi} \int_0^{2\pi} h(x_g + \rho_L e^{i\varphi_c}) d\varphi_c, \quad (1.8)$$

where ρ_L is the ion Larmor radius and $e^{i\varphi_c} = (\cos \varphi_c, \sin \varphi_c, 0)$. We also introduce the operator $\tilde{J}_{\rho_L}^0$ which stands for a position-velocity version of this average,

$$\tilde{J}_{\rho_L}^0 g(x_g, \rho_L, v_{\parallel}) = \frac{1}{2\pi} \int_0^{2\pi} g(x_g + \rho_L e^{i\varphi_c}, \rho_L e^{i(\varphi_c - \frac{\pi}{2})} + v_{\parallel} e_{\parallel}) d\varphi_c, \quad (1.9)$$

where $e_{\parallel} = (0, 0, 1)$.

Given the average on the gyrophase φ_c , the motion is reduced to a $5D$ space. The gyration radius $\rho_L = |v_{\perp}|$ (given the chosen normalization) is related to the magnetic moment $\mu = m_i \rho_L^2 / (2eB)$, a quantity that is an adiabatic invariant, *i.e.* a constant of motion in the large B limit. One can show that this invariant is the conjugate variable of the gyrophase. Here, as B is homogeneous, $\rho_L = |v_{\perp}|$ will remain constant in the limit $|B_{ext}| \rightarrow +\infty$. We thus obtain a $4D + 1D$ model, *i.e.* a $5D$ model with no dynamics in the variable derived from the magnetic moment. This reduction of the phase space is very interesting for numerical simulations. The following theorem states this reduction precisely.

Theorem 1.1 *Let us assume that f_{ε}^0 is uniformly bounded in L^q , $q > 1$, that it weakly converges toward $f^0 \in L^q$, and that E is a gradient and belongs to $L^p_{t,x}$ where $p^{-1} + q^{-1} = 1$. For each $\varepsilon > 0$, let f_{ε} be a solution of (1.4) with the initial condition f_{ε}^0 . Then, up to a subsequence (ε_n) , $(f_{\varepsilon}(t, x_g - v^{\perp}, v))$ weakly converges to \bar{f} in L^q , where \bar{f} only depends on $(t, x_g, v_{\parallel}, |v_{\perp}| = \rho_L)$ and is solution of*

$$\frac{\partial \bar{f}}{\partial t} + v_{\parallel} \partial_{x_{\parallel}} \bar{f} + J_{\rho_L}^0 E_{\parallel} \partial_{v_{\parallel}} \bar{f} + (J_{\rho_L}^0 E_{\perp})^{\perp} \cdot \nabla_{x_g} \bar{f} = 0, \quad (1.10)$$

with the initial condition $\tilde{J}_{\rho_L}^0(f^0)$.

Moreover, if $J_{\rho_L}^0 E \in BV(\mathbb{R}^3)$, for *a.e.* $\rho_L > 0$, then there is no need to extract a subsequence and the limit \bar{f} is the unique solution of (1.10) with the initial condition $\tilde{J}_{\rho_L}^0(f^0)$.

Remark 1.1 *The Bessel operator $J_{\rho_L}^0$ has some regularization properties. It goes from H^s to $H^{s+1/2}$ for all s , so that if $E \in H_{\parallel}^1 \times H_{\perp}^{1/2}$, then $J_{\rho_L}^0 E \in H^1$, a condition that ensures the uniqueness of the solution of equation (1.10).*

Remark 1.2 *The operator \tilde{J}^0 is important to perform the adaptation of the $6D$ initial condition to the $5D$ limit model. Indeed, the very fast Larmor gyration creates an initial layer that instantaneously adapts the initial condition to the limit model.*

Proof of Theorem 1.1.

The phase space in *position-velocity coordinates* is not well adapted to perform the fast gyration averaging. To handle it more easily, it is convenient to change the system of coordinates and consider the *gyro-coordinates* defined by

$$x_g = x + v^{\perp}, \quad v_g = v. \quad (1.11)$$

x_g is the position of the gyro-center and $\rho_L = |v^{\perp}|$ is the ion Larmor radius. To express the gradient in x and v in this new system of coordinates, one can conveniently remark that:

$$\nabla_x h(dx_g - dv_g^{\perp}) + \nabla_v h dv_g = \nabla_{x_g} \bar{h} dx_g + \nabla_{v_g} \bar{h} dv_g, \quad (1.12)$$

for any smooth function h , with the function \bar{h} defined by $\bar{h}(x_g, v_g) = h(x, v)$. Then $\nabla_x = \nabla_{x_g}$ and $\nabla_v = \nabla_{v_g} - \nabla_{x_g}^{\perp}$, where $\nabla_{x_g}^{\perp} = (\partial_{x_{g,2}}, -\partial_{x_{g,1}}, 0)$. Note that ∇^{\perp} stands for the gradient vector rotated by $-\pi/2$ and not $\pi/2$. The gradients are taken at the corresponding points. For instance, the first equality reads $\nabla_x h(x, v) = \nabla_{x_g} \bar{h}(x_g, v_g)$. The anti-symmetry of \perp , $a \cdot b^{\perp} = -a^{\perp} \cdot b$, has been used (and we shall make a wide use of it in the sequel). Given these relations, (1.4) can be modified leading one to an equation satisfied by the function $\bar{f}_{\varepsilon}(t, x_g, v_g) = f_{\varepsilon}(t, x, v)$,

$$\begin{aligned} & \frac{\partial \bar{f}_{\varepsilon}}{\partial t} + v_{\parallel} \partial_{x_{\parallel}} \bar{f}_{\varepsilon} + E_{\parallel}(t, x_g - v_g^{\perp}) \partial_{v_{\parallel}} \bar{f}_{\varepsilon} \\ & + E_{\perp}(t, x_g - v_g^{\perp}) \cdot (\nabla_{v_g} \bar{f}_{\varepsilon} - \nabla_{x_g}^{\perp} \bar{f}_{\varepsilon}) + \frac{1}{\varepsilon} v_g^{\perp} \cdot \nabla_{v_g} \bar{f}_{\varepsilon} = 0, \end{aligned} \quad (1.13)$$

with initial condition \bar{f}^0 . Here the subscript \perp stands for the perpendicular components to the magnetic field, for instance $E_{\perp} = (E_1, E_2, 0)$. Barred quantities are functions of the gyro-coordinates.

Since (1.6) is a conservative transport equation, *i.e.* it may be written as $\partial_t f + \text{div}(\dots) = 0$, the L^q -norms of f_ε and then of \bar{f}_ε are conserved. Thus, $\|\bar{f}_\varepsilon\|_{L_t^\infty(L_{x,v}^q)}$ is uniformly bounded. Then, up to the extraction of a subsequence, we may assume that \bar{f}_ε weakly converges towards some $\bar{f} \in L_t^\infty(L_{x,v}^q)$. Upon multiplying equation (1.13) by ε , in the limit $\varepsilon \rightarrow 0$, we obtain

$$v_g^\perp \cdot \nabla_{v_g} \bar{f} = 0,$$

All the other terms from Eq.(1.13) are bounded in the sense of distributions and their product with ε therefore vanishes in the limit $\varepsilon \rightarrow 0$. This reduced form of the Vlasov equation implies that the only dependence of \bar{f} on $v_{g,\perp}$ is on $|v_{g,\perp}|$. Hence with no dependence on the gyrophase.

Let us now consider equation (1.13) for \bar{f}_ε , when integrated against a smooth test function ϕ with support in time avoiding $t = 0$ (we will handle the initial conditions later). Let us further assume that the dependence on $v_{g,\perp}$ is restricted to a dependence on $|v_{g,\perp}|$. For such a function, one readily finds that

$$\frac{1}{\varepsilon} \int \bar{f}_\varepsilon v_g^\perp \nabla \phi dx_g dv_g = 0$$

for symmetry reasons. As a consequence, the projection of \bar{f}_ε takes the following form

$$\int \bar{f}_\varepsilon (\partial_t \phi + v_{g,\parallel} \partial_{x_{g,\parallel}} \phi + E_\parallel \partial_{v_{g,\parallel}} \phi + E_\perp \cdot \nabla_{v_g} \phi - E^\perp \cdot \nabla_{x_g} \phi) dx_g dv_g dt = 0, \quad (1.14)$$

keeping in mind that E is calculated at the point $x_g - v_g^\perp$. We have also used the relation $a^\perp \cdot b = -a \cdot b^\perp$ as well as the fact that equation (1.13) may be written in a conservative form because (if J is the matrix of the linear map $v \rightarrow v^\perp$),

$$\begin{aligned} \text{div}_{v_g} (E(t, x - v_g^\perp)) &= \text{Tr}((\nabla E) J) = -\partial_{x_2} E_1 + \partial_{x_1} E_2 = 0, \\ \text{div}_{x_g} E^\perp &= \text{Tr}(J (\nabla E)) = \text{Tr}((\nabla E) J) = 0, \end{aligned}$$

since E is a gradient. At this stage, we can take the limit and obtain that \bar{f} also satisfies equation (1.14).

This is not yet a proper equation in the sense of distributions, even though f depends on $|v_{g,\perp}|$. Indeed, the electric field still exhibits the dependence on the particle position $E \equiv E(t, x_g - v_g^\perp)$. To obtain an equation only depending on $|v_{g,\perp}|$, we use polar coordinates for $v_{g,\perp}$, *i.e.* $v_g = (\rho_L \cos \varphi_c, \rho_L \sin \varphi_c, v_\parallel)$ and Fubini's theorem to integrate the previous integral first in φ_c , then in the other variables. This leads one to the following equation:

$$\begin{aligned} \int_{x_g, v_{g,\parallel}, r} (2\pi \rho_L) \bar{f} (\partial_t \phi + v_{g,\parallel} \partial_{x_{g,\parallel}} \phi + J_{\rho_L}^0 E_\parallel \partial_{v_{g,\parallel}} \phi + (J_{\rho_L}^0 E)^\perp \cdot \nabla_{x_g} \phi \\ + R(t, x_g, v_{g,\parallel}, \rho_L) \cdot \nabla_{\rho_L} \phi) dx_g dv_{g,\parallel} d\rho_L = 0, \end{aligned} \quad (1.15)$$

where the the operator $J_{\rho_L}^0$ is defined in (1.8), and the term R is equal to

$$R(t, x_g, v_{g,\parallel}, \rho_L) = \int_0^{2\pi} E_\perp(x_g - \rho_L e^{i(\varphi_c + \frac{\pi}{2})}) \cdot e^{i\varphi_c} d\varphi_c,$$

with the previously introduced notation $e^{i\varphi_c} = (\cos \varphi_c, \sin \varphi_c, 0)$. It can be shown that R is null since it is the circulation of the electric field E along a gyrocircle. The latter vanishes in the electrostatic case since E is a gradient. With $R = 0$, (1.15) is identical to the equation (1.10) written distribution wise for $2\pi \rho_L \bar{f}$. As there is no dynamics in the ρ_L direction, the factor ρ_L is only a multiplicative constant. One can then introduce $\Phi / \sqrt{\rho_L^2 + \eta}$ as test function, and then obtain that $\bar{f} \rho_L / \sqrt{\rho_L^2 + \eta}$ satisfies the equation. Letting $\eta \rightarrow 0$ in that linear equation, we obtain the appropriate result for the equation governing the evolution of \bar{f} .

Regarding the initial conditions we use a similar projection technique with a test function Φ depending on $|v_{g,\parallel}|$, but not vanishing at $t = 0$. We then obtain equation (1.14), with the right-hand side replaced by

$$- \int \bar{f}_\varepsilon^0 \Phi(0, x_g, v_{g,\parallel}, \rho_L) dx_g dv_{g,\parallel} \rho_L d\rho_L d\varphi_c. \quad (1.16)$$

The asymptotic limit then leads one to:

$$\int \frac{1}{2\pi} \left(\int_0^{2\pi} \bar{f}^0(x_g, \rho_L e^{i\theta} + v_{\parallel} e_{\parallel}) d\varphi_c \right) \Phi(0, \cdot) dx_g dv_{g,\parallel} 2\pi \rho_L d\rho_L. \quad (1.17)$$

Changing coordinate to recover the *position-velocity* coordinates in the φ_c integral, we exactly obtain the initial condition $\tilde{J}_{\rho_L} f^0$ that is expected for \bar{f} . ■

This proof is valid for external electric and magnetic fields. The case where the equation (1.13), invariant in the parallel direction, is coupled to the Poisson equation given a Debye length of the same order as the Larmor radius has been treated by Frénod and Sonnendrücker in [FS01]. It corresponds to a Debye length of order $\sqrt{\varepsilon}$. Technically, we may handle that case with our technic to obtain weak solution as in Arsenev's work [Ars75]. The main point is to obtain L^p estimates on the density $n_i = \int f dv$. They can be obtained by classical estimates using upper bounds for the kinetic energy. The case where the Debye length is taken much smaller than the Larmor radius will be of greater interest, but the difficult problem is there to average the strong oscillations appearing at the scale of the Debye length. We refer to [Gre95], [Gre96] and [CG00] for more details on quasi-neutral plasma without magnetic field, and to [GSR03] for results in the guiding-center approximation with $\rho_L \propto \varepsilon$, $\lambda_D \propto \varepsilon$ and $\varepsilon \rightarrow 0$.

1.2 The electroneutrality equation. Heuristic approach.

In this section we address the self-consistent problem when linking the electric field to the charge distribution. The relevant equation is the Maxwell-Gauss equation that relates the divergence of the electric field to the local charge governed by the particle density of charged particles. The latter must be determined using the ion distribution function for the guiding centers that is solution of the gyrokinetic Vlasov equation. A similar treatment for the electrons must be done. We will follow heuristic arguments together with assumptions that are not justified rigorously. However, this heuristic derivation bares some interest. It is an alternative to the physicists' presentation of that equation based on the Fourier transform. Furthermore, it allows one to recover the electroneutrality equation (1.25) used in the GYSELA code to close the gyrokinetic equation (1.10).

For quasineutral plasmas, the electric field response to any charge separation governs a restoring force. Should the charge separation extend on a scale larger than the Debye length, the restoring force would be too strong to allow any significant charge build-up. The plasma can thus be considered to be everywhere with near zero charge, hence quasineutral. As a consequence, the density of negative charges en_e (n_e being the density of electrons) equals the density of positive charges en_i (n_i being the density of ions). In the electrostatic limit, neglecting the time dependence on the vector potential, one can recover this physics based argument as an asymptotic limit of the Poisson equation for the electric potential.

$$\frac{e}{T_e} \lambda_D^2 \Delta \Phi = \frac{n_e - n_i}{n_e},$$

The right hand side is dimensionless and so is $e\Phi/T_e$. The dimensionless control parameter on the left hand side operator thus appears as the square of the ratio of the Debye scale divided by the characteristic scale of the charge separation. In the limit $\lambda_D^2 \rightarrow 0$, one readily recovers the quasineutrality equation, namely: $n_e = n_i$. Let us consider the ions density. Given the Vlasov equation and its dependence on the electric field, it is likely that the ion distribution function (and thus the ion density) is an implicit function of the electrostatic potential. The same applies for the electron density. Then, in the limit $\lambda_D^2 \rightarrow 0$, there is no analytical dependence on the electric potential. The latter then becomes a Lagrangian multiplier associated to the quasineutrality equation. On such issues, we acknowledge the work of E. Grenier [Gre95], [Gre96] (the only work, at our knowledge, on that subject where the limiting model is kinetic and not fluid), Y. Brenier [BG94], S. Cordier [CG00] and N. Masmoudi [Mas01].

However, if we assume that the system is close to equilibrium, *i.e.* that the departure from a constant electric potential is small, then the electroneutrality equation $n_e = n_i$ provides an explicit dependence on a mean-field electric potential. For an electron-ion plasma, the mass ratio is such that the electrons are

far more mobile than the ions. One can then assume that their response to an electrostatic perturbation is adiabatic on magnetic field lines or surfaces. In that case, assuming that the equilibrium density of electrons $n_{e,0}$ and that of ions $n_{i,0}$ are equal to n_0 , one may write

$$n_e = n_0 e^{\frac{e}{T_e}(\Phi - \langle \Phi \rangle)} \approx n_0 \left(1 + \frac{e}{T_e} (\Phi - \langle \Phi \rangle) \right), \quad (1.18)$$

where $\langle \Phi \rangle$ is the average of Φ on a closed magnetic field line or surface. The expansion on the right hand side holds if $e\Phi \ll T_e$. In the very simple geometry that is considered here (with $\mathbf{B} = (0, 0, B)$), $\langle \Phi \rangle$ is the average in the x_{\parallel} direction: $\langle \Phi \rangle = \int \Phi(x) dx_{\parallel}$. The approximation of adiabatic electrons thus reintroduces an explicit dependence on the electrostatic potential.

For the perturbation of the ion distribution, the first difficulty is to obtain the distribution of ions in physical space from \bar{f} written in gyro-coordinates. If the distribution in physical space is assumed to be constant on gyrocircles as shown in the last section for the limit model, the following formula is obtained,

$$f(t, x, v) = \bar{f}(t, x + v^{\perp}, |v_{\perp}|, v_{\parallel}). \quad (1.19)$$

Taking the integral in v leads to

$$n_i(t, x) = n_0 \int J_{\rho_L}^0(\bar{f}(t, x, \rho_L, v_{\parallel})) 2\pi \rho_L d\rho_L dv_{\parallel}, \quad (1.20)$$

where $J_{\rho_L}^0$ only acts on the x variable (since $\rho_L = |v_{\perp}|$ with our conventions). The quasineutrality equation, $n_e = n_i$, then leads one to

$$1 + \frac{e}{T_e} (\Phi - \langle \Phi \rangle) = \int J_{\rho_L}^0(\bar{f}(t, x, \rho_L, v_{\parallel})) 2\pi \rho_L d\rho_L dv_{\parallel}. \quad (1.21)$$

However, the assumption that the distribution f is constant on the gyrocircles is unrealistic whenever the electric potential is not constant. In order to express the inhomogeneity of the density on gyrocircles, we add a perturbation to \bar{f} . It can be expressed as an adiabatic perturbation on the gyrocircles

$$-\frac{e}{T_i} (\Phi - \bar{\Phi}) n_0 f_i(v), \quad (1.22)$$

$\bar{\Phi}$ is the average of Φ over the gyrocircle of a given particle, and $f_i(v)$ is the equilibrium distribution of ions. Note that this adiabatic perturbation depends on v_{\perp} through the choice of the gyrocircle used in determining the average $\bar{\Phi}$.

$$\begin{aligned} \bar{\Phi}(t, x, v_{\perp}) &= \frac{1}{2\pi} \int \Phi(t, x + v^{\perp} + \rho_L e^{i\varphi_c}) d\varphi_c, \\ n_0 \int \bar{\Phi}(t, x, v_{\perp}) f_i(v) dv &= \frac{n_0}{2\pi} \int \Phi(t, x - \rho_L (e^{i\varphi'_c} + e^{i\varphi_c})) f_i(\rho_L, v_{\parallel}) d\varphi_c d\varphi'_c \rho_L d\rho_L dv_{\parallel} \\ &= n_0 \int (J_{\rho_L}^0)^2 \Phi(t, x) h_i(\rho_L) d\rho_L, \end{aligned} \quad (1.23)$$

where $e^{i\varphi_c} = (\cos \varphi_c, \sin \varphi_c, 0)$ and $h_i(\rho_L) = 2\pi \rho_L \int f_i(\rho_L, v_{\parallel}) dv_{\parallel}$. When the equilibrium distribution is a maxwellian, $f_i(v) = 1/(T_i \pi)^{3/2} e^{-|v|^2/T_i}$, so that $h_i(\rho_L) = 2\rho_L/T_i e^{-\rho_L^2/T_i}$. Adding the adiabatic perturbation of the gyrocircles, we finally obtain the following electroneutrality equation,

$$\begin{aligned} 1 + \frac{e}{T_e} (\Phi - \langle \Phi \rangle) &= \int J_{\rho_L}^0(\bar{f}(t, x, \rho_L, v_{\parallel})) 2\pi \rho_L d\rho_L dv_{\parallel} \\ &\quad - \frac{e}{T_i} \int (1 - (J_{\rho_L}^0)^2) \Phi(t, x) h_i(\rho_L) d\rho_L. \end{aligned} \quad (1.24)$$

Multiplying by T_e/e , then yields

$$\begin{aligned} (\Phi - \langle \Phi \rangle) + \frac{T_e}{T_i} \int (\Phi - (J_{\rho_L}^0)^2 \Phi) h_i(\rho_L) d\rho_L &= \\ \frac{T_e}{e} \left(\int J_{\rho_L}^0(\bar{f}(t, x, \rho_L, v_{\parallel})) 2\pi \rho_L d\rho_L dv_{\parallel} - 1 \right). \end{aligned} \quad (1.25)$$

Remark 1.3 *If the equilibrium density of ions is a Maxwellian, then $h_i(\rho_L) = 2\rho_L/T_i e^{-\rho_L^2/T_i}$ and the operator*

$$\int (J_{\rho_L}^0)^2 h_i(\rho_L) d\rho_L \quad (1.26)$$

is the convolution in the perpendicular plane with the radial function $H_{T_i}(r)$ defined by

$$H_{T_i}(r) = \frac{e^{-\frac{r^2}{4T_i}}}{2\pi^{3/2}\sqrt{T_i}r}. \quad (1.27)$$

See Appendix A for a detailed proof.

Since ρ_L is a parameter in the equation of motion (1.10), the equilibrium distribution and the initial perturbation remain concentrated on a unique value of ρ_L at all time provided it is the case for the initial conditions. A first step to solve the system might be to start under this assumption of single value of ρ_L . This would likely be the most difficult step, since the general case is a superposition of such cases. The gyrokinetic Vlasov equation (1.10) is unchanged when considering a single value of ρ_L , however the electroneutrality equation (1.25) may be simplified and becomes:

$$(\Phi - \langle \Phi \rangle) + \frac{T_e}{T_i} \left(1 - (J_{\rho_L}^0)^2\right) \Phi(t, x) = \frac{T_e}{e} \left(J_{\rho_L}^0 (\bar{n}_i(t, x)) - 1\right), \quad (1.28)$$

where $\bar{n}_i(t, x) = 2\pi\rho_L \int \bar{f}(t, x, v_{\parallel}) dv_{\parallel}$ is the density of ions in gyro-coordinates.

Still, after this further simplification, the gyrokinetic Vlasov equation (1.10) coupled to the electroneutrality equation (1.28) remains a very difficult mathematical problem. The main difficulties are the following.

- The lack of regularity in the parallel direction. The potential Φ has the regularity of f in the parallel direction. This is to be compared to the Vlasov-Poisson case where $D^2\Phi$ ($\Delta\Phi$ in the Poisson equation given above) has the regularity of f . We may overcome this problem by adding some viscosity in that direction, in other words by adding a term $-\lambda\partial_{v_{\parallel}}^2\bar{f}$ in equation (1.10).
- A less important lack of regularity lies in the perpendicular direction. In fact, only the gyro-average of Φ appears in (1.10). Moreover, the term $\langle \Phi \rangle$ is more regular than Φ , so that by (1.28) Φ has the regularity of $J_{\rho_L}^0(\bar{\rho})$. Hence $J_{\rho_L}^0(\Phi)$ has the regularity of $(J_{\rho_L}^0)^2(\bar{\rho})$. The operator $(J_{\rho_L}^0)^2$ sends L^2 into H^1 . That is “almost” enough, in the sense that we could reach a sufficient regularity provided $(J_{\rho_L}^0)^2$ were compact from L^2 in H^1 .

In view of these difficulties, we focus in the next section on the lack of regularity in the parallel direction, and consider time-independent solutions depending only on x_{\parallel} and v , of the form $f(x_{\parallel}, v_{\parallel})f_{\perp}(|v_{\perp}|)$.

2 Steady state solutions in the direction parallel to the magnetic field

Let us consider steady state solutions (f, Φ) to (1.10)-(1.25) and let us assume that the function f can be written in the form $f(x_{\parallel}, v_{\parallel})f_{\perp}(|v_{\perp}|)$, with $\int_0^{+\infty} f_{\perp}(|v_{\perp}|)2\pi|v_{\perp}|d|v_{\perp}| = 1$. Then the term f_i has no incidence in the evolution equation and can thus be factorized in (1.10). Taking $T_e = 1$ for the sake of simplicity, taking again into account the equilibrium density n_0 , not always constant in that section and therefore denoted by n , and replacing n_i by ρ , the electroneutrality equation (1.25) then reads:

$$\Phi - \langle \Phi \rangle = \frac{\rho}{n} - 1,$$

where $\rho(z) = \int f(z, v)dv$. To further simplify the notations, let $z = x_{\parallel}$ and $v = v_{\parallel}$. Solving (1.10)-(1.25) is then equivalent to finding a distribution function f solution of the following equation.

$$v \frac{\partial f}{\partial z} - \left(\frac{\rho}{n}\right)' \frac{\partial f}{\partial v} = 0 \quad (2.1)$$

Note that g' is the derivative of g with respect to z . Furthermore, we consider the problem in a slab geometry $z \in [-1, 1]$, and therefore require given f_{\pm} as boundary conditions,

$$f(-1, v) = f_-(v), \quad v > 0, \quad f(1, v) = f_+(v), \quad v < 0. \quad (2.2)$$

Lemma 2.1 *Given n positive, then any solution f to (2.1)-(2.2), such that $\frac{\rho}{n}$ is non decreasing, must satisfy:*

$$\begin{aligned} \rho(z) = & \int_{-\infty}^0 \frac{|u|f_+(u)}{\sqrt{u^2 + 2\frac{\rho}{n}(1) - 2\frac{\rho}{n}(z)}} du + 2 \int_{\sqrt{2\frac{\rho}{n}(z) - 2\frac{\rho}{n}(-1)}}^{\sqrt{2\frac{\rho}{n}(1) - 2\frac{\rho}{n}(-1)}} \frac{uf_-(u)}{\sqrt{u^2 - 2\frac{\rho}{n}(z) + 2\frac{\rho}{n}(-1)}} du \\ & + \int_{\sqrt{2\frac{\rho}{n}(1) - 2\frac{\rho}{n}(-1)}}^{+\infty} \frac{uf_-(u)}{\sqrt{u^2 - 2\frac{\rho}{n}(z) + 2\frac{\rho}{n}(-1)}} du \end{aligned} \quad (2.3)$$

Proof of Lemma 2.1 The characteristic (Z, V) of (2.1) starting from (z, v) is defined by:

$$\begin{aligned} Z'(s) &= V(s), \quad Z(0) = z, \\ V'(s) &= -\left(\frac{\rho}{n}\right)'(Z(s)), \quad V(0) = v. \end{aligned}$$

Hence

$$V^2(s) + 2\frac{\rho}{n}(Z(s)) = v^2 + 2\frac{\rho}{n}(z).$$

For $v > 0$, it crosses $\{(-1, u), u > 0\}$. For $v < 0$, it crosses $\{(1, u), u < 0\}$ if and only if there is a solution V to $V^2 + 2\frac{\rho}{n}(1) = v^2 + 2\frac{\rho}{n}(z)$, i.e. $v^2 \geq 2\left(\frac{\rho}{n}(1) - \frac{\rho}{n}(z)\right)$. Consequently,

$$\begin{aligned} f(z, v) &= f_-(V(s_-(z, v))) \quad \text{if } v > -\sqrt{2\left(\frac{\rho}{n}(1) - \frac{\rho}{n}(z)\right)}, \\ f(z, v) &= f_+(V(s_+(z, v))) \quad \text{if } v < -\sqrt{2\left(\frac{\rho}{n}(1) - \frac{\rho}{n}(z)\right)}, \end{aligned}$$

where

$$V^2(s_{\pm}(z, v)) + 2\frac{\rho}{n}(\pm 1) = v^2 + 2\frac{\rho}{n}(z), \quad V(s_-(z, v)) > 0, \quad V(s_+(z, v)) < 0.$$

Hence

$$\begin{aligned} f(z, v) &= f_-\left(\sqrt{2\left(\frac{\rho}{n}(z) - \frac{\rho}{n}(-1)\right) + v^2}\right) \quad \text{if } v > -\sqrt{2\left(\frac{\rho}{n}(1) - \frac{\rho}{n}(z)\right)}, \\ f(z, v) &= f_+\left(-\sqrt{2\left(\frac{\rho}{n}(z) - \frac{\rho}{n}(1)\right) + v^2}\right) \quad \text{if } v < -\sqrt{2\left(\frac{\rho}{n}(1) - \frac{\rho}{n}(z)\right)}. \end{aligned}$$

Consequently,

$$\begin{aligned} \rho(z) = & \int_{-\infty}^{-\sqrt{2\left(\frac{\rho}{n}(1) - \frac{\rho}{n}(z)\right)}} f_+\left(-\sqrt{2\left(\frac{\rho}{n}(z) - \frac{\rho}{n}(1)\right) + v^2}\right) dv \\ & + \int_{-\sqrt{2\left(\frac{\rho}{n}(1) - \frac{\rho}{n}(z)\right)}}^{+\infty} f_-\left(\sqrt{2\left(\frac{\rho}{n}(z) - \frac{\rho}{n}(-1)\right) + v^2}\right) dv. \end{aligned}$$

Changes of variables in both integrals lead to (2.3).

For a constant density n , trivial solutions to (2.1)-(2.2) are

$$f(z, v) = f_-(v), \quad v > 0, \quad f(z, v) = f_+(v), \quad v < 0.$$

Proving the existence of solutions to (2.1)-(2.2) satisfying $\left(\frac{\rho}{n}\right)' \geq 0$, for a non constant n , is the aim of this section. Denote by H_i , $1 \leq i \leq 4$, the following set of assumptions.

$$(H_1) \quad n' \leq 0.$$

$$(H_2) \quad \sup_{x \geq 0} \int_{\sqrt{x}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - x}} du =: \lambda < +\infty.$$

$$(H_3) \quad f_-(u) = 0, \quad 0 < u < 2\sqrt{\frac{2\mu}{n(1)}}, \quad \text{where } \mu = \int_{-\infty}^0 f_+(u) du + \lambda.$$

$$(H_4) \quad \int_0^{+\infty} f_-(u) du < \int_{-\infty}^0 f_+(u) du, \quad \int_{-\infty}^0 \frac{f_+(u)}{u^2} du < \frac{n(1)}{4}.$$

Notice that H_2 is satisfied when

$$\lim_{u \rightarrow +\infty} u f_-(u) < +\infty \quad \text{and} \quad \int_0^{+\infty} u |f'_-(u)| du < +\infty.$$

Remark 2.4 *The assumption (H2) ensures that ρ is a perturbation of the equilibrium n , so that with assumption (H1) the force-field will always be oriented rightward (no possibilities of trapped particles). The assumption (H3) ensures that there are two beams of ions (one coming from the right and the other from the left), and that all the ions coming from one side reach the other side (no turn-back). The (H4) assumption is more technical and enforces that the derivative of the density is bounded.*

Lemma 2.2 *Assume H_i , $1 \leq i \leq 4$, and $\frac{n'}{n^2} \in L^\infty$. Denote by*

$$K = \{\alpha \in W^{1,\infty}([-1, 1]); \alpha \geq 0, \alpha(1) - \alpha(-1) \leq \frac{2\mu}{n(1)}, 0 \leq \alpha' \leq 4\mu \|\frac{n'}{n^2}\|_\infty\}. \quad (2.4)$$

There is a solution $\alpha \in K$ to

$$\frac{n}{2}\alpha(z) = \int_{-\infty}^0 \frac{|u| f_+(u)}{\sqrt{u^2 + \alpha(1) - \alpha(z)}} du + \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - \alpha(z) + \alpha(-1)}} du, \quad z \in [-1, 1]. \quad (2.5)$$

Moreover, α is the unique non-decreasing solution of (2.5) such that

$$\alpha(1) - \alpha(-1) \in [0, \frac{2\mu}{n(1)}].$$

Proof of Lemma 2.2. Prove that the map F that maps $\alpha \in K$ in β defined by

$$\begin{aligned} \frac{n}{2}\beta(z) &= \int_{-\infty}^0 \frac{|u| f_+(u)}{\sqrt{u^2 + \alpha(1) - \alpha(z)}} du \\ &+ \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - \alpha(z) + \alpha(-1)}} du, \quad z \in [-1, 1], \end{aligned}$$

has a fixed point. First, F maps K in K . Indeed, $\beta \in W^{1,\infty}([-1, 1])$ like α , β is nonnegative, and

$$\begin{aligned} \frac{n(1)}{2}\beta(1) &= \int_{-\infty}^0 f_+(u) du + \int_{\sqrt{\alpha(1) - \alpha(-1)}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - \alpha(z) + \alpha(-1)}} du \\ &\leq \int_{-\infty}^0 f_+(u) du + \int_{\sqrt{\alpha(1) - \alpha(-1)}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - \alpha(1) + \alpha(-1)}} du \leq \mu. \end{aligned}$$

Hence, $\beta(1) - \beta(-1) \leq \frac{2\mu}{n(1)}$. Moreover,

$$\beta' = 2 \frac{|n'|}{n^2} X + \frac{\alpha'}{n} Y,$$

where

$$\begin{aligned} X &= \int_{-\infty}^0 \frac{|u| f_+(u)}{\sqrt{u^2 + \alpha(1) - \alpha(z)}} du + \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - \alpha(z) + \alpha(-1)}} du, \\ Y &= \int_{-\infty}^0 \frac{|u| f_+(u)}{(u^2 + \alpha(1) - \alpha(z))^{\frac{3}{2}}} du + \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{u f_-(u)}{(u^2 - \alpha(z) + \alpha(-1))^{\frac{3}{2}}} du. \end{aligned}$$

Since $\beta' \geq 0$, and, $X \leq \int_{-\infty}^0 f_+(u) du + \lambda = \mu$, then,

$$\sup_{x \geq 0} \int_{2\sqrt{x}}^{+\infty} \frac{u}{(u^2 - x)^{\frac{3}{2}}} f_-(u) du \leq \frac{n(1)}{3\sqrt{3}\mu} \int_0^{+\infty} f_-(u) du.$$

Indeed, either $x \geq \frac{2\mu}{n(1)}$ and then

$$u > 2\sqrt{x} \Rightarrow \frac{u}{(u^2 - x)^{\frac{3}{2}}} \leq \frac{2}{3\sqrt{3}x} \leq \frac{n(1)}{3\sqrt{3}\mu},$$

or $x < \frac{2\mu}{n(1)}$ and then $f_-(u) \neq 0$ implies that $u > 2\sqrt{\frac{2\mu}{n(1)}}$, hence

$$\frac{u}{(u^2 - x)^{\frac{3}{2}}} \leq \frac{2\sqrt{\frac{2\mu}{n(1)}}}{\left(\frac{8\mu}{n(1)} - x\right)^{\frac{3}{2}}} \leq \frac{n(1)}{3\sqrt{3}\mu}.$$

So that,

$$\sup_{x \geq 0} \int_{2\sqrt{x}}^{+\infty} \frac{u}{(u^2 - x)^{\frac{3}{2}}} f_-(u) du \leq \frac{n(1)}{3\sqrt{3} \int_{-\infty}^0 f_+(u) du} < \frac{n(1)}{4},$$

and by H_4 ,

$$Y \leq \int_{-\infty}^0 \frac{f_+(u)}{u^2} du + \frac{n(1)}{4} \leq \frac{n(1)}{2}.$$

Consequently, $\beta' \leq 4\mu \|\frac{n'}{n^2}\|_{\infty}$. And so, F maps K in K . Moreover, F is continuous for the topology of $C([-1, 1])$, by definition of β in terms of α . Finally, F is compact for the topology of $C([-1, 1])$, by the compact embedding of $W^{1,\infty}([-1, 1])$ in $C([-1, 1])$ and the boundedness of K in $W^{1,\infty}([-1, 1])$. It follows from a Schauder fixed point theorem that there is a fixed point for F in K .

Moreover, the solution of (2.5) is unique in the class of non-decreasing functions α such that $\alpha(1) - \alpha(-1) \in [0, \frac{2\mu}{n(1)}]$. Indeed, for any solution α of (2.5) in this class, $x := \alpha(1) - \alpha(-1)$ solves $G(x) = 0$, where

$$\begin{aligned} G(x) &:= x - \frac{2}{n(1)} \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - x}} du + \frac{2}{n(-1)} \int_{-\infty}^0 \frac{|u| f_+(u)}{\sqrt{u^2 + x}} du \\ &\quad - \frac{2}{n(1)} \int_{-\infty}^0 f_+(u) du + \frac{2}{n(-1)} \int_0^{+\infty} f_-(u) du. \end{aligned}$$

Using estimates very similar to (2.6) and (H_4) , we can show the function G is increasing. Moreover it follows from H_3 and H_4 that $G(0) \leq 0$ and $G(\frac{2\mu}{n(1)}) \geq 0$. Hence the value of $\alpha(1) - \alpha(-1)$ is unique, as well as $(\alpha(1), \alpha(-1))$, given by

$$\begin{aligned} \alpha(1) &= \frac{2}{n(1)} \left(\int_{-\infty}^0 f_+(u) du + \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{u f_-(u)}{\sqrt{u^2 - \alpha(1) + \alpha(-1)}} du \right), \\ \alpha(-1) &= \frac{2}{n(-1)} \left(\int_{-\infty}^0 \frac{|u| f_+(u)}{\sqrt{u^2 + \alpha(1) - \alpha(-1)}} du + \int_0^{+\infty} f_-(u) du \right). \end{aligned}$$

If α and β are two non-decreasing solutions of (2.5), then

$$(\alpha - \beta)(z)T(z) = 0, \quad z \in [-1, 1],$$

where

$$\begin{aligned} T(z) &= \frac{n(z)}{2} \\ &- \int_{-\infty}^0 \frac{|u|f_+(u)}{\sqrt{u^2 - \alpha(z) + \alpha(1)}\sqrt{u^2 - \beta(z) + \alpha(1)}(\sqrt{u^2 - \alpha(z) + \alpha(1)} + \sqrt{u^2 - \beta(z) + \alpha(1)})} du \\ &- \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{uf_-(u)}{\sqrt{u^2 - \alpha(z) + \alpha(-1)}\sqrt{u^2 - \beta(z) + \alpha(-1)}(\sqrt{u^2 - \alpha(z) + \alpha(-1)} + \sqrt{u^2 - \beta(z) + \alpha(-1)})} du. \end{aligned}$$

By H_1 , H_3 and H_4 ,

$$\begin{aligned} 2T(z) &\geq n(1) - \int_{-\infty}^0 \frac{f_+(u)}{u^2} du - \int_{2\sqrt{\frac{2\mu}{n(1)}}}^{+\infty} \frac{uf_-(u)}{(u^2 - \frac{2\mu}{n(1)})^{\frac{3}{2}}} du \\ &\geq n(1) - \int_{-\infty}^0 \frac{f_+(u)}{u^2} du - \frac{n(1)}{3\sqrt{3}\mu} \int_0^{+\infty} f_-(u) du > 0, \quad z \in [-1, 1]. \end{aligned}$$

Hence $\alpha = \beta$.

Theorem 2.3 Assume H_i , $1 \leq i \leq 4$, and $f_{\pm} \in L^{\infty}$.

There is a unique solution $f \in L^{\infty}([-1, 1] \times \mathbb{R})$ to (2.1)-(2.2), such that $\frac{\rho}{n} \in K$.

Proof of Theorem 2.3 Let $\alpha \in K$ be the solution to (2.5). The distribution function f defined by

$$\begin{aligned} f(z, v) &= f_-(\sqrt{\alpha(z) - \alpha(-1) + v^2}), \quad v > -\sqrt{\alpha(1) - \alpha(z)}, \\ f(z, v) &= f_+(\sqrt{\alpha(z) - \alpha(1) + v^2}), \quad v < -\sqrt{\alpha(1) - \alpha(z)}, \end{aligned}$$

is the unique solution to (2.1)-(2.2) in $L^{\infty}([-1, 1] \times \mathbb{R})$ such that $\frac{\rho}{n} \in K$.

A Appendix: The polarization operator as a convolution

In this section we restate and prove the result announced in Remark 1.3.

We define

$$F_T^0 = \frac{2}{T} \int_0^{+\infty} \rho e^{-\rho^2/T} (J_{\rho}^0)^2 d\rho, \quad (\text{A.6})$$

where T is the temperature (of the ions) and J_{ρ}^0 is the gyro-average operator defined in (1.8). We forget the subscript L in the Larmor radius for convenience. We use here the measure $(2\rho/T)e^{-\rho^2/T}$ and not an usual Maxwellian, because we start from a 2D Maxwellian and perform an integration over the angular variable in polar coordinates.

The operator F_0 appears in the electroneutrality equation (1.25). Here we will prove the following proposition:

Proposition A.1 The operator F_T^0 is the convolution with the radial function H_T , defined by

$$H_T(r) = \frac{e^{-\frac{r^2}{4T}}}{2\pi^{3/2}r\sqrt{T}}. \quad (\text{A.7})$$

Proof of the proposition. First, the square of the operator J_{ρ}^0 is

$$(J_{\rho}^0)^2(f)(x_g) = \frac{1}{4\pi^2} \int_0^{2\pi} \int_0^{2\pi} f(x_g + \rho e^{i\varphi_c} + \rho e^{i\varphi'_c}) d\varphi_c d\varphi'_c.$$

To simplify it, we use the equality

$$\rho(e^{i\varphi_c} + e^{i\varphi'_c}) = 2\rho \cos\left(\frac{\varphi_c - \varphi'_c}{2}\right) e^{i\frac{\varphi_c + \varphi'_c}{2}},$$

which helps defining the polar coordinates (r, ϕ)

$$\begin{cases} r = 2\rho \left| \cos\left(\frac{\varphi_c - \varphi'_c}{2}\right) \right|, \\ \phi = \frac{\varphi_c + \varphi'_c}{2} + \varepsilon, \end{cases} \quad (\text{A.8})$$

where $\varepsilon = 0$ or π depending on the sign of the cosine. This is not exactly a change of variable since a couple (r, ϕ) has exactly two pre-images (φ_c, φ'_c) and (φ'_c, φ_c) . But since it is always bi-valued, we can use the formula for the change of variables with a factor 2. The presence of ε does not introduce specific difficulties, if the intervals of integration are distinguished. The Jacobian of this transformation is

$$\rho \left| \sin\left(\frac{\varphi_c - \varphi'_c}{2}\right) \right| = \sqrt{\rho^2 - \frac{r^2}{4}},$$

so that

$$(J_\rho^0)^2(f)(x_g) = \frac{1}{\pi^2} \int_0^{2\rho} \int_0^{2\pi} f(x_g + r e^{i\phi}) \frac{d\phi dr}{\sqrt{4\rho^2 - r^2}},$$

The next step is to introduce cartesian coordinates. This is a standard transform with Jacobian r , so that

$$(J_\rho^0)^2(f)(x_g) = \int_{B(2\rho)} f(x_g + y) \frac{dy}{\pi^2 r \sqrt{4\rho^2 - r^2}},$$

with $y \in \mathbb{R}^2$, $r = |y|$ and the notation $B(a)$ for the ball of center 0 and radius a . This is exactly the convolution with the radial function

$$h_\rho(r) = \frac{\chi_{(0,2\rho)}(r)}{\pi^2 r \sqrt{4\rho^2 - r^2}},$$

where χ_A denote the characteristic function of A . It can be checked that

$$\int h_\rho(r) 2\pi r dr = 1.$$

The last step is to perform the integration in ρ . As $(J_\rho^0)^2$ is the convolution with the radial function h_ρ , $F_T^0 = \frac{2}{T} \int \rho e^{-\rho^2/T} (J_\rho^0)^2 d\rho$ is the convolution with the radial function

$$H_T(r) = \frac{2}{T} \int_0^{+\infty} h_\rho(r) \rho e^{-\rho^2/T} d\rho.$$

In other words,

$$H_T(r) = \frac{2}{\pi^2 r T} \int_{r/2}^{+\infty} \frac{\rho e^{-\rho^2/T}}{\sqrt{4\rho^2 - r^2}} d\rho.$$

We perform the change of variable $\rho' = \sqrt{4\rho^2 - r^2}$. Hence

$$\begin{aligned} H_T(r) &= \frac{e^{-\frac{r^2}{4T}}}{2\pi^2 r T} \int_0^{+\infty} e^{-\frac{\rho'^2}{4T}} d\rho' \\ &= \frac{e^{-\frac{r^2}{4T}}}{2\pi^{3/2} r \sqrt{T}}. \end{aligned}$$

It can be checked that H_T has total mass one. Indeed,

$$\int_0^\infty H_T(r) 2\pi r dr = \frac{1}{\sqrt{\pi T}} \int_0^\infty e^{-\frac{r^2}{4T}} dr = 1$$

■

References

- [Amb04] Luigi Ambrosio. Transport equation and Cauchy problem for BV vector fields. *Invent. Math.*, 158(2):227–260, 2004.
- [Ars75] A. A. Arsen'ev. Existence in the large of a weak solution of Vlasov's system of equations. *Ž. Vychisl. Mat. i Mat. Fiz.*, 15:136–147, 276, 1975.
- [BG94] Yann Brenier and Emmanuel Grenier. Limite singulière du système de Vlasov-Poisson dans le régime de quasi neutralité: le cas indépendant du temps. *C. R. Acad. Sci. Paris Sér. I Math.*, 318(2):121–124, 1994.
- [BH07] A. J. Brizard and T. S. Hahm. Foundations of nonlinear gyrokinetic theory. *Reviews of Modern Physics*, 79:421–468, April 2007.
- [Bos09] Mihai Bostan. The Vlasov-Poisson system with strong external magnetic field. Finite Larmor radius regime. *Asymptot. Anal.*, 61(2):91–123, 2009.
- [Bre00] Y. Brenier. Convergence of the Vlasov-Poisson system to the incompressible Euler equations. *Comm. Partial Differential Equations*, 25(3-4):737–754, 2000.
- [Cat78] P. J. Catto. Linearized gyro-kinetics. *Plasma Physics*, 20:719–722, July 1978.
- [CG00] Stéphane Cordier and Emmanuel Grenier. Quasineutral limit of an Euler-Poisson system arising from plasma physics. *Comm. Partial Differential Equations*, 25(5-6):1099–1113, 2000.
- [CTB81] P. J. Catto, W. M. Tang, and D. E. Baldwin. Generalized gyrokinetics. *Plasma Physics*, 23:639–650, July 1981.
- [DL89a] R. J. DiPerna and P.-L. Lions. Global weak solutions of Vlasov-Maxwell systems. *Comm. Pure Appl. Math.*, 42(6):729–757, 1989.
- [DL89b] R. J. DiPerna and P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.*, 98(3):511–547, 1989.
- [FS00a] E. Frénod and E. Sonnendrücker. Long time behavior of the two-dimensional Vlasov equation with a strong external magnetic field. *Math. Models Methods Appl. Sci.*, 10(4):539–553, 2000.
- [FS00b] Emmanuel Frénod and Eric Sonnendrücker. Approximation “rayon de Larmor fini” pour l'équation de Vlasov. *C. R. Acad. Sci. Paris Sér. I Math.*, 330(5):421–426, 2000.
- [FS01] Emmanuel Frénod and Eric Sonnendrücker. The finite Larmor radius approximation. *SIAM J. Math. Anal.*, 32(6):1227–1247, 2001.
- [GIVT09] X. Garbet, Y. Idomura, L. Villard, and Watanabe T. Gyrokinetic simulations of turbulent transport. *Nuclear Fusion*, 2009. In Preparation.
- [GLV08] P. Ghendrih, R. Lima, and R. Vilela Mendes. Reduction and approximation in guiding-center dynamics. *Journal of Physics A Mathematical General*, 41:465501, November 2008.
- [Gre95] E. Grenier. Defect measures of the Vlasov-Poisson system in the quasineutral regime. *Comm. Partial Differential Equations*, 20(7-8):1189–1215, 1995.
- [Gre96] Emmanuel Grenier. Oscillations in quasineutral plasmas. *Comm. Partial Differential Equations*, 21(3-4):363–394, 1996.
- [Gre97] E. Grenier. Pseudo-differential energy estimates of singular perturbations. *Comm. Pure Appl. Math.*, 50(9):821–865, 1997.
- [GSG+06a] V. Grandgirard, Y. Sarazin, X. Garbet, G. Dif-Pradalier, P. Ghendrih, N. Crouseilles, G. Latu, E. Sonnendrücker, N. Besse, and P. Bertrand. Computing ITG turbulence with a full-f semi-Lagrangian code. 2006.

- [GSG⁺06b] V. Grandgirard, Y. Sarazin, X. Garbet, G. Dif-Pradalier, P. Ghendrih, N. Crouseilles, G. Latu, E. Sonnendrücker, N. Besse, and P. Bertrand. GYSELA, a full-f global gyrokinetic Semi-Lagrangian code for ITG turbulence simulations. In O. Sauter, editor, *Theory of Fusion Plasmas*, volume 871 of *American Institute of Physics Conference Series*, pages 100–111, November 2006.
- [GSR03] F. Golse and L. Saint-Raymond. The Vlasov-Poisson system with strong magnetic field in quasineutral regime. *Math. Models Methods Appl. Sci.*, 13(5):661–714, 2003.
- [Hah88] T. S. Hahm. Nonlinear gyrokinetic equations for tokamak microturbulence. *Physics of Fluids*, 31:2670–2673, September 1988.
- [Hor93] E. Horst. On the asymptotic growth of the solutions of the Vlasov-Poisson system. *Math. Methods Appl. Sci.*, 16(2):75–86, 1993.
- [III99] ITER Physics Expert Groups on Confinement and Transport and Confinement Modelling and Database, ITER Physics Basis Editors, and ITER EDA. Chapter 2: Plasma confinement and transport. *Nuclear Fusion*, 49(12):2179–2249, December 1999.
- [LEHT02] Z. Lin, S. Ethier, T. S. Hahm, and W. M. Tang. Size Scaling of Turbulent Transport in Magnetically Confined Plasmas. *Physical Review Letters*, 88(19):195004–+, May 2002.
- [LP91] P.-L. Lions and B. Perthame. Propagation of moments and regularity for the 3-dimensional Vlasov-Poisson system. *Invent. Math.*, 105(2):415–430, 1991.
- [Mas01] Nader Masmoudi. From Vlasov-Poisson system to the incompressible Euler system. *Comm. Partial Differential Equations*, 26(9-10):1913–1928, 2001.
- [Nic83] D. R. Nicholson. *Introduction to plasma theory*. Wiley series in plasma physics. John Wiley & Sons, New York, 1983.