

HLA-DRB1 Genotypes and the Risk of Developing Anti Citrullinated Protein Antibody (ACPA) Positive Rheumatoid Arthritis

Nathalie Balandraud^{1,2}, Christophe Picard³, Denis Reviron³, Cyril Landais⁶, Eric Toussiroot^{4,5}, Nathalie Lambert², Emmanuel Telle², Caroline Charpin^{1,2}, Daniel Wendling⁵, Etienne Pardoux⁶, Isabelle Auger², Jean Roudier^{1,2*}

1 Rhumatologie, Hôpital Sainte Marguerite, Assistance Publique Hôpitaux de Marseille, Marseille, France, **2** Institut National des Sciences et de la Recherche Médicale, Unité Mixte de Recherche en Santé 1097, Aix Marseille Université, Marseille, France, **3** Laboratoire d'Immunogénétique et d'Histocompatibilité, Etablissement Français du Sang Alpes-Méditerranée, Marseille, France, **4** Centre d'Investigation Clinique 506, Hôpital Saint Jacques, Besançon, France, **5** Rhumatologie, Hôpital Minjoz, Besançon, France, **6** Centre National pour la Recherche Scientifique, Unité Mixte de Recherche 6632, Équipe Évolution Biologique et Modélisation, Aix Marseille Université, Marseille, France

Abstract

Objective: To provide a table indicating the risk for developing anti citrullinated protein antibody (ACPA) positive rheumatoid arthritis (RA) according to one's HLA-DRB1 genotype.

Methods: We HLA-DRB1 genotyped 857 patients with ACPA positive RA and 2178 controls from South Eastern and Eastern France and calculated Odds Ratios (OR) for developing RA for 106 of 132 possible genotypes accounting for 97% of subjects.

Results: HLA-DRB1 genotypic ORs for developing ACPA positive RA range from 28 to 0.19. HLA-DRB1 genotypes with HLA-DRB1*04SE (HLA-DRB1*0404, HLA-DRB1*0405, HLA-DRB1*0408), HLA-DRB1*04:01, HLA-DRB1*01 are usually associated with high risk for developing RA. The second HLA-DRB1 allele in genotype somewhat modulates shared epitope associated risk. We did not identify any absolutely protective allele. Neither the Reviron, nor the du Montcel models accurately explains our data which are compatible with the shared epitope hypothesis and suggest a dosage effect among shared epitope positive HLA-DRB1 alleles, double dose genotypes carrying higher ORs than single dose genotypes.

Conclusion: HLA-DRB1 genotypic risk for developing ACPA positive RA is influenced by both HLA-DRB1 alleles in genotype. We provide an HLA-DRB1 genotypic risk table for ACPA positive RA.

Citation: Balandraud N, Picard C, Reviron D, Landais C, Toussiroot E, et al. (2013) HLA-DRB1 Genotypes and the Risk of Developing Anti Citrullinated Protein Antibody (ACPA) Positive Rheumatoid Arthritis. PLoS ONE 8(5): e64108. doi:10.1371/journal.pone.0064108

Editor: Sunil K. Ahuja, South Texas Veterans Health Care System and University Health Science Center San Antonio, United States of America

Received: November 26, 2012; **Accepted:** April 9, 2013; **Published:** May 30, 2013

Copyright: © 2013 Balandraud et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was funded by INSERM (Institut National des Sciences et de la Recherche Médicale), Arthritis Fondation Clarins and Société Française de Rhumatologie. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interest exist.

* E-mail: jean.roudier@inserm.fr

Introduction

Susceptibility for developing rheumatoid arthritis (RA) is associated with particular HLA-DRB1 alleles like HLA-DRB1*04, HLA-DRB1*01 and HLA-DRB1*10 [1]. A molecular basis for this association was provided by Gregersen and al. who showed that RA associated HLA-DRB1 alleles contain a conserved 5 amino acid stretch, the “shared epitope” (SE) in the third hypervariable region of their DRB1 chain [2]. This led to a simple model in which shared epitope positive HLA-DRB1 alleles carried susceptibility for developing RA and shared epitope negative alleles were considered neutral.

Since 1987, numerous studies have confirmed the association of RA with shared epitope positive HLA-DRB1 alleles. However, it has been suggested that a few shared epitope negative HLA-DRB1 alleles protect against the development of RA, whereas most others are neutral [3–5].

Different models have been proposed to predict whether a given genotype (ie both genes present in an individual) will be susceptible, neutral or protective towards the development of RA. These models are based on the classification of HLA-DRB1 alleles in different categories depending on the sequence of their third hypervariable region [3–5]. The Reviron model proposes that shared epitope positive HLA-DRB1 alleles which have a very positive charge in their P4 pocket (HLA-DRB1*0401, HLA-DRB1*0404, HLA-DRB1*0405, HLA-DRB1*0408, HLA-DRB1*0101) predispose to RA whereas among shared epitope negative alleles, those which have a positive charge in their P4 pocket, called XP4p (HLA-DRB1*03, HLA-DRB1*09, HLA-DRB1*14, HLA-DRB1*15, HLA-DRB1*16), are neutral and those with a negative or neutral charge in their P4 pocket, called XP4n (HLA-DRB1*07, HLA-DRB1*08, HLA-DRB1*11, HLA-DRB1*0402), protect against RA [4]. Sophie Tézenas du Montcel proposed a model which considers 4 groups of HLA-DRB1 alleles

according to the sequence of their third hypervariable region: S2 (QKRAA), S3P (QRRAA, RRRAA), S3D (DRRAA), S1 (ARAA and ERAA) and X (non RAA). In the du Montcel model, S2 alleles confer very high risk, S3P alleles confer high risk and S1, S3D and X alleles confer low risk to develop RA [5]. The main difference between the two models is the absence of protective alleles in the du Montcel model.

However, the validity of these models has not been established. Here, we decided to calculate the risk for developing RA given by any individual HLA-DRB1 genotype, with no a priori model.

Complexity of the RA/HLA-DRB1 association further increased with the discovery that RA can be divided into two subtypes according to the presence in a patient's serum of autoantibodies directed at citrullin residues on different proteins [6]. Indeed, the sera of two thirds of patients with RA contain antibodies to citrullinated proteins. Citrullin is an amino acid generated by the posttranslational modification of arginyl residues by peptidyl arginine deiminases. These autoantibodies are called anti citrullinated protein antibodies (ACPA) [6]. Presence or absence of ACPAs define two subtypes of RA. ACPA positive RA is well defined and the 2010 ACR criteria for the diagnosis of RA include a positive ACPA test [7]. ACPA negative RA is much more heterogeneous [8]. The association of RA with shared epitope positive HLA-DRB1 alleles is stronger in ACPA positive RA than in ACPA negative RA [8,9].

Here, we studied a series of 857 ACPA positive RA patients and compared them with a series of 2178 controls.

Every patient and control, all from South Eastern France was HLA-DR typed for 20 different HLA-DRB1 alleles. Bayesian statistics were used to define susceptible and protective genotypes and to calculate accurate confidence intervals for the associated Odds Ratios. We calculated Odds Ratios (OR) for 102 of 136 possible HLA-DRB1 genotypes, for which the number of patients plus controls was at least 5. Thirty genotypes had ORs significantly higher than 1 and were considered high risk, 45 had ORs not significantly different from 1 and were considered neutral, 27 genotypes had ORs significantly lower than 1 and were considered low risk.

Patients and Methods

Patients and Controls

Ethical approval was obtained for this study from CPP Marseille II. All participants gave their informed consent. All participants signed informed consent according to the Declaration of Helsinki.

In this cohort study, we took blood samples from 857 ACPA positive RA patients and 2178 controls from Marseilles (South Eastern France) and Besançon (Eastern France). RA patients fulfilled the 2010 EULAR/ACR criteria [7] and tested positive for anti cyclic-citrullinated peptide antibodies (ACPA) using the anti CCP2 kit which is routinely used in our patients with arthritis. The control group included voluntary blood or bone marrow donors from the same area. Sex ratio and age were not significantly different between patients and controls. Seventy one percent of patients were positive for at least one shared epitope positive HLA-DRB1 allele (Table 1).

ACPA and Rheumatoid Factor Testing

Positivity for anti citrullinated peptide antibodies (ACPA) was used to define mainstream, classical RA [8]. ACPA were detected by anti CCP2 Enzyme-linked immuno sorbent assay (ELISA) (Immunoscan RA, Euro-Diagnostica, Arnhem, the Netherlands). This kit is currently the most commonly used in Southern France and positivity is defined by a cutoff value of 25 Units/ml at a

Table 1. Patients' and controls' characteristics.

	Patients	Controls
Number	857	2178
Female	74.40%	69%
Male	26%	31%
ACPA	100%	0%
RF pos	91%	NC
SE double dose	20.40%	4.50%
SE single dose	51%	37%
Age at diagnosis	43.6	42

RF: rheumatoid factor, pos: positive, SE: Shared Epitope, ACPA: anti citrullinated protein antibodies, NT: not tested

doi:10.1371/journal.pone.0064108.t001

dilution of 1/50. 91% of our patients selected for ACPA positivity by anti CCP2 testing tested positive for rheumatoid factor by ELISA using the Orgentec Kit (Mainz, Germany) with a 20 Units/ml cutoff at a 1/100 dilution.

HLA-DRB1 Genotyping

Low resolution HLA-DRB1 typing was carried out according to the manufacturer's specification for LABType SSO (One Lambda Inc, USA) and the retrieved output was analyzed by HLA Fusion v 1.2.1. software (One Lambda Inc, USA) for allele identification. HLA-DRB1*04-positive samples were subtyped by PCR sequence-specific primers (Olerup SSP HLA-DRB1*04, Genovision, Vienna, Austria).

We typed for the following alleles or group of alleles: HLA-DRB1*01, HLA-DRB1*03, HLA-DRB1*04:01, HLA-DRB1*04:02, HLA-DRB1*04:03, HLA-DRB1*04:04, HLA-DRB1*04:05, HLA-DRB1*04:06, HLA-DRB1*04:07, HLA-DRB1*04:08, HLA-DRB1*07, HLA-DRB1*08, HLA-DRB1*09, HLA-DRB1*1001, HLA-DRB1*11, HLA-DRB1*12, HLA-DRB1*13, HLA-DRB1*14, HLA-DRB1*15, HLA-DRB1*16. HLA-DRB1*04:04, HLA-DRB1*04:05 and HLA-DRB1*04:08 were considered as one allelic group called HLA-DRB1*04SE. The rationale for this regrouping is that these three HLA-DRB1*04 alleles encode a common amino acid sequence, QRRAA in their third hypervariable region (HV3). This HV3 sequence is different from that encoded by HLA-DRB1*04:01, QKRAA. Therefore, HLA-DRB1*04:01 is not included in this HLA-DRB1*04 allelic subgroup that we called HLA-DRB1*04SE.

After this regrouping, we ended up with 16 allelic groups defining 136 genotypes.

Statistical Analysis

The statistical method used in this study to estimate the Odds Ratio and its confidence interval for the different genotypes is based on the Bayesian theory. This method is more stringent than the one using the maximization of the likelihood function especially when there are less than 5 individuals in one of the two classes of genotypes (patient or control). This method was applied with the Matlab® Software. Data were simulated according to the posteriori density of Bayesian statistics in order to obtain the confidence intervals of the different odds ratios.

Results

HLA-DRB1 Alleles Associated with RA

Seventy one percent of patients and 41% of controls were positive for the HLA-DRB1 shared epitope. Fifty one percent of patients and 37% of controls expressed one shared epitope positive allele. Twenty percent of patients and 4.5% of controls expressed two shared epitope positive HLA-DRB1 alleles.

RA associated HLA-DRB1 alleles were, by order of Odds Ratios, HLA-DRB1*04:08 (OR = 10.3), HLA-DRB1*04:01 (OR = 3.3), HLA-DRB1*04:05 (OR = 3.3), HLA-DRB1*10 (OR = 2.9), HLA-DRB1*04:04 (OR = 2.8), HLA-DRB1*09 (OR = 2.1), HLA-DRB1*01 (OR = 1.5) (Table 2). For genotype risk analysis, HLA-DRB1*04:04, HLA-DRB1*04:05, HLA-DRB1*04:08 were pooled into the HLA-DRB1*04SE group.

Alleles significantly underrepresented in RA patients (thus potentially protective) were HLA-DRB1*08 (OR = 0.42), HLA-DRB1*13 (OR = 0.58), HLA-DRB1*03 (OR = 0.51), HLA-DRB1*11 (OR = 0.54), HLA-DRB1*14 (OR = 0.58), HLA-DRB1*07 (OR = 0.66) (Table 2).

HLA-DRB1 Genotypes and the Risk for Developing ACPA Positive RA

Sixteen HLA-DRB1 allelic groups define 136 genotypes. We arbitrarily decided not to calculate Odds Ratios for genotypes for which the total number of subjects (patients plus controls) was less than 5. This allowed us calculate Odds Ratios for 102 of 136 possible genotypes (Figure 1). Genotypic Odds Ratios ranged from 28 (HLA-DRB1*04:01/*10) to 0.19 (HLA-DRB1*03/*03)

(Figure 1). Thirty genotypes (29%) had ORs significantly higher than 1 and were considered “high risk”, 45 (44%) had ORs not significantly different from 1 and were considered neutral, 27 (26%) had ORs significantly lower than 1 and were considered “low risk” (Figure 2).

High Risk and Low Risk Genotypes

Of 30 “high risk” genotypes, 18 contain one and 10 contain two RA associated HLA-DRB1 alleles. Only two: HLA-DRB1*07/*16 and HLA-DRB1*07/*04:02 do not contain any RA associated allele (Figure 2).

Of 27 “low risk” genotypes, 26 contain no RA associated allele. Only HLA-DRB1*01/*13 contains RA associated HLA-DRB1*01 (Figure 2).

Risk carried by RA associated allele is modulated by second allele in genotype as observed in genotypes containing HLA-DRB1*04SE, HLA-DRB1*04:01, HLA-DRB1*01. For instance, ORs associated with HLA-DRB1*04SE genotypes range from 13.4 (HLA-DRB1*04SE/04:01) to 1 (HLA-DRB1*04SE/08) (Figure 3).

Discussion

RA Associated and Non RA Associated HLA-DRB1 Alleles

In this study of HLA-DRB1 alleles expressed in 857 patients with ACPA positive RA and 2178 normal controls from South Eastern France, we confirm the classical association of RA with shared epitope positive HLA-DRB1 alleles.

Seven HLA-DRB1 alleles are associated with RA in our population, HLA-DRB1*04:01, HLA-DRB1*04:04, HLA-

Table 2. HLA-DRB1 allelic frequencies in patients and controls.

	Patient number n = 1714	Frequency (%)	Control number n = 4356	Frequency (%)	Odds ratio	P-value (FISHER TEST)	Confidence interval
DRB1*01:XX	263	15.34	495	11.36	1.52*	6.6×10 ⁻⁷	1.28–1.79
DRB1*03:XX	101	5.89	470	10.79	0.51*	1.1×10 ⁻⁹	0.41–0.64
DRB1*04:01	246	14.35	208	4.78	3.31*	5.1×10 ⁻³³	2.71–4.04
DRB1*04:02	18	1.05	38	0.87	1.2	0.55	0.64–2.17
DRB1*04:03	18	1.05	63	1.45	0.7	0.31	0.41–1.27
DRB1*04:04	120	7.00	114	2.62	2.80*	3.6×10 ⁻¹⁴	2.13–3.67
DRB1*04:05	69	4.03	55	1.26	3.27*	1.12×10 ⁻¹⁰	2.26–4.78
DRB1*04:06	1	0.06	8	0.18	0.31	0.46	0.007–2.37
DRB1*04:07	10	0.58	41	0.94	0.55	0.11	0.23–1.16
DRB1*04:08	32	1.87	8	0.18	10.33*	1.34×10 ⁻¹¹	4.70–26.0
DRB1*07:XX	180	10.50	654	15.01	0.66*	2.31×10 ⁻⁶	0.55–0.79
DRB1*08:XX	21	1.23	123	2.82	0.42*	0.0001	0.25–0.68
DRB1*09:XX	30	1.75	37	0.85	2.07*	0.0038	1.24–3.47
DRB1*10:XX	56	3.27	50	1.15	2.9*	8.5×10 ⁻⁸	1.94–4.36
DRB1*11:XX	142	8.28	626	14.37	0.54*	3.3×10 ⁻¹¹	0.44–0.65
DRB1*12:XX	21	1.23	71	1.63	0.7	0.29	0.43–1.23
DRB1*13:XX	130	7.58	542	12.44	0.58*	2.63×10 ⁻⁸	0.47–0.70
DRB1*14:XX ¹	39	2.28	169	3.88	0.58*	0.0016	0.39–0.82
DRB1*15:XX	161	9.39	465	10.67	0.87	0.14	0.7–1.05
DRB1*16:XX	56	3.27	119	2.73	1.20	0.26	0.85–1.67

*statistically significant.

ORs and p values by Fisher's test.

doi:10.1371/journal.pone.0064108.t002






Figure 3. Influence of second allele on genotypic ORs in genotypes with HLA-DRB1*04SE, HLA-DRB1*04:01, HLA-DRB1*01. Genotypes with HLA-DRB1*04SE and HLA-DRB1*04:01 keep high ORs despite modulation by second allele. doi:10.1371/journal.pone.0064108.g003

prove useful for genetic counseling in families affected by rheumatoid arthritis, indicating which individual to monitor for emergence of ACPA response and possibly early treatment or even prevention.

Finally, our genotypic risk table does not necessarily reflect the effect and interaction of HLA-DRB1 genes only, but may also involve non HLA-DRB1 genes on HLA-DRB1 haplotypes.

The Remaining Third: ACPA Negative RA

This study did not approach the very controversial issue of ACPA negative RA and its HLA-DRB1 association(s). Indeed, ACPA negative RA is somewhat heterogeneous and careful analysis of patients' files often ends up with diagnosis other than RA. Still, one large scale study of HLA-DRB1 associations in ACPA negative RA patients suggests that half of them test positive for rheumatoid factors and show HLA-DRB1 association similar to ACPA positive RA [11]. Such patients will be the objects of future studies.

References

- Ollier W, Thomson W (1992) Population genetics of rheumatoid arthritis. *Rheum Dis Clin North Am* 18: 741–759.
- Gregersen PK, Silver J, Winchester RJ (1987) The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum* 30: 1205–1213.
- Mattey DL, Dawes PT, Gonzalez-Gay MA, Garcia-Porrúa C, Thomson W, et al. (2001) HLA-DRB1 alleles encoding an aspartic acid at position 70 protect against development of rheumatoid arthritis. *J Rheumatol* 28: 232–239.
- Reviron D, Perdriger A, Toussiro E, Wendling D, Balandraud N, et al. (2001) Influence of shared epitope-negative HLA-DRB1 alleles on genetic susceptibility to rheumatoid arthritis. *Arthritis Rheum* 44: 535–540.
- du Montcel ST, Michou L, Petit-Teixeira E, Osorio J, Lemaire I, et al. (2005) New classification of HLA-DRB1 alleles supports the shared epitope hypothesis of rheumatoid arthritis susceptibility. *Arthritis Rheum* 52: 1063–1068.
- Klareskog L, Rönnelid J, Lundberg K, Padykov L, Alfredsson L (2008) Immunity to citrullinated proteins in rheumatoid arthritis. *Annu Rev Immunol* 26: 651–675.
- Aletaha D, Neogi T, Silman AJ, Funovits J, Felson DT, et al. (2010) Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum* 62: 2569–81.
- De Rooy DP, Willemze A, Mertens B, Huizinga TW, Van der Helm-van Hil AH, et al. (2011) Can anti-cyclic citrullinated peptide antibody-negative RA be subdivided into clinical subphenotypes? *Arthritis Res Ther* 13: R180.
- de Hair MJ, Lehmann KA, van de Sande MG, Maijjer KL, Gerlag DM, et al. (2012) The clinical picture of rheumatoid arthritis according to the 2010 American College of Rheumatology/European League Against Rheumatism criteria: is this still the same disease? *Arthritis Rheum* 64: 389–393.
- Benazet JF, Reviron D, Mercier P, Roux H, Roudier J (1995) HLA-DRB1 alleles associated with rheumatoid arthritis in southern France. Absence of extraarticular disease despite expression of the shared epitope. *J Rheumatol* 22: 607–610.
- Mackie SL, Taylor JC, Martin SG, YEAR Consortium, UKRAG Consortium, et al. (2012) A spectrum of susceptibility to rheumatoid arthritis within HLA-DRB1: stratification by autoantibody status in a large UK population. *Genes Immun* 13: 120–128.
- Raychaudhuri S, Sandor C, Stahl EA, Freudenberg J, Lee HS, et al. (2012) Five aminoacids in three HLA proteins explain most of the association between MHC and seropositive rheumatoid arthritis. *Nature Genetics* 44: 291–296.

Other Polymorphisms in the HLA Region may be Relevant to Susceptibility for Developing ACPA Positive RA

A recent very large-scale association study using 3000 SNPs in the MHC confirmed the importance of polymorphisms in the HLA-DRB1 chain, pinpointing three polymorphic residues, two of which are included in the shared epitope region. However, this study did not consider the genotypic effect, that is how BOTH HLA-DRB1 alleles interact to modulate risk for developing RA [12]. Here, we did not focus on the identification of new polymorphic residues in HLA-DRB1 alleles but on the interaction of individual alleles inside genotypes to influence RA susceptibility.

Author Contributions

Conceived and designed the experiments: JR NB DR IA CP E. Toussiro DW CC NL. Performed the experiments: CL EP DR CP. Analyzed the data: JR NB CP DR IA NL. Contributed reagents/materials/analysis tools: JR E. Telle DW CC. Wrote the paper: JR NB.