

Hydrodynamic limits for kinetic flocking models of Cucker-Smale type

P. Aceves-Sánchez ^{*}, M. Bostan [†], J.A. Carrillo [‡], P. Degond [§]

(April 22, 2019)

Abstract

We analyse the asymptotic behavior for kinetic models describing the collective behavior of animal populations. We focus on models for self-propelled individuals, whose velocity relaxes toward the mean orientation of the neighbors. The self-propelling and friction forces together with the alignment and the noise are interpreted as a collision/interaction mechanism acting with equal strength. We show that the set of generalized collision invariants, introduced in [39], is equivalent in our setting to the more classical notion of collision invariants, i.e., the kernel of a suitably linearized collision operator. After identifying these collision invariants, we derive the fluid model, by appealing to the balances for the particle concentration and orientation. We investigate the main properties of the macroscopic model for a general potential with radial symmetry.

Keywords: Vlasov-like equations, Swarming, Cucker-Smale model, Vicsek model.

AMS classification: 92D50, 82C40, 92C10.

1 Introduction

Flocking is observed in large populations of social agents such as birds [1], fish [51] or insects [19]. It refers to the emergence of large scale spatio-temporal structures which are not directly encoded in the individual agents' behavior. Understanding how large scale structures appear from individual behavior has sparked a huge literature in the recent years concerned with both

^{*}Department of Mathematics, Imperial College London, London SW7 2AZ UK. E-mail : p.aceves-sanchez@imperial.ac.uk

[†]Aix Marseille Université, CNRS, Centrale Marseille, Institut de Mathématiques de Marseille, UMR 7373, Château Gombert 39 rue F. Joliot Curie, 13453 Marseille Cedex 13 FRANCE. E-mail : mihai.bostan@univ-amu.fr

[‡]Department of Mathematics, Imperial College London, London SW7 2AZ UK. E-mail : carrillo@imperial.ac.uk

[§]Department of Mathematics, Imperial College London, London SW7 2AZ UK. E-mail : p.degond@imperial.ac.uk.

modelling [6, 27, 31, 58, 59, 60] and experiments [18, 28]. We refer the reader to the reviews [47, 52, 61]. Yet this phenomenon is still poorly understood. The modelling of individual behavior at the microscopic scale requires to resolve the motion of each individual in the course of time. This leads to so-called Individual-Based Models (IBM) (or particle models) consisting of a huge number of coupled ordinary or stochastic differential equations. By contrast, given the very large number of agents involved, the modelling of the macroscopic scale is best done through Continuum Models (CM). They describe the system as a fluid through average quantities such as the density or mean velocity of the agents. At the mesoscopic scale, Kinetic Models (KM) are intermediates between IBM and CM. They describe the agents dynamics through the statistical distribution of their positions and velocities. In the literature, IBM, KM or CM are often chosen according to the authors' preferences rather than following an explicit rationale. However, understanding flocking requires understanding how the individuals' microscale impacts on the system's macroscale and consequently demands the use of a consistent sequence of IBM, KM and CM models. This consistency can only be guaranteed if the passage from IBM to KM and from KM to CM can be systematically established. But this issue is seldom considered. One goal of this paper is to establish the consistency between a KM and a CM of flocking based on a variant of the celebrated Cucker-Smale model of consensus (Eqs. (1), (2) below). KM have been introduced in the last years for the mesoscopic description of collective behavior of agents/particles with applications in collective behavior of cell and animal populations, see [23, 39, 48, 55] and the references therein for a general overview on this active field. These models usually include alignment, attraction and repulsion as basic bricks of interactions between individuals.

In this paper, we will discard the microscopic scale, i.e. the IBM. We will directly consider the mesoscale, i.e. the KM and will focus on the derivation of the CM from the KM. Of course, our KM has an underlying IBM. However the techniques involved in the passage from IBM to KM are quite different from those needed to pass from KM to CM. For this reason, considering the IBM would have brought us beyond the scope of the present paper. We refer to [7, 8, 17, 20, 24, 25, 40, 45, 56] and references therein for a derivation of KM from IBM. The derivation of the CM from the KM requires a spatio-temporal rescaling. Indeed, the KM still describes - although in a statistical way compared to the IBM - the microscopic dynamics of the particles. In particular, it is written in space and time units that are of the order of the particle interaction distances and times. To describe the macroscopic scale, one needs to introduce a change of variables by which the space and time units become of the order of the system scale, which is much wider than the particle-related scales. This rescaling introduces a small parameter, the ratio of the microscopic to the macroscopic space units. The derivation of the CM from the KM consists in finding the limit of the KM when this small parameter tends to zero. This is the goal of this paper. We will see that the CM (Eqs (5), (6) below) corresponding to the considered modified Cucker-Smale model is a system consisting of equations for the density and mean orientation of the particles as functions of space and time. This system, referred to as the 'Self-Organized Hydrodynamics (SOH)' bears analogies with the Euler equations of isothermal compressible gas dynamics, with the important difference that the average velocity is replaced by the average orientation, i.e. a vector of norm one. The SOH appears in a variety of contexts related to alignment interactions, such as repulsion [33], nematic alignment [37], suspensions [38], solid orientation [36], and can be seen as a basic CM

of collective dynamics.

Here, we focus on the derivation of macroscopic equations (SOH) for the collective motion of self-propelled particles with alignment and noise when a cruise speed for individuals is imposed asymptotically for large times as in [2, 3, 21, 26, 29, 30, 41]. More precisely, in the presence of friction and self-propulsion and the absence of other interactions, individuals/particles accelerate or break to achieve a cruise speed exponentially fast in time. The alignment between particles is imposed via localized versions of the Cucker-Smale or Motsch-Tadmor reorientation procedure [22, 23, 32, 44, 46, 54] leading to relaxation terms to the mean velocity modulated or not by the density of particles. By scaling the relaxation time towards the asymptotic cruise speed, or equivalently, penalizing the balance between friction and self-propulsion, this alignment interaction leads asymptotically to variations of the classical kinetic Vicsek-Fokker-Planck equation with velocities on the sphere, see [13, 14, 34, 35, 39, 43, 60]. It was shown in [2, 3] that particular versions of the localized kinetic Cucker-Smale model can lead to phase transitions driven by noise. Moreover, these phase transitions are numerically stable in this asymptotic limit converging towards the phase transitions of the limiting versions of the corresponding kinetic Vicsek-Fokker-Planck equation.

In this work, we choose a localized and normalized version of the Cucker-Smale model not showing phase transition. More precisely, let us denote by $f = f(t, x, v) \geq 0$ the particle density in the phase space $(x, v) \in \mathbb{R}^d \times \mathbb{R}^d$, with $d \geq 2$. The standard self-propulsion/friction mechanism leading to the cruise speed of the particles in the absence of alignment is given by the term $\operatorname{div}_v \{f(\alpha - \beta|v|^2)v\}$ with $\alpha, \beta > 0$, and the relaxation toward the normalized mean velocity writes $\operatorname{div}_v \{f(v - \Omega[f])\}$. Here, for any particle density $f(x, v)$, the notation $\Omega[f]$ stands for the orientation of the mean velocity

$$\Omega[f] := \begin{cases} \frac{\int_{\mathbb{R}^d} f(\cdot, v)v \, dv}{\left| \int_{\mathbb{R}^d} f(\cdot, v)v \, dv \right|}, & \text{if } \int_{\mathbb{R}^d} f(\cdot, v)v \, dv \in \mathbb{R}^d \setminus \{0\}, \\ 0, & \text{if } \int_{\mathbb{R}^d} f(\cdot, v)v \, dv = 0. \end{cases}$$

Notice that we always have

$$\rho u[f] := \int_{\mathbb{R}^d} f(\cdot, v)v \, dv = \left| \int_{\mathbb{R}^d} f(\cdot, v)v \, dv \right| \Omega[f] \quad \text{with } \rho := \int_{\mathbb{R}^d} f(\cdot, v) \, dv.$$

Let us remark that the standard localized Cucker-Smale model would lead to $\rho \operatorname{div}_v \{f(v - u[f])\}$ while the localized Motsch-Tadmor model would lead to $\operatorname{div}_v \{f(v - u[f])\}$. Our relaxation term towards the normalized local velocity $\Omega[f]$ does not give rise to phase transition in the homogeneous setting on the limiting Vicsek-Fokker-Planck-type model on the sphere according to [34] and it produces a competition to the cruise speed term comprising a tendency towards unit speed. Including random Brownian fluctuations in the velocity variable leads to the kinetic Fokker-Planck type equation

$$\partial_t f + v \cdot \nabla_x f + \operatorname{div}_v \{f(\alpha - \beta|v|^2)v\} = \operatorname{div}_v \{\sigma \nabla_v f + f(v - \Omega[f])\}, \quad (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}^d \times \mathbb{R}^d.$$

We include this equation in a more general family of equations written in a compact form as

$$\partial_t f + v \cdot \nabla_x f = Q(f), \tag{1}$$

where

$$Q(f) = \operatorname{div}_v \{ \sigma \nabla_v f + f(v - \Omega[f]) + \eta f \nabla_v V \}, \quad (2)$$

for any density distribution f with V a general confining potential in the velocity variables and $\eta > 0$ (see Lemma 2.1 for more information on the type of potentials that we consider). In the particular example considered above we take $V = V_{\alpha,\beta}(|v|) := \beta \frac{|v|^4}{4} - \alpha \frac{|v|^2}{2}$.

We investigate the large time and space scale regimes of the kinetic transport equation (1) with collision operator given by (2). Namely, we study the asymptotic behavior when $\varepsilon \rightarrow 0$ of

$$\partial_t f^\varepsilon + v \cdot \nabla_x f^\varepsilon = \frac{1}{\varepsilon} Q(f^\varepsilon), \quad (3)$$

supplemented with the initial condition

$$f^\varepsilon(0, x, v) = f^{\text{in}}(x, v), \quad (x, v) \in \mathbb{R}^d \times \mathbb{R}^d.$$

The rescaling taken in the kinetic transport equation (3) with confining potential $V_{\alpha,\beta}$ can be seen as an intermediate scaling between the ones proposed in [14] and [2]. The difference being that we have a relaxation towards the normalized mean velocity $\Omega[f]$ rather than the mean velocity $u[f]$ as in [2, 14]. This difference is important since in the first case there is no phase transition in the homogeneous limiting setting on the sphere as we mentioned above, while in the second there is, see [3, 14, 34]. In fact, in [14] the scaling corresponds to $\eta = 1/\varepsilon$ in (3), that is the relaxation to the cruise speed is penalized with a term of the order of $1/\varepsilon^2$. Whereas in [2] the scaling corresponds to $\eta = \varepsilon$, that is the cruise speed is not penalized at all.

The methodology followed in [14] lies within the context of measure solutions by introducing a projection operator onto the set of measures supported in the sphere whose radius is the critical speed $r = \sqrt{\alpha/\beta}$. These technicalities are needed because the zeroth order expansion of f_ε lives on the sphere. This construction followed closely the average method in gyro-kinetic theory [10, 11, 12].

However, in our present case we will show in contrast to [2, 14] that there are no phase transitions which is in accordance with the results obtained in [35] for the kinetic Vicsek-Fokker-Planck equation with analogous alignment operator on the sphere. A modified version of (1)-(2) in which phase-transitions occur was studied in [3] whose analysis is postponed to a future work to focus here on the mathematical difficulties of the asymptotic analysis. Another difference in the present case is that the zeroth order expansion of f_ε will be parameterized by Von Mises-Fisher distributions in the whole velocity space, that is $f(t, x, v) = \rho(t, x) M_{\Omega(t,x)}(v)$, with ρ and Ω being, respectively, the density and the mean orientation of the particles. And where for any $\Omega \in \mathbb{S}^{d-1}$ we define (see Sect. 2)

$$M_\Omega(v) = \frac{1}{Z_\Omega} \exp\left(-\frac{\Phi_\Omega(v)}{\sigma}\right), \quad \text{with } Z_\Omega = \int_{\mathbb{R}^d} \exp\left(-\frac{\Phi_\Omega(v')}{\sigma}\right) dv' \quad (4)$$

and

$$\Phi(v) = \frac{|v - \Omega|^2}{2} + V(|v|).$$

The main result of this paper is the asymptotic analysis of the singularly perturbed kinetic transport equation of Cucker-Smale type (3). The particle density ρ and the orientation Ω obey the hydrodynamic type equations given in the following result.

Theorem 1.1 *Let $f^{\text{in}} \geq 0$ be a smooth initial particle density with nonvanishing orientation at any $x \in \mathbb{R}^d$. For any $\varepsilon > 0$ we consider the problem*

$$\partial_t f^\varepsilon + v \cdot \nabla_x f^\varepsilon = \frac{1}{\varepsilon} \operatorname{div}_v \{ \sigma \nabla_v f^\varepsilon + f^\varepsilon (v - \Omega[f^\varepsilon]) + f^\varepsilon \nabla_v V(|v|) \}, \quad (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}^d \times \mathbb{R}^d,$$

with initial condition

$$f^\varepsilon(0, x, v) = f^{\text{in}}(x, v), \quad (x, v) \in \mathbb{R}^d \times \mathbb{R}^d.$$

At any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$ the leading order term in the Hilbert expansion $f_\varepsilon = f + \varepsilon f_1 + \dots$ is an equilibrium distribution of Q , that is $f(t, x, v) = \rho(t, x) M_{\Omega(t, x)}(v)$ with $M_{\Omega(t, x)}(v)$ defined in (4), where the concentration ρ and the orientation Ω satisfy

$$\partial_t \rho + \operatorname{div}_x (\rho c_1 \Omega) = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d, \quad (5)$$

$$\partial_t \Omega + c_2 (\Omega \cdot \nabla_x) \Omega + \sigma (I_d - \Omega \otimes \Omega) \frac{\nabla_x \rho}{\rho} = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d, \quad (6)$$

with initial conditions

$$\rho(0, x) = \int_{\mathbb{R}^d} f^{\text{in}}(x, v) \, dv, \quad \Omega(0, x) = \frac{\int_{\mathbb{R}^d} f^{\text{in}}(x, v) v \, dv}{\left| \int_{\mathbb{R}^d} f^{\text{in}}(x, v) v \, dv \right|}, \quad x \in \mathbb{R}^d.$$

The constants c_1, c_2 are given by

$$c_1 = \frac{\int_{\mathbb{R}_+} r^d \int_0^\pi \cos \theta e(\cos \theta, r) \sin^{d-2} \theta \, d\theta \, dr}{\int_{\mathbb{R}_+} r^{d-1} \int_0^\pi e(\cos \theta, r) \sin^{d-2} \theta \, d\theta \, dr},$$

$$c_2 = \frac{\int_{\mathbb{R}_+} r^{d+1} \int_0^\pi \cos \theta \chi(\cos \theta, r) e(\cos \theta, r) \sin^{d-1} \theta \, d\theta \, dr}{\int_{\mathbb{R}_+} r^d \int_0^\pi \chi(\cos \theta, r) e(\cos \theta, r) \sin^{d-1} \theta \, d\theta \, dr},$$

and the function χ solves

$$\begin{aligned} -\sigma \partial_c \left[r^{d-3} (1 - c^2)^{\frac{d-1}{2}} e(c, r) \partial_c \chi \right] - \sigma \partial_r \left[r^{d-1} (1 - c^2)^{\frac{d-3}{2}} e(c, r) \partial_r \chi \right] + \sigma (d-2) r^{d-3} (1 - c^2)^{\frac{d-5}{2}} e \chi \\ = r^d (1 - c^2)^{\frac{d-2}{2}} e(c, r), \end{aligned}$$

where $e(c, r) = \exp(rc/\sigma) \exp(-(r^2 + 1)/(2\sigma) - V(r)/\sigma)$.

Our article is organized as follows. First, in Section 2 we state auxiliary results allowing us to discuss the kernel of the collision operator. Then in Section 3 we concentrate on the characterization of the collision invariants. We prove that the generalized collision invariants introduced in [39] coincide with the kernel of a suitable linearised collision operator. We explicitly describe the collision invariants in Section 4 and investigate their symmetries. Finally, the limit fluid model is determined in Section 5 and we analyse its main properties.

2 Preliminaries

Plugging into (3) the Hilbert expansion

$$f^\varepsilon = f + \varepsilon f^1 + \dots,$$

we obtain at the leading order

$$Q(f) = 0, \tag{7}$$

whereas to the next order we get

$$\partial_t f + v \cdot \nabla_x f = \lim_{\varepsilon \searrow 0} \frac{1}{\varepsilon} \{Q(f^\varepsilon) - Q(f)\} = dQ_f(f^1) =: \mathcal{L}_f(f^1), \tag{8}$$

where dQ_f denotes the first variation of Q with respect to f . The constraint (7) leads immediately to the equilibrium

$$M_\Omega(v) = \frac{1}{Z_\Omega} \exp\left(-\frac{\Phi_\Omega(v)}{\sigma}\right), \quad \text{with } Z_\Omega = \int_{\mathbb{R}^d} \exp\left(-\frac{\Phi_\Omega(v')}{\sigma}\right) dv',$$

where

$$\Phi_\Omega(v) = \frac{|v - \Omega|^2}{2} + V(|v|). \tag{9}$$

Indeed, by using the identity

$$\nabla_v M_\Omega = -\frac{M_\Omega(v)}{\sigma} \nabla_v \Phi_\Omega = -\frac{M_\Omega(v)}{\sigma} (v - \Omega + \nabla_v V(|v|)), \tag{10}$$

we can recast the operator Q as

$$Q(f) = \operatorname{div}_v (\sigma \nabla_v f + f \nabla_v \Phi_{\Omega[f]}) = \sigma \operatorname{div}_v \left[M_{\Omega[f]} \nabla_v \left(\frac{f}{M_{\Omega[f]}} \right) \right].$$

We denote by \mathbb{S}^{d-1} the set of unit vectors in \mathbb{R}^d . For any $\Omega \in \mathbb{S}^{d-1}$, we consider the weighted spaces

$$L_{M_\Omega}^2 = \left\{ \chi : \mathbb{R}^d \rightarrow \mathbb{R} \text{ measurable, } \int_{\mathbb{R}^d} (\chi(v))^2 M_\Omega(v) dv < \infty \right\},$$

and

$$H_{M_\Omega}^1 = \left\{ \chi : \mathbb{R}^d \rightarrow \mathbb{R} \text{ measurable, } \int_{\mathbb{R}^d} [(\chi(v))^2 + |\nabla_v \chi|^2] M_\Omega(v) dv < \infty \right\}.$$

The nonlinear operator Q should be understood in the distributional sense, and is defined for any particle density $f = f(v)$ in the domain

$$\begin{aligned} D(Q) &= \left\{ f : \mathbb{R}^d \rightarrow \mathbb{R}_+ \text{ measurable, } f/M_{\Omega[f]} \in H_{M_{\Omega[f]}}^1 \right\} \\ &= \left\{ f : \mathbb{R}^d \rightarrow \mathbb{R}_+ \text{ measurable, } \int_{\mathbb{R}^d} \left\{ \left(\frac{f}{M_{\Omega[f]}} \right)^2 + \left| \nabla_v \left(\frac{f}{M_{\Omega[f]}} \right) \right|^2 \right\} M_{\Omega[f]}(v) dv < \infty \right\}. \end{aligned}$$

We introduce the usual scalar products

$$(\chi, \theta)_{M_\Omega} = \int_{\mathbb{R}^d} \chi(v) \theta(v) M_\Omega(v) \, dv, \quad \chi, \theta \in L^2_{M_\Omega},$$

$$((\chi, \theta))_{M_\Omega} = \int_{\mathbb{R}^d} (\chi(v) \theta(v) + \nabla_v \chi \cdot \nabla_v \theta) M_\Omega(v) \, dv, \quad \chi, \theta \in H^1_{M_\Omega},$$

and we denote by $|\cdot|_{M_\Omega}, \|\cdot\|_{M_\Omega}$ the associated norms. We make the following hypotheses on the potential V . We assume that for any $\Omega \in \mathbb{S}^{d-1}$ we have

$$Z_\Omega = \int_{\mathbb{R}^d} \exp\left(-\frac{1}{\sigma} \left[\frac{|v - \Omega|^2}{2} + V(|v|)\right]\right) \, dv < \infty. \quad (11)$$

Clearly (11) holds true for the potentials $V_{\alpha,\beta}$. Notice that in that case $1 \in L^2_{M_\Omega}$ and $|1|_{M_\Omega} = 1$ for any $\Omega \in \mathbb{S}^{d-1}$. Moreover, we need a Poincaré inequality, that is, for any $\Omega \in \mathbb{S}^{d-1}$ there is $\lambda_\Omega > 0$ such that for all $\chi \in H^1_{M_\Omega}$ we have

$$\sigma \int_{\mathbb{R}^d} |\nabla_v \chi|^2 M_\Omega(v) \, dv \geq \lambda_\Omega \int_{\mathbb{R}^d} \left| \chi(v) - \int_{\mathbb{R}^d} \chi(v') M_\Omega(v') \, dv' \right|^2 M_\Omega(v) \, dv. \quad (12)$$

A sufficient condition for (12) to hold comes from the well-known equivalence between the Fokker-Planck and Schrödinger operators (see for instance [9]). Namely, for any $\Omega \in \mathbb{S}^{d-1}$ we have

$$-\frac{\sigma}{\sqrt{M_\Omega}} \operatorname{div}_v \left(M_\Omega \nabla_v \left(\frac{u}{\sqrt{M_\Omega}} \right) \right) = -\sigma \Delta_v u + \left[\frac{1}{4\sigma} |\nabla_v \Phi_\Omega|^2 - \frac{1}{2} \Delta_v \Phi_\Omega \right] u.$$

The operator $\mathcal{H}_\Omega = -\sigma \Delta_v + \left[\frac{1}{4\sigma} |\nabla_v \Phi_\Omega|^2 - \frac{1}{2} \Delta_v \Phi_\Omega \right]$ is defined in the domain

$$D(\mathcal{H}_\Omega) = \left\{ u \in L^2(\mathbb{R}^d), \left[\frac{1}{4\sigma} |\nabla_v \Phi_\Omega|^2 - \frac{1}{2} \Delta_v \Phi_\Omega \right] u \in L^2(\mathbb{R}^d) \right\}.$$

Using classical results for Schrödinger operators (see for instance Theorem XIII.67 in [57]), we have a spectral decomposition of the operator \mathcal{H}_Ω under suitable confining assumptions.

Lemma 2.1 *Assume that for Φ_Ω defined in (9) the function $v \rightarrow \frac{1}{4\sigma} |\nabla_v \Phi_\Omega|^2 - \frac{1}{2} \Delta_v \Phi_\Omega$ satisfies the following:*

- a) *it belongs to $L^1_{\text{loc}}(\mathbb{R}^d)$,*
- b) *it is bounded from below,*
- c)

$$\lim_{|v| \rightarrow \infty} \left[\frac{1}{4\sigma} |\nabla_v \Phi_\Omega|^2 - \frac{1}{2} \Delta_v \Phi_\Omega \right] = \infty.$$

Then \mathcal{H}_Ω^{-1} is a self adjoint compact operator in $L^2(\mathbb{R}^d)$ and \mathcal{H}_Ω admits a spectral decomposition, that is a nondecreasing sequence of real numbers $(\lambda_\Omega^n)_{n \in \mathbb{N}}$, $\lim_{n \rightarrow \infty} \lambda_\Omega^n = \infty$, and a $L^2(\mathbb{R}^d)$ -orthonormal basis $(\psi_\Omega^n)_{n \in \mathbb{N}}$ such that $\mathcal{H}_\Omega \psi_\Omega^n = \lambda_\Omega^n \psi_\Omega^n$, $n \in \mathbb{N}$, $\lambda_\Omega^0 = 0$, $\lambda_\Omega^1 > 0$.

Let us note that the spectral gap of the Schrödinger operator \mathcal{H}_Ω is the Poincaré constant in the Poincaré inequality (12). Notice also that the hypotheses in Lemma 2.1 are satisfied by the potentials $V_{\alpha, \beta}$, and therefore (12) holds true in that case. It is easily seen that the set of equilibrium distributions of Q is parametrized by d parameters as stated in the following result.

Lemma 2.2 *Let $f = f(v) \geq 0$ be a function in $D(Q)$. Then f is an equilibrium for Q if and only if there are $(\rho, \Omega) \in \mathbb{R}_+ \times (\mathbb{S}^{d-1} \cup \{0\})$ such that $f = \rho M_\Omega$. Moreover we have $\rho = \rho[f] := \int_{\mathbb{R}^d} f(v) dv$ and $\Omega = \Omega[f]$.*

Proof.

If f is an equilibrium for Q , we have

$$\sigma \int_{\mathbb{R}^d} \left| \nabla_v \left(\frac{f}{M_{\Omega[f]}} \right) \right|^2 M_{\Omega[f]}(v) dv = 0,$$

and therefore there is $\rho \in \mathbb{R}$ such that $f = \rho M_{\Omega[f]}$. Obviously $\rho = \int_{\mathbb{R}^d} f(v) dv \geq 0$ and $\Omega[f] \in \mathbb{S}^{d-1} \cup \{0\}$. Conversely, we claim that for any $(\rho, \Omega) \in \mathbb{R}_+ \times (\mathbb{S}^{d-1} \cup \{0\})$, the particle density $f = \rho M_\Omega$ is an equilibrium for Q . Indeed, we have

$$\sigma \nabla_v (\rho M_\Omega) + \rho M_\Omega (v - \Omega + \nabla_v V) = \rho (\sigma \nabla_v M_\Omega + M_\Omega \nabla_v \Phi_\Omega) = 0.$$

We are done if we prove that $\Omega[f] = \Omega$. If $\Omega = 0$, it is easily seen that

$$\int_{\mathbb{R}^d} f(v) v dv = \rho \int_{\mathbb{R}^d} \frac{1}{Z_0} \exp \left(-\frac{\Phi_0(v)}{\sigma} \right) v dv = \frac{\rho}{Z_0} \int_{\mathbb{R}^d} \exp \left(-\frac{1}{\sigma} \left(\frac{|v|^2}{2} + V(|v|) \right) \right) v dv = 0,$$

implying $\Omega[f] = 0 = \Omega$. Assume now that $\Omega \in \mathbb{S}^{d-1}$. For any $\xi \in \mathbb{S}^{d-1}$, $\xi \cdot \Omega = 0$, we consider the orthogonal transformation $\mathcal{O}_\xi = I_d - 2\xi \otimes \xi$. Thanks to the change of variable $v = \mathcal{O}_\xi v'$, we write

$$\begin{aligned} \int_{\mathbb{R}^d} (v \cdot \xi) f(v) dv &= \frac{\rho}{Z_\Omega} \int_{\mathbb{R}^d} (v \cdot \xi) M_\Omega(v) dv = \frac{\rho}{Z_\Omega} \int_{\mathbb{R}^d} (\mathcal{O}_\xi v' \cdot \xi) M_\Omega(\mathcal{O}_\xi v') dv' \\ &= -\frac{\rho}{Z_\Omega} \int_{\mathbb{R}^d} (v' \cdot \xi) M_\Omega(v') dv' = -\int_{\mathbb{R}^d} (v' \cdot \xi) f(v') dv', \end{aligned}$$

where we have used the radial symmetry of V , $\mathcal{O}_\xi \xi = -\xi$ and $\mathcal{O}_\xi \Omega = \Omega$. We deduce that $\int_{\mathbb{R}^d} f(v) v dv = \int_{\mathbb{R}^d} (v \cdot \Omega) f(v) dv \Omega$. We claim that $\int_{\mathbb{R}^d} (v \cdot \Omega) f(v) dv > 0$. Indeed we have

$$\begin{aligned} \int_{\mathbb{R}^d} (v \cdot \Omega) f(v) dv &= \frac{\rho}{Z_\Omega} \int_{v \cdot \Omega > 0} (v \cdot \Omega) M_\Omega(v) dv + \frac{\rho}{Z_\Omega} \int_{v \cdot \Omega < 0} (v \cdot \Omega) M_\Omega(v) dv \\ &= \frac{\rho}{Z_\Omega} \int_{v \cdot \Omega > 0} (v \cdot \Omega) \left[\exp \left(-\frac{\Phi_\Omega(v)}{\sigma} \right) - \exp \left(-\frac{\Phi_\Omega(-v)}{\sigma} \right) \right] dv. \end{aligned}$$

Obviously, we have for any $v \in \mathbb{R}^d$ such that $v \cdot \Omega > 0$

$$-\frac{\Phi_\Omega(v)}{\sigma} + \frac{\Phi_\Omega(-v)}{\sigma} = -\frac{|v - \Omega|^2}{2\sigma} + \frac{|-v - \Omega|^2}{2\sigma} = 2\frac{v \cdot \Omega}{\sigma} > 0,$$

implying that $\int_{\mathbb{R}^d} (v \cdot \Omega) f(v) dv > 0$ and

$$\Omega[f] = \frac{\int_{\mathbb{R}^d} f(v) v dv}{\left| \int_{\mathbb{R}^d} f(v) v dv \right|} = \frac{\int_{\mathbb{R}^d} (v \cdot \Omega) f(v) dv \Omega}{\left| \int_{\mathbb{R}^d} (v \cdot \Omega) f(v) dv \Omega \right|} = \Omega.$$

□

3 Characterization of the collision invariants

In [39], the following notion of generalized collision invariant (GCI) has been introduced.

Definition 3.1 (GCI)

Let $\Omega \in \mathbb{S}^{d-1}$ be a fixed orientation. A function $\psi = \psi(v)$ is called a generalized collision invariant of Q associated to Ω , if and only if

$$\int_{\mathbb{R}^d} Q(f)(v) \psi(v) dv = 0,$$

for all f such that $(I_d - \Omega \otimes \Omega) \int_{\mathbb{R}^d} f(v) v dv = 0$, that is such that $\int_{\mathbb{R}^d} f(v) v dv \in \mathbb{R}\Omega$.

In order to obtain the hydrodynamic limit of (3), for any fixed $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, we multiply (8) by a function $v \rightarrow \psi_{t,x}(v)$ and integrate with respect to v yielding

$$\begin{aligned} \int_{\mathbb{R}^d} \partial_t f(t, x, v) \psi_{t,x}(v) dv + \int_{\mathbb{R}^d} v \cdot \nabla_x f(t, x, v) \psi_{t,x}(v) dv &= \int_{\mathbb{R}^d} \mathcal{L}_{f(t,x,\cdot)}(f^1(t, x, \cdot)) \psi_{t,x}(v) dv \\ &= \int_{\mathbb{R}^d} f^1(t, x, v) (\mathcal{L}_{f(t,x,\cdot)}^* \psi_{t,x})(v) dv. \end{aligned} \quad (13)$$

The above computation leads naturally to the following extension of the notion of collision invariant, see also [14].

Definition 3.2 Let $f = f(v) \geq 0$ be an equilibrium of Q . A function $\psi = \psi(v)$ is called a collision invariant for Q associated to the equilibrium f , if and only if $\mathcal{L}_f^* \psi = 0$, that is

$$\int_{\mathbb{R}^d} (\mathcal{L}_f g)(v) \psi(v) dv = 0 \quad \text{for any function } g = g(v).$$

We are looking for a good characterization of the linearized collision operator \mathcal{L}_f and its adjoint with respect to the leading order particle density f . Motivated by (7), we need to determine the structure of the equilibria of Q which are given by Lemma 2.2.

By Lemma 2.2, we know that for any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, there are $(\rho(t, x), \Omega(t, x)) \in \mathbb{R}_+ \times (\mathbb{S}^{d-1} \cup \{0\})$ such that $f(t, x, \cdot) = \rho(t, x)M_{\Omega(t, x)}$, where

$$\rho(t, x) = \rho[f(t, x, \cdot)] \quad \text{and} \quad \Omega(t, x) = \Omega[f(t, x, \cdot)].$$

The evolution of the macroscopic quantities ρ and Ω follows from (8) and (13), by appealing to the moment method [4, 5, 15, 16, 49, 50]. Next, we explicitly determine the linearization of the collision operator Q around its equilibrium distributions. For any orientation $\Omega \in \mathbb{S}^{d-1} \cup \{0\}$ we introduce the pressure tensor

$$\mathcal{M}_\Omega := \int_{\mathbb{R}^d} (v - \Omega) \otimes (I_d - \Omega \otimes \Omega)(v - \Omega) M_\Omega(v) \, dv,$$

and the quantity

$$c_1 := \int_{\mathbb{R}^d} (v \cdot \Omega) M_\Omega(v) \, dv > 0.$$

We will check later, see Lemma 3.2, that the pressure tensor \mathcal{M}_Ω is symmetric.

Proposition 3.1 *Let $f = f(v) \geq 0$ be an equilibrium distribution of Q with nonvanishing orientation, that is*

$$f = \rho M_\Omega, \quad \text{where} \quad \rho = \rho[f], \quad \text{and} \quad \Omega = \Omega[f] \in \mathbb{S}^{d-1}.$$

1. *The linearization $\mathcal{L}_f = dQ_f$ is given by*

$$\mathcal{L}_f g = \operatorname{div}_v \left\{ \sigma \nabla_v g + g \nabla_v \Phi_\Omega - \frac{f}{\int_{\mathbb{R}^d} (v \cdot \Omega) f(v) \, dv} P_f \int_{\mathbb{R}^d} g(v) v \, dv \right\},$$

where $P_f := I_d - \Omega[f] \otimes \Omega[f]$ is the orthogonal projection onto $\{\xi \in \mathbb{R}^d : \xi \cdot \Omega[f] = 0\}$. In particular $\mathcal{L}_{\rho M_\Omega} = \mathcal{L}_{M_\Omega}$.

2. *The formal adjoint of \mathcal{L}_f is given by*

$$\mathcal{L}_f^* \psi = \sigma \frac{\operatorname{div}_v (M_\Omega \nabla_v \psi)}{M_\Omega} + P_f v \cdot W[\psi], \quad W[\psi] := \frac{\int_{\mathbb{R}^d} M_\Omega(v) \nabla_v \psi \, dv}{\int_{\mathbb{R}^d} (v \cdot \Omega) M_\Omega(v) \, dv}. \quad (14)$$

3. *We have the identity*

$$\mathcal{L}_f(f(v - \Omega)) = \sigma \nabla_v f - \operatorname{div}_v \left(f \frac{\mathcal{M}_\Omega}{c_1} \right).$$

Note that div_v refers to the divergence operator acting on matrices defined as applying the divergence operator over rows.

Proof.

1. By standard computations we have

$$\mathcal{L}_f g = \frac{d}{ds} \Big|_{s=0} Q(f + sg) = \operatorname{div}_v \left\{ \sigma \nabla_v g + g(v - \Omega[f] + \nabla_v V) - f \frac{d}{ds} \Big|_{s=0} \Omega[f + sg] \right\},$$

and

$$\frac{d}{ds}\Big|_{s=0} \Omega[f + sg] = \frac{(I_d - \Omega[f] \otimes \Omega[f])}{\left| \int_{\mathbb{R}^d} f(v)v \, dv \right|} \int_{\mathbb{R}^d} g(v)v \, dv.$$

Therefore we obtain

$$\mathcal{L}_f g = \operatorname{div}_v \left\{ \sigma \nabla_v g + g \nabla_v \Phi_\Omega - \frac{f}{\int_{\mathbb{R}^d} (v \cdot \Omega) f(v) \, dv} P_f \int_{\mathbb{R}^d} g(v)v \, dv \right\}.$$

2. We have

$$\begin{aligned} & \int_{\mathbb{R}^d} (\mathcal{L}_f g)(v) \psi(v) \, dv \\ &= - \int_{\mathbb{R}^d} \left\{ \sigma \nabla_v g + g \nabla_v \Phi_\Omega - \frac{f}{\int_{\mathbb{R}^d} (v' \cdot \Omega) f(v') \, dv'} P_f \int_{\mathbb{R}^d} g(v')v' \, dv' \right\} \cdot \nabla_v \psi \, dv \\ &= \int_{\mathbb{R}^d} g [\sigma \operatorname{div}_v \nabla_v \psi - \nabla_v \psi \cdot \nabla_v \Phi_\Omega] \, dv + \int_{\mathbb{R}^d} g(v') P_f v' \cdot \frac{\int_{\mathbb{R}^d} f(v) \nabla_v \psi(v) \, dv}{\int_{\mathbb{R}^d} (v \cdot \Omega) f(v) \, dv} \, dv', \end{aligned}$$

implying

$$\mathcal{L}_f^* \psi = \sigma \frac{\operatorname{div}_v (M_\Omega \nabla_v \psi)}{M_\Omega} + P_f v \cdot W[\psi].$$

3. For any $i \in \{1, \dots, d\}$ we have

$$\mathcal{L}_f(f(v - \Omega)_i) = \operatorname{div}_v \left\{ (v - \Omega)_i (\sigma \nabla_v f + f \nabla_v \Phi_\Omega) + \sigma f e_i - f \frac{\int_{\mathbb{R}^d} M_\Omega(v') (v' - \Omega)_i P_f v' \, dv'}{\int_{\mathbb{R}^d} (v' \cdot \Omega) M_\Omega(v') \, dv'} \right\},$$

and therefore, since $f = \rho M_\Omega$ satisfies $\sigma \nabla_v f + f \nabla_v \Phi_\Omega = 0$, we get

$$\begin{aligned} \mathcal{L}_f(f(v - \Omega)_i) &= \sigma \nabla_v f - \operatorname{div}_v \left(f \frac{\int_{\mathbb{R}^d} M_\Omega(v') (v' - \Omega)_i \otimes P_f v' \, dv'}{\int_{\mathbb{R}^d} (v' \cdot \Omega) M_\Omega(v') \, dv'} \right) \\ &= \sigma \nabla_v f - \operatorname{div}_v \left(f \frac{\mathcal{M}_\Omega}{c_1} \right). \end{aligned}$$

□

Notice that at any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, the function $h = 1$ is a collision invariant for Q , associated to $f(t, x, \cdot)$. Indeed, for any $g = g(v)$ we have

$$\int_{\mathbb{R}^d} Q(f(t, x, \cdot) + sg) \, dv = 0,$$

implying that $\int_{\mathbb{R}^d} (\mathcal{L}_{f(t,x,\cdot)} g)(v) \, dv = 0$ and therefore $\mathcal{L}_{f(t,x,\cdot)}^* 1 = 0$, for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$. Once we have determined a collision invariant $\psi = \psi(t, x, v)$ at any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, we deduce, thanks to (13), a balance for the macroscopic quantities $\rho(t, x) = \rho[f(t, x, \cdot)]$ and $\Omega(t, x) = \Omega[f(t, x, \cdot)]$, given by the relationship

$$\int_{\mathbb{R}^d} \partial_t (\rho M_{\Omega(t,x)}) \psi(t, x, v) \, dv + \int_{\mathbb{R}^d} v \cdot \nabla_x (\rho M_{\Omega(t,x)}) \psi(t, x, v) \, dv = 0. \quad (15)$$

When taking as collision invariant the function $h(t, x, v) = 1$, we obtain the local mass conservation equation

$$\partial_t \rho + \operatorname{div}_x \left(\rho \int_{\mathbb{R}^d} (v \cdot \Omega(t, x)) M_{\Omega(t, x)}(v) \, dv \, \Omega \right) = 0. \quad (16)$$

As usual, we are looking also for the conservation of the total momentum, however, the nonlinear operator Q does not preserve momentum. In other words, v is not a collision invariant. Indeed, if $f = \rho M_{\Omega}$ is an equilibrium with nonvanishing orientation, we have

$$\mathcal{L}_f^* v = \sigma \frac{\nabla_v M_{\Omega}}{M_{\Omega}} + \frac{P_f v}{\int_{\mathbb{R}^d} (v' \cdot \Omega) M_{\Omega}(v') \, dv'} = -\nabla_v \Phi_{\Omega} + \frac{P_f v}{\int_{\mathbb{R}^d} (v' \cdot \Omega) M_{\Omega}(v') \, dv'},$$

and therefore v is not a collision invariant.

We concentrate next on the resolution of (14). We will use the notation $\partial_v \xi = \left(\frac{\partial \xi_i}{\partial v_j} \right)$ for the Jacobian matrix of a vector field ξ and div_v for the divergence operator in v of both vectors and matrices with the convention of taking the divergence over the rows of the matrix. With this convention, we have

$$\int_{\mathbb{R}^d} g \operatorname{div}_v A \, dv = - \int_{\mathbb{R}^d} A \nabla_v g \, dv \quad \text{and} \quad \int_{\mathbb{R}^d} \xi \operatorname{div}_v \eta \, dv = - \int_{\mathbb{R}^d} \partial_v \xi \eta \, dv \quad (17)$$

for all smooth functions g , vector fields ξ, η , and matrices A . We now focus in finding a parameterization of the kernel of the operator \mathcal{L}_f^* .

Lemma 3.1 *Let $f = \rho M_{\Omega}$ be an equilibrium of Q with nonvanishing orientation. The following two statements are equivalent:*

1. $\psi = \psi(v)$ is a collision invariant for Q associated to f .
2. ψ satisfies

$$\sigma \frac{\operatorname{div}_v (M_{\Omega} \nabla_v \psi)}{M_{\Omega}} + P_f v \cdot W = 0, \quad (18)$$

for some vector $W \in \ker(\mathcal{M}_{\Omega} - \sigma c_1 I_d)$.

Moreover, the linear map $W : \ker(\mathcal{L}_f^*) \rightarrow \ker(\mathcal{M}_{\Omega} - \sigma c_1 I_d)$, with $W[\psi] := \int_{\mathbb{R}^d} M_{\Omega}(v) \nabla_v \psi \, dv / c_1$ induces an isomorphism between the vector spaces $\ker(\mathcal{L}_f^*) / \ker W$ and $\ker(\mathcal{M}_{\Omega} - \sigma c_1 I_d)$, where $\ker W$ is the set of the constant functions.

Proof.

1. \implies 2. Since ψ is a collision invariant associated to f , i.e. $\mathcal{L}_f^* \psi = 0$, and by the third statement in Proposition 3.1 we deduce (using also the first formula in (17) with $f \mathcal{M}_{\Omega} / c_1$ and ψ)

$$\begin{aligned} 0 &= \int_{\mathbb{R}^d} \mathcal{L}_f^* \psi \, f(v - \Omega) \, dv = \int_{\mathbb{R}^d} \psi(v) \mathcal{L}_f (f(v - \Omega)) \, dv \\ &= \int_{\mathbb{R}^d} \psi(v) \left[\sigma \nabla_v f - \operatorname{div}_v \left(f \frac{\mathcal{M}_{\Omega}}{c_1} \right) \right] \, dv = -\sigma \int_{\mathbb{R}^d} f(v) \nabla_v \psi \, dv + \mathcal{M}_{\Omega} \int_{\mathbb{R}^d} \frac{f(v) \nabla_v \psi}{c_1} \, dv \\ &= -\rho \sigma c_1 W[\psi] + \rho \mathcal{M}_{\Omega} W[\psi]. \end{aligned}$$

Note that if $\rho = 0$ then $f = 0$ and $\int v f dv = 0$, implying that $\Omega[f] = 0$. Hence, since $\Omega \neq 0$, we have $\rho > 0$ and thus $W[\psi] \in \ker(\mathcal{M}_\Omega - \sigma c_1 I_d)$, saying that (18) holds true with $W = W[\psi] \in \ker(\mathcal{M}_\Omega - \sigma c_1 I_d)$.

2. \implies 1. Let ψ be a function satisfying (18) for some vector $W \in \ker(\mathcal{M}_\Omega - \sigma c_1 I_d)$. Multiplying (18) by $f(v - \Omega)$ and integrating with respect to v yields (thanks to the second formula in (17))

$$-\sigma \rho \int_{\mathbb{R}^d} \partial_v(v - \Omega) \nabla_v \psi M_\Omega(v) dv + \rho \mathcal{M}_\Omega W = 0,$$

which implies $W[\psi] = W$ since $\mathcal{M}_\Omega W = \sigma c_1 W$ by the assumption $W \in \ker(\mathcal{M}_\Omega - \sigma c_1 I_d)$. Therefore ψ is a collision invariant for Q , associated to f

$$\mathcal{L}_f^* \psi = \sigma \frac{\operatorname{div}_v(M_\Omega \nabla_v \psi)}{M_\Omega} + P_f v \cdot W[\psi] = \sigma \frac{\operatorname{div}_v(M_\Omega \nabla_v \psi)}{M_\Omega} + P_f v \cdot W = 0.$$

□

Remark 3.1 For any non negative measurable function $\chi = \chi(c, r) :]-1, +1[\times]0, \infty[\rightarrow \mathbb{R}$ and any $\Omega \in \mathbb{S}^{d-1}$, for $d \geq 2$, we have

$$\int_{\mathbb{R}^d} \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) dv = |\mathbb{S}^{d-2}| \int_{\mathbb{R}_+} \int_0^\pi \chi(\cos \theta, r) r^{d-1} \sin^{d-2} \theta d\theta dr,$$

where $|\mathbb{S}^{d-2}|$ is the surface of the unit sphere in \mathbb{R}^{d-1} , for $d \geq 3$, and $|\mathbb{S}^0| = 2$ for $d = 2$. In particular we have the formula

$$\begin{aligned} \int_{\mathbb{R}^d} \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) M_\Omega(v) dv &= \frac{\int_{\mathbb{R}_+} r^{d-1} \int_0^\pi \chi(\cos \theta, r) e(\cos \theta, r) \sin^{d-2} \theta d\theta dr}{\int_{\mathbb{R}_+} r^{d-1} \int_0^\pi e(\cos \theta, r) \sin^{d-2} \theta d\theta dr} \\ &= \frac{\int_{\mathbb{R}_+} r^{d-1} \int_{-1}^+ \chi(c, r) e(c, r) (1 - c^2)^{\frac{d-3}{2}} dc dr}{\int_{\mathbb{R}_+} r^{d-1} \int_{-1}^+ e(c, r) (1 - c^2)^{\frac{d-3}{2}} dc dr}, \end{aligned} \quad (19)$$

where $e(c, r) = \exp(rc/\sigma) \exp(-(r^2 + 1)/(2\sigma) - V(r)/\sigma)$.

Notice that thanks to (19) the coefficient c_1 does not depend upon $\Omega \in \mathbb{S}^{d-1}$

$$c_1 = \frac{\int_{\mathbb{R}^d} (v \cdot \Omega) \exp \left(-\frac{|v-\Omega|^2}{2\sigma} - \frac{V(|v|)}{\sigma} \right) dv}{\int_{\mathbb{R}^d} \exp \left(-\frac{|v-\Omega|^2}{2\sigma} - \frac{V(|v|)}{\sigma} \right) dv} = \frac{\int_{\mathbb{R}_+} r^d \int_0^\pi \cos \theta e(\cos \theta, r) \sin^{d-2} \theta d\theta dr}{\int_{\mathbb{R}_+} r^{d-1} \int_0^\pi e(\cos \theta, r) \sin^{d-2} \theta d\theta dr}.$$

In order to determine all the collision invariants, we focus on the spectral decomposition of the pressure tensor \mathcal{M}_Ω for any $\Omega \in \mathbb{S}^{d-1}$. In particular, the next lemma will imply the symmetry of the pressure tensor.

Lemma 3.2 (Spectral decomposition of \mathcal{M}_Ω) For any $\Omega \in \mathbb{S}^{d-1}$ we have $\mathcal{M}_\Omega = \sigma c_1 (I_d - \Omega \otimes \Omega)$. In particular we have $\ker(\mathcal{M}_\Omega - \sigma c_1 I_d) = (\mathbb{R}\Omega)^\perp$ and thus $\dim(\ker(\mathcal{L}_f^*)/\ker W) = \dim \ker(\mathcal{M}_\Omega - \sigma c_1 I_d) = d - 1$, cf. Lemma 3.1.

Proof. Let us consider $\{E_1, \dots, E_{d-1}\}$ an orthonormal basis of $(\mathbb{R}\Omega)^\perp$. By using the decomposition

$$v - \Omega = (\Omega \otimes \Omega)(v - \Omega) + \sum_{i=1}^{d-1} (E_i \otimes E_i)(v - \Omega) = (\Omega \otimes \Omega)(v - \Omega) + \sum_{i=1}^{d-1} (E_i \otimes E_i)v,$$

one gets

$$\mathcal{M}_\Omega = \int_{\mathbb{R}^d} \left[(\Omega \otimes \Omega)(v - \Omega) + \sum_{i=1}^{d-1} (E_i \otimes E_i)v \right] \otimes \left[\sum_{j=1}^{d-1} (E_j \otimes E_j)v \right] M_\Omega(v) dv. \quad (20)$$

We claim that the following equalities hold true

$$\int_{\mathbb{R}^d} [\Omega \cdot (v - \Omega)](E_j \cdot v) M_\Omega(v) dv = 0, \quad 1 \leq j \leq d-1, \quad (21)$$

$$\int_{\mathbb{R}^d} (E_i \cdot v)(E_j \cdot v) M_\Omega(v) dv = \delta_{ij} \int_{\mathbb{R}^d} \frac{|v|^2 - (v \cdot \Omega)^2}{d-1} M_\Omega(v) dv, \quad 1 \leq i, j \leq d-1. \quad (22)$$

Formula (21) is obtained by using the change of variable $v = (I_d - 2E_j \otimes E_j)v'$. It is easily seen that

$$\Omega \cdot (v - \Omega) = \Omega \cdot (v' - \Omega), \quad E_j \cdot v = -E_j \cdot v', \quad M_\Omega(v) = M_\Omega(v'), \quad 1 \leq j \leq d-1,$$

and therefore we have

$$\int_{\mathbb{R}^d} [\Omega \cdot (v - \Omega)](E_j \cdot v) M_\Omega(v) dv = - \int_{\mathbb{R}^d} [\Omega \cdot (v' - \Omega)](E_j \cdot v') M_\Omega(v') dv'$$

which implies (21). For the formulae (22) with $i \neq j$, we appeal to the orthogonal transformation

$$v = \mathcal{O}_{ij}v', \quad \mathcal{O}_{ij} = \Omega \otimes \Omega + \sum_{k \notin \{i,j\}} E_k \otimes E_k + E_i \otimes E_j - E_j \otimes E_i.$$

Notice that $\mathcal{O}_{ij}\xi = \xi$, for all $\xi \in (\text{span}\{E_i, E_j\})^\perp$, $\mathcal{O}_{ij}E_i = -E_j$, $\mathcal{O}_{ij}E_j = E_i$ and therefore

$$(E_i \cdot v)(E_j \cdot v) = -(E_j \cdot v')(E_i \cdot v').$$

After this change of variable we deduce that

$$\int_{\mathbb{R}^d} (E_i \cdot v)(E_j \cdot v) M_\Omega(v) dv = 0, \quad 1 \leq i, j \leq d-1, \quad i \neq j,$$

and also

$$\int_{\mathbb{R}^d} (E_i \cdot v)^2 M_\Omega(v) dv = \int_{\mathbb{R}^d} (E_j \cdot v)^2 M_\Omega(v) dv, \quad 1 \leq i, j \leq d-1.$$

Thanks to the equality $\sum_{i=1}^{d-1}(E_i \cdot v)^2 = |v|^2 - (v \cdot \Omega)^2$, one gets

$$\int_{\mathbb{R}^d} (E_i \cdot v)(E_j \cdot v)M_\Omega(v) \, dv = \delta_{ij} \int_{\mathbb{R}^d} \frac{|v|^2 - (v \cdot \Omega)^2}{d-1} M_\Omega(v) \, dv, \quad 1 \leq i, j \leq d-1.$$

Coming back to (20) we obtain

$$\mathcal{M}_\Omega = \sum_{i=1}^{d-1} \left(\int_{\mathbb{R}^d} (E_i \cdot v)^2 M_\Omega(v) \, dv \right) E_i \otimes E_i = \int_{\mathbb{R}^d} \frac{|v|^2 - (v \cdot \Omega)^2}{d-1} M_\Omega(v) \, dv (I_d - \Omega \otimes \Omega).$$

We are done if we prove that

$$\int_{\mathbb{R}^d} \frac{|v|^2 - (v \cdot \Omega)^2}{d-1} M_\Omega(v) \, dv = \sigma c_1.$$

Notice that, using (10):

$$((|v|^2 I_d - v \otimes v) \Omega) \cdot \nabla_v M_\Omega = \frac{|v|^2 - (v \cdot \Omega)^2}{\sigma} M_\Omega(v),$$

and therefore

$$\begin{aligned} \int_{\mathbb{R}^d} \frac{|v|^2 - (v \cdot \Omega)^2}{\sigma} M_\Omega(v) \, dv &= - \int_{\mathbb{R}^d} \operatorname{div}_v [(|v|^2 I_d - v \otimes v) \Omega] M_\Omega(v) \, dv \\ &= - \int_{\mathbb{R}^d} [2(v \cdot \Omega) - d(v \cdot \Omega) - (v \cdot \Omega)] M_\Omega(v) \, dv \\ &= (d-1)c_1. \end{aligned}$$

□

By Lemmas 3.1 and 3.2 the computation of the collision invariants is reduced to the resolution of (18) for any $W \in (\mathbb{R}\Omega)^\perp$. Hence, if we denote by E_1, E_2, \dots, E_{d-1} any orthonormal basis of $(\mathbb{R}\Omega)^\perp$, we obtain a set of $d-1$ collision invariants $\psi_{E_1}, \psi_{E_2}, \dots, \psi_{E_{d-1}}$ for Q associated to the equilibrium distribution f such that $E_i = W[\psi_{E_i}]$, $i = 1, \dots, d-1$. This set of collision invariants forms a basis for the $\ker(\mathcal{L}_f^*)$. In the next section we will characterize this set of collision invariants and provide an easy manner to compute them (see Lemma 4.1).

We conclude this section by showing that in our case the set of all GCIs of the operator Q coincide with the kernel of the operator \mathcal{L}_f^* .

Theorem 3.1 *Let M_Ω be an equilibrium of Q with nonvanishing orientation $\Omega \in \mathbb{S}^{d-1}$. The set of collision invariants of Q associated to M_Ω coincides with the set of the generalized collision invariants of Q associated to Ω .*

Proof.

Let $\psi = \psi(v)$ be a generalized collision invariant of Q associated to Ω . We denote by $\{e_1, \dots, e_d\}$ the canonical basis of \mathbb{R}^d . For any $f = f(v)$ satisfying $(I_d - \Omega \otimes \Omega)e_i \cdot \int_{\mathbb{R}^d} f(v)v \, dv = 0$, $1 \leq i \leq d$, we have

$$\int_{\mathbb{R}^d} f(\sigma \Delta_v \psi - \nabla_v \Phi_\Omega \cdot \nabla_v \psi) \, dv = \int_{\mathbb{R}^d} Q(f)(v) \psi(v) \, dv = 0.$$

Therefore the linear form $f \rightarrow \int_{\mathbb{R}^d} f(\sigma \Delta_v \psi - \nabla_v \Phi_\Omega \cdot \nabla_v \psi) dv$ is a linear combination of the linear forms $f \rightarrow (I_d - \Omega \otimes \Omega) e_i \cdot \int_{\mathbb{R}^d} f(v) v dv$. We deduce that there is a vector $\tilde{W} = (\tilde{W}_1, \dots, \tilde{W}_d) \in \mathbb{R}^d$ such that

$$\int_{\mathbb{R}^d} f(\sigma \Delta_v \psi - \nabla_v \Phi_\Omega \cdot \nabla_v \psi) dv + (I_d - \Omega \otimes \Omega) \tilde{W} \cdot \int_{\mathbb{R}^d} f(v) v dv = 0,$$

for any f and thus

$$\sigma \Delta_v \psi - \nabla_v \Phi_\Omega \cdot \nabla_v \psi + (I_d - \Omega \otimes \Omega) v \cdot \tilde{W} = 0,$$

implying that ψ satisfies (18) with the vector $W = (I_d - \Omega \otimes \Omega) \tilde{W} \in (\mathbb{R}\Omega)^\perp$, that is, ψ is a collision invariant of Q associated to M_Ω .

Conversely, let $\psi = \psi(v)$ be a collision invariant of Q associated to M_Ω . By Lemmas 3.1, 3.2 we know that there is $W \in (\mathbb{R}\Omega)^\perp$ such that

$$\sigma \Delta_v \psi - \nabla_v \Phi_\Omega \cdot \nabla_v \psi + v \cdot W = 0.$$

Multiplying by any function f such that $(I_d - \Omega \otimes \Omega) \int_{\mathbb{R}^d} f(v) v dv = 0$ one gets

$$\int_{\mathbb{R}^d} Q(f)(v) \psi(v) dv = \int_{\mathbb{R}^d} f(v) (\sigma \Delta_v \psi - \nabla_v \Phi_\Omega \cdot \nabla_v \psi) dv = - \int_{\mathbb{R}^d} f(v) v dv \cdot W = 0,$$

implying that ψ is a generalized collision invariant of Q associated to Ω . □

4 Identification of the collision invariants

In this section we investigate the structure of the collision invariants of Q associated to an equilibrium distribution $f = \rho M_\Omega$. By Lemmas 3.1, 3.2, we need to solve the elliptic problem

$$- \sigma \operatorname{div}_v (M_\Omega \nabla_v \psi) = (v \cdot W) M_\Omega(v), \quad v \in \mathbb{R}^d, \quad (23)$$

for any $W \in (\mathbb{R}\Omega)^\perp$. We appeal to a variational formulation by considering the continuous bilinear symmetric form $a_\Omega : H_{M_\Omega}^1 \times H_{M_\Omega}^1 \rightarrow \mathbb{R}$ defined as

$$a_\Omega(\chi, \theta) = \sigma \int_{\mathbb{R}^d} \nabla_v \chi \cdot \nabla_v \theta M_\Omega(v) dv, \quad \chi, \theta \in H_{M_\Omega}^1,$$

and the linear form $l : H_{M_\Omega}^1 \rightarrow \mathbb{R}$, $l(\theta) = \int_{\mathbb{R}^d} \theta(v) (v \cdot W) M_\Omega(v) dv$, $\theta \in H_{M_\Omega}^1$. Notice that l is well defined and bounded on $H_{M_\Omega}^1$ provided that the additional hypothesis $|v| \in L_{M_\Omega}^2$ holds true, that is

$$\int_{\mathbb{R}^d} |v|^2 \exp\left(-\frac{|v - \Omega|^2}{2\sigma} - \frac{V(|v|)}{\sigma}\right) dv < \infty.$$

The above hypothesis is obviously satisfied by the potentials $V_{\alpha,\beta}$. We say that $\psi \in H_{M_\Omega}^1$ is a variational solution of (23) if and only if

$$a_\Omega(\psi, \theta) = l(\theta) \quad \text{for any } \theta \in H_{M_\Omega}^1. \quad (24)$$

Proposition 4.1 *A necessary and sufficient condition for the existence and uniqueness of variational solution to (23) is*

$$\int_{\mathbb{R}^d} (v \cdot W) M_\Omega(v) \, dv = 0. \quad (25)$$

Proof. The necessary condition for the solvability of (23) is obtained by taking $\theta = 1$ (which belongs to $H_{M_\Omega}^1$ thanks to (11)) in (24) leading to (25). This condition is satisfied for any $W \in (\mathbb{R}\Omega)^\perp$ since we have

$$\int_{\mathbb{R}^d} (v \cdot W) M_\Omega(v) \, dv = \int_{\mathbb{R}^d} (v \cdot \Omega) M_\Omega(v) \, dv (\Omega \cdot W) = 0.$$

The condition (25) also guarantees the solvability of (23). Indeed, under the hypotheses (11), (12), the bilinear form a_Ω is coercive on the Hilbert space $\tilde{H}_{M_\Omega}^1 := \{\chi \in H_{M_\Omega}^1 : ((\chi, 1))_{M_\Omega} = 0\}$, i.e. we have:

$$a_\Omega(\chi, \chi) \geq \frac{\min\{\sigma, \lambda_\Omega\}}{2} \|\chi\|_{M_\Omega}^2, \quad \chi \in \tilde{H}_{M_\Omega}^1.$$

Applying the Lax-Milgram lemma to (24) with $\psi, \theta \in \tilde{H}_{M_\Omega}^1$ yields a unique function $\psi \in \tilde{H}_{M_\Omega}^1$ such that

$$a_\Omega(\psi, \tilde{\theta}) = l(\tilde{\theta}) \quad \text{for any } \tilde{\theta} \in \tilde{H}_{M_\Omega}^1. \quad (26)$$

Actually, the compatibility condition $l(1) = 0$ allows us to extend (26) to $H_{M_\Omega}^1$. This follows by applying (26) with $\tilde{\theta} = \theta - ((\theta, 1))_{M_\Omega}$, for $\theta \in H_{M_\Omega}^1$. Moreover, the uniqueness of the solution for the problem on $\tilde{H}_{M_\Omega}^1$ implies the uniqueness, up to a constant, of the solution for the problem on $H_{M_\Omega}^1$. \square

As observed in (13), the fluid equations for ρ and Ω will follow by appealing to the collision invariants associated to the orientation $\Omega \in \mathbb{S}^{d-1}$, for any $W \in (\mathbb{R}\Omega)^\perp$. When $W = 0$, the solutions of (23) are all the constants, and we obtain the particle number balance (16). Consider now $W \in (\mathbb{R}\Omega)^\perp \setminus \{0\}$ and ψ_W a solution of (23). Obviously we have $\psi_W = \tilde{\psi}_W + \int_{\mathbb{R}^d} \psi_W(v) M_\Omega(v) \, dv$, where $\tilde{\psi}_W$ is the unique solution of (23) in $\tilde{H}_{M_\Omega}^1$. It is easily seen, thanks to (16) and the linearity of (15), that the balances corresponding to ψ_W and $\tilde{\psi}_W$ are equivalent. Therefore for any $W \in (\mathbb{R}\Omega)^\perp$ it is enough to consider only the solution of (23) in $\tilde{H}_{M_\Omega}^1$. From now on, for any $W \in (\mathbb{R}\Omega)^\perp$, we denote by ψ_W the unique variational solution of (23) verifying $\int_{\mathbb{R}^d} \psi_W(v) M_\Omega(v) \, dv = 0$. The structure of the solutions $\psi_W, W \in (\mathbb{R}\Omega)^\perp \setminus \{0\}$ comes by the symmetry of the equilibrium M_Ω . Analyzing the rotations leaving invariant the vector Ω , we prove as in [14] the following result.

Proposition 4.2 *Consider $W \in (\mathbb{R}\Omega)^\perp \setminus \{0\}$. For any orthogonal transformation \mathcal{O} of \mathbb{R}^d leaving Ω invariant, that is $\mathcal{O}\Omega = \Omega$, we have*

$$\psi_W(\mathcal{O}v) = \psi_{{}^t\mathcal{O}W}(v), \quad v \in \mathbb{R}^d,$$

where ${}^t\mathcal{O}$ denotes the transpose of the matrix \mathcal{O} .

Proof. First of all notice that ${}^t\mathcal{O}W \in (\mathbb{R}\Omega)^\perp \setminus \{0\}$. We know that ψ_W is the minimum point of the functional

$$J_W(z) = \frac{\sigma}{2} \int_{\mathbb{R}^d} |\nabla_v z|^2 M_\Omega(v) \, dv - \int_{\mathbb{R}^d} (v \cdot W) z(v) M_\Omega(v) \, dv, \quad z \in \tilde{H}_{M_\Omega}^1.$$

It is easily seen that, for any orthogonal transformation \mathcal{O} of \mathbb{R}^d leaving the orientation Ω invariant, and any function $z \in \tilde{H}_{M_\Omega}^1$, we have, by defining $z_\mathcal{O} := z \circ \mathcal{O} \in \tilde{H}_{M_\Omega}^1$

$$M_\Omega \circ \mathcal{O} = M_\Omega, \quad \nabla z_\mathcal{O} = {}^t\mathcal{O}(\nabla z) \circ \mathcal{O}.$$

Moreover, we obtain with the change of variables $v' = \mathcal{O}v$ and using that $M_\Omega(v) = M_\Omega(v')$:

$$\begin{aligned} J_{{}^t\mathcal{O}W}(z_\mathcal{O}) &= \frac{\sigma}{2} \int_{\mathbb{R}^d} |{}^t\mathcal{O}(\nabla z)(\mathcal{O}v)|^2 M_\Omega(v) \, dv - \int_{\mathbb{R}^d} (v \cdot {}^t\mathcal{O}W) z(\mathcal{O}v) M_\Omega(v) \, dv \\ &= \frac{\sigma}{2} \int_{\mathbb{R}^d} |(\nabla z)(v')|^2 M_\Omega(v') \, dv' - \int_{\mathbb{R}^d} (v' \cdot W) z(v') M_\Omega(v') \, dv' \\ &= J_W(z). \end{aligned}$$

Finally, we deduce that

$$\psi_W \circ \mathcal{O} \in \tilde{H}_{M_\Omega}^1, \quad J_{{}^t\mathcal{O}W}(\psi_W \circ \mathcal{O}) = J_W(\psi_W) \leq J_W(z \circ {}^t\mathcal{O}) = J_{{}^t\mathcal{O}W}(z),$$

for any $z \in \tilde{H}_{M_\Omega}^1$, implying that $\psi_W \circ \mathcal{O} = \psi_{{}^t\mathcal{O}W}$. \square

The computation of the collision invariants $\{\psi_W : W \in (\mathbb{R}\Omega)^\perp \setminus \{0\}\}$ can be reduced to the computation of one scalar function. For any orthonormal basis $\{E_1, \dots, E_{d-1}\}$ of $(\mathbb{R}\Omega)^\perp$ we define the vector field $F = \sum_{i=1}^{d-1} \psi_{E_i} E_i$. This vector field does not depend upon the basis $\{E_1, \dots, E_{d-1}\}$ and has the following properties, see [14].

Lemma 4.1 *The vector field F does not depend on the orthonormal basis $\{E_1, \dots, E_{d-1}\}$ of $(\mathbb{R}\Omega)^\perp$ and for any orthogonal transformation \mathcal{O} of \mathbb{R}^d , preserving Ω , we have $F \circ \mathcal{O} = \mathcal{O}F$. There is a function χ such that*

$$F(v) = \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) \frac{(I_d - \Omega \otimes \Omega)(v)}{\sqrt{|v|^2 - (v \cdot \Omega)^2}}, \quad v \in \mathbb{R}^d \setminus (\mathbb{R}\Omega),$$

and thus, for any $i \in \{1, \dots, d-1\}$, we have

$$\psi_{E_i}(v) = F(v) \cdot E_i = \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) \frac{v \cdot E_i}{\sqrt{|v|^2 - (v \cdot \Omega)^2}}, \quad v \in \mathbb{R}^d \setminus (\mathbb{R}\Omega). \quad (27)$$

Proof. Let $\{F_1, \dots, F_{d-1}\}$ be another orthonormal basis of $(\mathbb{R}\Omega)^\perp$. The following identities hold

$$E_1 \otimes E_1 + \dots + E_{d-1} \otimes E_{d-1} + \Omega \otimes \Omega = I_d, \quad F_1 \otimes F_1 + \dots + F_{d-1} \otimes F_{d-1} + \Omega \otimes \Omega = I_d,$$

and therefore

$$\begin{aligned}
\sum_{i=1}^{d-1} \psi_{E_i} E_i &= \sum_{i=1}^{d-1} \psi_{\sum_{j=1}^{d-1} (E_i \cdot F_j) F_j} E_i \\
&= \sum_{i=1}^{d-1} \sum_{j=1}^{d-1} (E_i \cdot F_j) \psi_{F_j} E_i \\
&= \sum_{j=1}^{d-1} \psi_{F_j} \sum_{i=1}^{d-1} (E_i \cdot F_j) E_i \\
&= \sum_{j=1}^{d-1} \psi_{F_j} F_j.
\end{aligned}$$

For any orthogonal transformation of \mathbb{R}^d such that $\mathcal{O}\Omega = \Omega$ we obtain thanks to Proposition 4.2

$$F \circ \mathcal{O} = \sum_{i=1}^{d-1} (\psi_{E_i} \circ \mathcal{O}) E_i = \sum_{i=1}^{d-1} \psi_{{}^t\mathcal{O}E_i} E_i = \mathcal{O} \sum_{i=1}^{d-1} \psi_{{}^t\mathcal{O}E_i} {}^t\mathcal{O}E_i = \mathcal{O}F,$$

where, the last equality holds true since $\{{}^t\mathcal{O}E_1, \dots, {}^t\mathcal{O}E_{d-1}\}$ is an orthonormal basis of $(\mathbb{R}\Omega)^\perp$. Let $v \in \mathbb{R}^d \setminus (\mathbb{R}\Omega)$ and consider

$$E(v) = \frac{(Id - \Omega \otimes \Omega)v}{\sqrt{|v|^2 - (\Omega \cdot v)^2}}.$$

Notice that $E \cdot \Omega = 0, |E| = 1$. When $d = 2$, since the vector $F(v)$ is orthogonal to Ω , there exists a function $\Lambda = \Lambda(v)$ such that

$$F(v) = \Lambda(v)E = \Lambda(v) \frac{(I_2 - \Omega \otimes \Omega)v}{\sqrt{|v|^2 - (\Omega \cdot v)^2}}, \quad v \in \mathbb{R}^2 \setminus (\mathbb{R}\Omega).$$

If $d \geq 3$, let us denote by ${}^\perp E$, any unitary vector orthogonal to E and Ω . Introducing the orthogonal matrix $\mathcal{O} = I_d - 2 {}^\perp E \otimes {}^\perp E$ (which leaves Ω invariant), we obtain $F \circ \mathcal{O} = \mathcal{O}F$. Observe that

$$0 = {}^\perp E \cdot E = {}^\perp E \cdot \frac{v - (v \cdot \Omega)\Omega}{\sqrt{|v|^2 - (v \cdot \Omega)^2}} = \frac{{}^\perp E \cdot v}{\sqrt{|v|^2 - (v \cdot \Omega)^2}}, \quad \mathcal{O}v = v,$$

and thus

$$F(v) = F(\mathcal{O}v) = \mathcal{O}F(v) = (I_d - 2 {}^\perp E \otimes {}^\perp E)F(v) = F(v) - 2({}^\perp E \cdot F(v)) {}^\perp E,$$

from which it follows that ${}^\perp E \cdot F(v) = 0$, for any vector ${}^\perp E$ orthogonal to E and Ω . Hence, there exists a function $\Lambda(v)$ such that

$$F(v) = \Lambda(v)E(v) = \Lambda(v) \frac{(Id - \Omega \otimes \Omega)v}{\sqrt{|v|^2 - (v \cdot \Omega)^2}}, \quad v \in \mathbb{R}^d \setminus (\mathbb{R}\Omega).$$

We will show that $\Lambda(v)$ depends only on $v \cdot \Omega/|v|$ and $|v|$. Indeed, for any $d \geq 2$, and any orthogonal transformation \mathcal{O} such that $\mathcal{O}\Omega = \Omega$ we have $F(\mathcal{O}v) = \mathcal{O}F(v)$, $E(\mathcal{O}v) = \mathcal{O}E(v)$ because

$$(I_d - \Omega \otimes \Omega)\mathcal{O}v = \mathcal{O}v - (\Omega \cdot \mathcal{O}v)\Omega = \mathcal{O}(I_d - \Omega \otimes \Omega)v,$$

$$\sqrt{|\mathcal{O}v|^2 - (\mathcal{O}v \cdot \Omega)^2} = \sqrt{|v|^2 - (v \cdot \Omega)^2},$$

implying that $\Lambda(\mathcal{O}v) = \Lambda(v)$, for any $v \in \mathbb{R}^d \setminus (\mathbb{R}\Omega)$. We are done if we prove that $\Lambda(v) = \Lambda(v')$ for any $v, v' \in \mathbb{R}^d \setminus (\mathbb{R}\Omega)$ such that $v \cdot \Omega/|v| = v' \cdot \Omega/|v'|$, $|v| = |v'|$, $v \neq v'$. It is enough to consider the rotation \mathcal{O} such that

$$\mathcal{O}E = E', \quad (\mathcal{O} - I_d)_{\text{span}\{E, E'\}^\perp} = 0, \quad E = \frac{(I_d - \Omega \otimes \Omega)v}{\sqrt{|v|^2 - (v \cdot \Omega)^2}}, \quad E' = \frac{(I_d - \Omega \otimes \Omega)v'}{\sqrt{|v'|^2 - (v' \cdot \Omega)^2}}.$$

The equality $\mathcal{O}E = E'$ implies that $\mathcal{O}v = v'$ and therefore $\Lambda(v') = \Lambda(\mathcal{O}v) = \Lambda(v)$, showing that there exists a function χ such that $\Lambda(v) = \chi(v \cdot \Omega/|v|, |v|)$, $v \in \mathbb{R}^d \setminus \{0\}$. \square

In the last part of this section we concentrate on the elliptic problem satisfied by the function $(c, r) \rightarrow \chi(c, r)$ introduced in Lemma 4.1. Even if ψ_{E_i} are eventually singular on $\mathbb{R}\Omega$, it will be no difficulty to define a Hilbert space on which solving for the profile χ . We again proceed using the minimization of quadratic functionals.

Proposition 4.3 *The function χ constructed in Lemma 4.1 solves the problem*

$$\begin{aligned} -\sigma \partial_c \{r^{d-3}(1-c^2)^{\frac{d-1}{2}} e(c, r) \partial_c \chi\} - \sigma \partial_r \{r^{d-1}(1-c^2)^{\frac{d-3}{2}} e(c, r) \partial_r \chi\} + \sigma(d-2)r^{d-3}(1-c^2)^{\frac{d-5}{2}} e\chi \\ = r^d(1-c^2)^{\frac{d-2}{2}} e(c, r), \quad (c, r) \in]-1, +1[\times]0, \infty[. \end{aligned} \quad (28)$$

Proof. For any $i \in \{1, \dots, d-1\}$, let us consider $\psi_{E_i, h}(v) = h(v \cdot \Omega/|v|, |v|) \frac{v \cdot E_i}{\sqrt{|v|^2 - (v \cdot \Omega)^2}}$ where $v \rightarrow h(v \cdot \Omega/|v|, |v|)$ is a function such that $\psi_{E_i, h} \in H_{M_\Omega}^1$. Observe that if $h = \chi$, then $\psi_{E_i, h}$ coincides with ψ_{E_i} . Note that generally $\psi_{E_i, h}$ are not collision invariants, but perturbations of them, corresponding to profiles h . In this way, the minimization problem (29) will lead to a minimization problem on h , whose solution will be χ . Notice that once that $\psi_{E_i, h} \in H_{M_\Omega}^1$, then $\int_{\mathbb{R}^d} \psi_{E_i, h} M_\Omega(v) dv = 0$, saying that $\psi_{E_i, h} \in \tilde{H}_{M_\Omega}^1$. We know that ψ_{E_i} is the minimum point of J_{E_i} on $\tilde{H}_{M_\Omega}^1$ and therefore

$$J_{E_i}(\psi_{E_i}) \leq J_{E_i}(\psi_{E_i, h}). \quad (29)$$

A straightforward computation shows that

$$\begin{aligned} \nabla_v \psi_{E_i, h} &= \frac{v \cdot E_i}{\sqrt{|v|^2 - (v \cdot \Omega)^2}} \left[\partial_c h \frac{I_d - \frac{v \otimes v}{|v|^2}}{|v|} \Omega + \partial_r h \frac{v}{|v|} \right] \\ &+ h \left(\frac{v \cdot \Omega}{|v|}, |v| \right) \left[I_d - \frac{(v - (v \cdot \Omega)\Omega) \otimes v}{|v|^2 - (v \cdot \Omega)^2} \right] \frac{E_i}{\sqrt{|v|^2 - (v \cdot \Omega)^2}}, \end{aligned}$$

and

$$|\nabla_v \psi_{E_i, h}|^2 = \frac{(v \cdot E_i)^2}{|v|^4} (\partial_c h)^2 + \frac{(v \cdot E_i)^2}{|v|^2 - (v \cdot \Omega)^2} (\partial_r h)^2 + \frac{|v|^2 - (v \cdot \Omega)^2 - (v \cdot E_i)^2}{(|v|^2 - (v \cdot \Omega)^2)^2} h^2 \left(\frac{v \cdot \Omega}{|v|}, |v| \right).$$

The condition $\psi_{E_i, h} \in H_{M_\Omega}^1$ writes

$$\int_{\mathbb{R}^d} (\psi_{E_i, h})^2 M_\Omega(v) \, dv < \infty, \quad \int_{\mathbb{R}^d} |\nabla_v \psi_{E_i, h}|^2 M_\Omega(v) \, dv < \infty,$$

which is equivalent, thanks to the Poincaré inequality (12) to

$$\int_{\mathbb{R}^d} |\nabla_v \psi_{E_i, h}|^2 M_\Omega(v) \, dv < \infty.$$

Based on formula (19), we have

$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla_v \psi_{E_i, h}|^2 M_\Omega(v) \, dv &= \int_{\mathbb{R}^d} \left[\frac{|v|^2 - (v \cdot \Omega)^2}{(d-1)|v|^4} (\partial_c h)^2 + \frac{(\partial_r h)^2}{d-1} + \frac{(d-2)h^2}{(d-1)(|v|^2 - (v \cdot \Omega)^2)} \right] M_\Omega \, dv \\ &= \frac{\int_{\mathbb{R}_+} r^{d-1} \int_{-1}^{+1} \left[\frac{1-c^2}{r^2} (\partial_c h)^2 + (\partial_r h)^2 + \frac{(d-2)h^2}{r^2(1-c^2)} \right] e(c, r) (1-c^2)^{\frac{d-3}{2}} \, dc \, dr}{(d-1) \int_{\mathbb{R}_+} r^{d-1} \int_{-1}^{+1} e(c, r) (1-c^2)^{\frac{d-3}{2}} \, dc \, dr} \end{aligned}$$

and therefore we consider the Hilbert space $H_d = \{h :]-1, +1[\times]0, \infty[\rightarrow \mathbb{R}, \|h\|_d^2 < \infty\}$, endowed with the scalar product

$$(g, h)_d = \int_{\mathbb{R}_+} r^{d-1} \int_{-1}^{+1} \left[\frac{1-c^2}{r^2} \partial_c g \partial_c h + \partial_r g \partial_r h + \frac{(d-2)gh}{r^2(1-c^2)} \right] e(c, r) (1-c^2)^{\frac{d-3}{2}} \, dc \, dr$$

for g and h in H_d and the norm given by

$$\|h\|_d = \sqrt{(h, h)_d}.$$

The expression $J_{E_i}(\psi_{E_i, h})$ writes as functional of h

$$\begin{aligned} J_{E_i}(\psi_{E_i, h}) &= \frac{\sigma}{2} \int_{\mathbb{R}^d} |\nabla_v \psi_{E_i, h}|^2 M_\Omega(v) \, dv - \int_{\mathbb{R}^d} \frac{(v \cdot E_i)^2 h \left(\frac{v \cdot \Omega}{|v|}, |v| \right)}{\sqrt{|v|^2 - (v \cdot \Omega)^2}} M_\Omega(v) \, dv \\ &= \frac{\sigma}{2} \int_{\mathbb{R}^d} |\nabla_v \psi_{E_i, h}|^2 M_\Omega(v) \, dv - \int_{\mathbb{R}^d} \frac{h \left(\frac{v \cdot \Omega}{|v|}, |v| \right) \sqrt{|v|^2 - (v \cdot \Omega)^2}}{d-1} M_\Omega(v) \, dv \\ &= \frac{J(h)}{(d-1) \int_{\mathbb{R}_+} r^{d-1} \int_{-1}^{+1} e(c, r) (1-c^2)^{\frac{d-3}{2}} \, dc \, dr}, \end{aligned}$$

where

$$\begin{aligned} J(h) &= \frac{\sigma}{2} \int_{\mathbb{R}_+} r^{d-1} \int_{-1}^{+1} \left[\frac{1-c^2}{r^2} (\partial_c h)^2 + (\partial_r h)^2 + \frac{(d-2)h^2}{r^2(1-c^2)} \right] e(c, r) (1-c^2)^{\frac{d-3}{2}} \, dc \, dr \\ &\quad - \int_{\mathbb{R}_+} r^{d-1} \int_{-1}^{+1} h(c, r) r \sqrt{1-c^2} e(c, r) (1-c^2)^{\frac{d-3}{2}} \, dc \, dr. \end{aligned}$$

Coming back to (29) and using (27), we deduce that

$$\chi \in H_d \quad \text{and} \quad J(\chi) \leq J(h) \quad \text{for any } h \in H_d.$$

Therefore, by the Lax-Milgram lemma, we deduce that χ solves the problem (28). \square

5 Hydrodynamic equations

After identifying the collision invariants, we determine the fluid equations satisfied by the macroscopic quantities entering the dominant particle density $f(t, x, v) = \rho(t, x)M_{\Omega(t, x)}(v)$. As seen before the balance for the particle density follows thanks to the collision invariant $\psi = 1$. The other balances follow by appealing to the vector field F (cf. Lemma 4.1) and the details are given in Sect. 5.1.

5.1 Proof of Theorem 1.1

Applying (13) with $\psi = 1$ leads to (16). For any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$ we consider the vector field

$$v \rightarrow F(t, x, v) = \chi\left(\frac{v \cdot \Omega(t, x)}{|v|}, |v|\right) \frac{(I_d - \Omega(t, x) \otimes \Omega(t, x))v}{\sqrt{|v|^2 - (v \cdot \Omega(t, x))^2}}.$$

By the definition of $F(t, x, \cdot)$, we know that

$$\mathcal{L}_{f(t, x, \cdot)}^* F(t, x, \cdot) = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d,$$

and therefore (13) implies

$$\int_{\mathbb{R}^d} \partial_t f F(t, x, v) dv + \int_{\mathbb{R}^d} v \cdot \nabla_x f F(t, x, v) dv = 0, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^d. \quad (30)$$

It remains to compute $\int_{\mathbb{R}^d} \partial_t f F(t, x, v) dv$ and $\int_{\mathbb{R}^d} v \cdot \nabla_x f F(t, x, v) dv$ in terms of $\rho(t, x)$ and $\Omega(t, x)$. By a direct computation we obtain

$$\partial_t f = \partial_t \rho M_{\Omega} + \rho \frac{M_{\Omega}}{\sigma} (v - \Omega) \cdot \partial_t \Omega,$$

implying, thanks to the equalities $\partial_t \Omega \cdot \Omega = 0$ and $v \cdot \partial_t \Omega = P_f v \cdot \partial_t \Omega$,

$$\begin{aligned} \int_{\mathbb{R}^d} \partial_t f F dv &= \int_{\mathbb{R}^d} \left(\partial_t \rho + \frac{\rho}{\sigma} (v - \Omega) \cdot \partial_t \Omega \right) \chi\left(\frac{v \cdot \Omega}{|v|}, |v|\right) M_{\Omega}(v) \frac{P_f v}{|P_f v|} dv \\ &= \partial_t \rho \int_{\mathbb{R}^d} \chi\left(\frac{v \cdot \Omega}{|v|}, |v|\right) M_{\Omega}(v) \frac{P_f v}{|P_f v|} dv + \frac{\rho}{\sigma} \int_{\mathbb{R}^d} \chi\left(\frac{v \cdot \Omega}{|v|}, |v|\right) M_{\Omega}(v) \frac{P_f v \otimes P_f v}{|P_f v|} dv \partial_t \Omega. \end{aligned} \quad (31)$$

It is an easy exercise to show that the integral $\int_{\mathbb{R}^d} \chi\left(\frac{v \cdot \Omega}{|v|}, |v|\right) M_{\Omega}(v) \frac{P_f v}{|P_f v|} dv$ vanishes and that the following relationship holds

$$\begin{aligned} \int_{\mathbb{R}^d} \chi\left(\frac{v \cdot \Omega}{|v|}, |v|\right) M_{\Omega}(v) \frac{P_f v \otimes P_f v}{|P_f v|} dv \\ = \int_{\mathbb{R}^d} \chi\left(\frac{v \cdot \Omega}{|v|}, |v|\right) \frac{\sqrt{|v|^2 - (v \cdot \Omega)^2}}{d-1} M_{\Omega}(v) dv (I_d - \Omega \otimes \Omega). \end{aligned}$$

Therefore, by taking into account that $\partial_t \Omega \cdot \Omega = 0$, the equality (31) becomes

$$\int_{\mathbb{R}^d} \partial_t f F dv = \tilde{c}_1 \frac{\rho}{\sigma} \partial_t \Omega, \quad \tilde{c}_1 = \int_{\mathbb{R}^d} \chi\left(\frac{v \cdot \Omega}{|v|}, |v|\right) \frac{\sqrt{|v|^2 - (v \cdot \Omega)^2}}{d-1} M_{\Omega}(v) dv. \quad (32)$$

Similarly we write (for any smooth vector field $\xi(x)$, the notation $\partial_x \xi$ stands for the Jacobian matrix of ξ , i.e. $(\partial_x \xi)_{i,j} = \partial_{x_j} \xi_i$)

$$v \cdot \nabla_x f = (v \cdot \nabla_x \rho) M_\Omega + \frac{\rho}{\sigma} v \cdot ({}^t \partial_x \Omega (v - \Omega)) M_\Omega,$$

implying

$$\begin{aligned} \int_{\mathbb{R}^d} (v \cdot \nabla_x f) F \, dv &= \int_{\mathbb{R}^d} \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) M_\Omega(v) \frac{P_f v \otimes P_f v}{|P_f v|} \, dv \nabla_x \rho \\ &+ \frac{\rho}{\sigma} \int_{\mathbb{R}^d} (v \cdot \Omega) \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) M_\Omega(v) \frac{P_f v \otimes P_f v}{|P_f v|} \, dv (\partial_x \Omega) \Omega \\ &= \tilde{c}_1 (I_d - \Omega \otimes \Omega) \nabla_x \rho + \frac{\rho}{\sigma} \tilde{c}_2 \partial_x \Omega \Omega, \end{aligned} \quad (33)$$

where

$$\tilde{c}_2 = \int_{\mathbb{R}^d} (v \cdot \Omega) \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) \frac{\sqrt{|v|^2 - (v \cdot \Omega)^2}}{d-1} M_\Omega(v) \, dv.$$

Notice that in the above computations we have used $({}^t \partial_x \Omega) \Omega = 0$ and

$$\int_{\mathbb{R}^d} (v \cdot E_i)(v \cdot E_j)(v \cdot E_k) \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) M_\Omega(v) \, dv = 0,$$

for any $i, j, k \in \{1, \dots, d-1\}$. Combining (30), (32) and (33) yields

$$\tilde{c}_1 \frac{\rho}{\sigma} \partial_t \Omega + \frac{\rho}{\sigma} \tilde{c}_2 \partial_x \Omega \Omega + \tilde{c}_1 (I_d - \Omega \otimes \Omega) \nabla_x \rho = 0,$$

or equivalently

$$\partial_t \Omega + c_2 \partial_x \Omega \Omega + \sigma (I_d - \Omega \otimes \Omega) \frac{\nabla_x \rho}{\rho} = 0, \quad (34)$$

where

$$\begin{aligned} c_2 &= \frac{\tilde{c}_2}{\tilde{c}_1} = \frac{\int_{\mathbb{R}^d} (v \cdot \Omega) \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) \sqrt{|v|^2 - (v \cdot \Omega)^2} M_\Omega(v) \, dv}{\int_{\mathbb{R}^d} \chi \left(\frac{v \cdot \Omega}{|v|}, |v| \right) \sqrt{|v|^2 - (v \cdot \Omega)^2} M_\Omega(v) \, dv} \\ &= \frac{\int_{\mathbb{R}_+} r^{d+1} \int_0^\pi \cos \theta \chi(\cos \theta, r) e(\cos \theta, r) \sin^{d-1} \theta \, d\theta \, dr}{\int_{\mathbb{R}_+} r^d \int_0^\pi \chi(\cos \theta, r) e(\cos \theta, r) \sin^{d-1} \theta \, d\theta \, dr}. \end{aligned}$$

5.2 Properties of the hydrodynamic equations

Let us start by noticing that the system (5)-(6) is hyperbolic as a consequence of Theorem 4.1 in [42]. On the other hand, the orientation balance equation (34) propagates the constraint $|\Omega| = 1$. Indeed, let $\Omega = \Omega(t, x)$ be a smooth solution of (34), satisfying $|\Omega(0, \cdot)| = 1$. Multiplying by $\Omega(t, x)$ we obtain

$$\frac{1}{2} \partial_t |\Omega|^2 + \frac{c_2}{2} (\Omega(t, x) \cdot \nabla_x) |\Omega|^2 = 0,$$

implying that $|\Omega|$ is constant along the characteristics of the vector field $c_2\Omega(t, x) \cdot \nabla_x$ and thus $|\Omega(t, \cdot)| = |\Omega(0, \cdot)| = 1$, for all $t \geq 0$.

The rescaled equation (3) can be considered as an intermediate model between the equations introduced in [2] and [14] when there are no phase transitions. In [14], the authors considered a strong relaxation towards the 'terminal speed' (or cruise speed). Whereas in [2] the authors do not impose a penalization on the self-propelled/friction term. Our result could be applied to obtain the results in [14] without resorting to measures supported on the sphere by doing a double passage to the limit. First, passing to the limit $\varepsilon \rightarrow 0$ in (3), taking $V = V_{\alpha, \beta}$, we obtain (5)-(6). Afterwards, we rescale V to $\lambda\tilde{V}$ in the system (5)-(6) and study the limit when $\lambda \rightarrow \infty$. This amounts to study the behavior of the coefficients

$$c_{1, \lambda} = \frac{\int_{\mathbb{R}_+} r^d \int_0^\pi \cos \theta e_\lambda(\cos \theta, r) \sin^{d-2} \theta \, d\theta \, dr}{\int_{\mathbb{R}_+} r^{d-1} \int_0^\pi e_\lambda(\cos \theta, r) \sin^{d-2} \theta \, d\theta \, dr},$$

and

$$c_{2, \lambda} = \frac{\int_{\mathbb{R}_+} r^{d+1} \int_0^\pi \cos \theta \chi_\lambda(\cos \theta, r) e_\lambda(\cos \theta, r) \sin^{d-1} \theta \, d\theta \, dr}{\int_{\mathbb{R}_+} r^d \int_0^\pi \chi_\lambda(\cos \theta, r) e_\lambda(\cos \theta, r) \sin^{d-1} \theta \, d\theta \, dr},$$

when $\lambda \rightarrow \infty$, where the function χ_λ solves the elliptic problem

$$\begin{aligned} & -\sigma \partial_c \left[r^{d-3} (1-c^2)^{\frac{d-1}{2}} e_\lambda(c, r) \partial_c \chi_\lambda \right] \\ & -\sigma \partial_r \left[r^{d-1} (1-c^2)^{\frac{d-3}{2}} e_\lambda(c, r) \partial_r \chi_\lambda \right] + \sigma (d-2) r^{d-3} (1-c^2)^{\frac{d-5}{2}} e_\lambda \chi_\lambda \\ & = r^d (1-c^2)^{\frac{d-2}{2}} e_\lambda(c, r), \end{aligned}$$

and $e_\lambda(c, r) = \exp(rc/\sigma) \exp(-(r^2 + 1)/(2\sigma) - \lambda\tilde{V}(r)/\sigma)$.

In order to analyse the asymptotic behavior of $c_{1, \lambda}$ we introduce the following result.

Lemma 5.1 *Let $\varphi \in C^2((0, \infty); \mathbb{R})$ and $g \in C^0((0, \infty) \times (0, \pi); \mathbb{R})$. Let us assume that*

- i) $\int_{\mathbb{R}_+} \int_0^\pi \exp(\lambda\varphi(r)) |g(r, \theta)| \, d\theta \, dr < \infty$,
- ii) *The function φ has a unique global maximum at an interior point r_0 ,*
- iii) $\int_0^\pi g(r_0, \theta) \, d\theta \neq 0$.

Then the function $G(\lambda)$ defined as

$$G(\lambda) = \int_{\mathbb{R}_+} \int_0^\pi \exp(\lambda\varphi(r)) g(r, \theta) \, d\theta \, dr,$$

has the the following asymptotic behavior

$$G(\lambda) \sim \sqrt{\frac{2\pi}{|\varphi''(r_0)|}} \frac{\exp(\lambda\varphi(r_0))}{\sqrt{\lambda}} \int_0^\pi g(r_0, \theta) \, d\theta,$$

as $\lambda \rightarrow \infty$.

The proof of this result is a direct application of the Laplace method, see for instance [53]. As an immediate consequence of Lemma 5.1 we obtain

$$\lim_{\lambda \rightarrow \infty} c_{1,\lambda} = \frac{r_0 \int_0^\pi \cos \theta \exp(r_0 \cos \theta / \sigma) \sin^{d-2} \theta \, d\theta}{\int_0^\pi \exp(r_0 \cos \theta / \sigma) \sin^{d-2} \theta \, d\theta},$$

where r_0 is the minimum of the potential function $V_{\alpha,\beta}(r)$. Let us note that the asymptotic study of the coefficient $c_{1,\lambda}$ when $\lambda \rightarrow \infty$ can also be performed using Lemma 5.1 for more general potentials than $V_{\alpha,\beta}(r)$. In particular, we could also consider smooth potentials $V(|v|)$ having a unique global minimum r_0 such that $V'(r) < 0$, for $0 < r < r_0$, and $V'(r) > 0$, for $r > r_0$. On the other hand, the asymptotic study of $c_{2,\lambda}$ could be addressed following similar techniques as in [42], however, we do not dwell upon this matter here and leave it for a future work.

6 Conclusion

In this paper, we have considered a flocking model consisting of a modified Cucker-Smale involving noise, alignment and self-propulsion. We have investigated its macroscopic limit when the time and space variables are rescaled to the macroscopic scale. In this limit, the velocity distribution converges to an equilibrium which depends on the local density and local mean orientation of the particles. The density and mean orientation evolve in space and time according to a hyperbolic system of equations named the Self-Organized Hydrodynamics. This system is akin to the system of isothermal compressible gas dynamics with the important difference that the velocity is a vector of unit norm. It is structurally identical with that obtained from the Vicsek dynamics, with differences only in the values of the coefficients. An interesting open problem would be to investigate whether the coefficients of the two models coincide in the limit where the Cucker-Smale model converges to the Vicsek model. Future work will investigate different variants in the Cucker-Smale model, possibly involving phase transitions (which have been avoided in the present work by an appropriate expression of the alignment force).

Acknowledgments

PAS and PD acknowledge support by the Engineering and Physical Sciences Research Council (EPSRC) under grants no. EP/M006883/1 and EP/P013651/1. This work was accomplished during the visit of MB to Imperial College London with an ICL-CNRS Fellowship. JAC was partially supported by the EPSRC grant number EP/P031587/1. PD also acknowledges support by the Royal Society and the Wolfson Foundation through a Royal Society Wolfson Research Merit Award no. WM130048 and by the National Science Foundation (NSF) under grant no. RNMS11-07444 (KI-Net). PD is on leave from CNRS, Institut de Mathématiques de Toulouse, France. PAS and PD would like to thank the Isaac Newton Institute for Mathematical Sciences for support and hospitality during the programme ‘‘Growth form and self-organisation’’ when part of this work was carried over which was supported by: EPSRC grant numbers

EP/K032208/1 and EP/R014604/1.

Data statement: no new data were collected in the course of this research.

References

- [1] M. Ballerini et al, Interaction ruling animal collective behavior depends on topological rather than metric distance: evidence from a field study, *Proc. Natl. Acad. Sci. USA* 105(2008) 1232-1237.
- [2] A. B. T. Barbaro and P. Degond, Phase transition and diffusion among socially interacting self-propelled agents, *Discrete Contin. Dyn. Syst. Ser. B*, 19 (2014), pp. 1249–1278.
- [3] A. B. T. Barbaro, J.A. Cañizo, J.A. Carrillo, P. Degond, Phase transitions in a kinetic flocking model of Cucker-Smale type, *Multiscale Model. Simul.* 14 (2016) 1063–1088.
- [4] C. Bardos, F. Golse, D. Levermore, Fluid dynamic limits of kinetic equations. I. Formal derivations, *J. Stat. Phys.* 63(1991) 323-344.
- [5] C. Bardos, F. Golse, D. Levermore, Fluid dynamic limits of kinetic equations. II. Convergence proofs for the Boltzmann equation, *Comm. Pure Appl. Math.* 46(1993) 667-753.
- [6] E. Bertin, M. Droz and G. Grégoire, Hydrodynamic equations for self-propelled particles: Microscopic derivation and stability analysis, *J. Phys. A, Math. Theor.* 42(2009) 445001.
- [7] F. Bolley, J.A. Cañizo, J.A. Carrillo, Stochastic Mean-Field Limit: Non-Lipschitz Forces & Swarming, *Math. Models Methods Appl. Sci.* 21(2011) 2179-2210.
- [8] F. Bolley, J.A. Cañizo, J.A. Carrillo, Mean-field limit for the stochastic Vicsek model, *Appl. Math. Lett.* 25(2012) 339-343.
- [9] V. Bonnaillie-Noël, J.A. Carrillo, T. Goudon, G.A. Pavliotis, Efficient numerical calculation of drift and diffusion coefficients in the diffusion approximation of kinetic equations, *IMA J. Numer. Anal.* (2016) 1-34.
- [10] M. Bostan, The Vlasov-Poisson system with strong external magnetic field. Finite Larmor radius regime, *Asymptot. Anal.* 61(2009) 91-123.
- [11] M. Bostan, Transport equations with disparate advection fields. Application to the gyrokinetic models in plasma physics, *J. Differential Equations* 249(2010) 1620-1663.
- [12] M. Bostan, Gyro-kinetic Vlasov equation in three dimensional setting. Second order approximation, *SIAM J. Multiscale Model. Simul.* 8(2010) 1923-1957.
- [13] M. Bostan, J.-A. Carrillo, Asymptotic fixed-speed reduced dynamics for kinetic equations in swarming, *Math. Models Methods Appl. Sci.* 23(2013) 2353-2393.

- [14] M. Bostan, J.-A. Carrillo, Reduced fluid models for self-propelled populations, interacting through alignment, *Math. Models Methods Appl. Sci.* 27(2017) 1255-1299.
- [15] M. Bostan, On the Boltzmann equation for charged particle beams under the effect of strong magnetic fields, *Discrete Contin. Dyn. Syst. Ser. B* 20(2015) 339-371.
- [16] M. Bostan, High magnetic field equilibria for the Fokker-Planck-Landau equation, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 33(2016) 899-931.
- [17] W. Braun, K. Hepp, The Vlasov dynamics and its fluctuations in the $1/N$ limit of interacting classical particles, *Comm. Math. Phys.* 56(1977) 101-113.
- [18] A. Bricard, J-B. Caussin, N. Desreumaux, O. Dauchot, D. Bartolo, Emergence of macroscopic directed motion in populations of motile colloids, *Nature* 503(2013) 95.
- [19] J. Buhl et al, From disorder to order in marching locusts, *Science* 312(2006) 1402-1406.
- [20] J.A. Cañizo, J.A. Carrillo, J. Rosado, A well-posedness theory in measures for some kinetic models of collective motion, *Math. Models Methods Appl. Sci.* 21(2011) 515-539.
- [21] J.A. Carrillo, M.R. D’Orsogna, V. Panferov, Double milling in a self-propelled swarms from kinetic theory, *Kinet. Relat. Models* 2(2009) 363-378.
- [22] J.A. Carrillo, M. Fornasier, J. Rosado, G. Toscani, Asymptotic flocking dynamics for the kinetic Cucker-Smale model, *SIAM J. Math. Anal.* 42(2010) 218-236.
- [23] J. A. Carrillo, M. Fornasier, G. Toscani, F. Vecil, Particle, Kinetic, and Hydrodynamic Models of Swarming, *Mathematical Modeling of Collective Behavior in Socio-Economic and Life Sciences*, Series: Modelling and Simulation in Science and Technology, Birkhauser (2010), 297-336.
- [24] J. A. Carrillo, Y.-P. Choi, M. Hauray, The derivation of swarming models: mean-field limit and Wasserstein distances, in *Collective dynamics from bacteria to crowds*, volume 553 *CISM Courses and Lect.*, pages 1–46. Springer, Vienna, 2014.
- [25] J. A. Carrillo, Y.-P. Choi, M. Hauray, S. Salem, Mean-field limit for collective behavior models with sharp sensitivity regions, *J. Eur. Math. Soc.* 21(2019) 121-161.
- [26] J.A. Carrillo, A. Klar, S. Martin, S. Tiwari, Self-propelled interacting particle systems with roosting force, *Math. Models Methods Appl. Sci* 20 (2010) 1533-1552.
- [27] H. Chaté, F. Ginelli, G. Grégoire, F. Raynaud, Collective motion of self-propelled particles interacting without cohesion, *Phys. Rev. E* 77(2008) 046113.
- [28] A. Creppy et al, Symmetry-breaking phase transitions in highly concentrated semen, *J. R. Soc. Interface* 13(2016) 20160575.
- [29] Y.-L. Chuang, M.R. D’Orsogna, D. Marthaler, A.L. Bertozzi, L. Chayes, State transitions and the continuum limit for a 2D interacting, self-propelled particle system, *Phys. D* 232(2007) 33-47.

- [30] Y.-L. Chuang, Y.R. Huang, M.R. D’Orsogna, A.L. Bertozzi, Multi-vehicle flocking: scalability of cooperative control algorithms using pairwise potentials, *IEEE International Conference on Robotics and Automation* (2007) 2292-2299.
- [31] I.D. Couzin, J. Krause, N.R. Franks, S.A. Levin, Effective leadership and decision making in animal groups on the move, *Nature* 433(2005) 513-516.
- [32] F. Cucker and S. Smale. Emergent behavior in flocks, *IEEE Trans. Automat. Control* 52(2007) 852-862.
- [33] P. Degond, G Dimarco, T. B. N. Mac, N. Wang, [Macroscopic models of collective motion with repulsion, *Comm. Math. Sci.* 13\(2015\)pp. 1615 – 1638.](#)
- [34] P. Degond, A. Frouvelle, J.-G. Liu, Macroscopic limits and phase transition in a system of self-propelled particles, *J. Nonlinear Sci.* 23 (2013) 427-456.
- [35] P. Degond, A. Frouvelle, J.-G. Liu, Phase transitions, hysteresis, and hyperbolicity for self-organized alignment dynamics, *Arch. Ration. Mech. Anal.* 216 (2015) 63–115.
- [36] P. Degond, A. Frouvelle, S. Merino-Aceituno, A. Trescases, [Quaternions in collective dynamics. *Multiscale Model. Simul.* 16 \(2018\) 28-77.](#)
- [37] P. Degond, A. Manhart, H. Yu, [A continuum model for nematic alignment of self-propelled particles, *Discrete Contin. Dyn. Syst. Ser. B* 22\(2017\) 1295–1327.](#)
- [38] P. Degond, S. Merino-Aceituno, F. Vergnet, H. Yu, [Coupled Self-Organized Hydrodynamics and Stokes models for suspensions of active particles. *J. Math. Fluid Mech.* 21\(2019\) 6.](#)
- [39] P. Degond, S. Motsch, Continuum limit of self-driven particles with orientation interaction, *Math. Models Methods Appl. Sci.* 18(2008) 1193-1215.
- [40] R. Dobrushin, Vlasov equations, *Funct. Anal. Appl.* 13(1979) 115-123.
- [41] M.R. D’Orsogna, Y.-L. Chuang, A.L. Bertozzi, L. Chayes, Self-propelled particles with soft-core interactions : Patterns, stability and collapse, *Phys. Rev. Lett.* 96(2006) 104302-1/4.
- [42] A. Frouvelle, A continuum model for alignment of self-propelled particles with anisotropy and density-dependent parameters, *Math. Models Methods Appl. Sci.* 22 (2012) 1250011.
- [43] A. Frouvelle, J.-G. Liu, Dynamics in a kinetic model of oriented particles with phase transition, *SIAM J. Math. Anal.* 44 (2012) 791-826.
- [44] S.-Y. Ha, J.-G. Liu, A simple proof of the Cucker-Smale flocking dynamics and mean-field limit, *Commun. Math. Sci.* 7(2009) 297-325.
- [45] S.-Y. Ha, K. Lee, D. Levy, Emergence of Time-Asymptotic Flocking in a Stochastic Cucker-Smale System, *Commun. Math. Sci.* 7(2009) 453-469.

- [46] S.-Y. Ha, E. Tadmor, From particle to kinetic and hydrodynamic descriptions of flocking, *Kinet. Relat. Models* 1 (2008) 415-435.
- [47] D. L. Koch, G. Subramanian, Collective hydrodynamics of swimming microorganisms: living fluids, *Annu. Rev. Fluid Mech.* 43 (2011) 637–659.
- [48] T. Kolokolnikov, J. A. Carrillo, A. Bertozzi, R. Fetecau, M. Lewis, Emergent behaviour in multi-particle systems with non-local interactions, *Phys. D* 260 (2013), 1–4.
- [49] D. Levermore, Entropic convergence and the linearized limit for the Boltzmann equation, *Comm. Partial Differential Equations* 18(1993) 1231-1248.
- [50] D. Levermore, Moment closure hierarchies for kinetic theories, *J. Stat. Phys.* 83 (1996) 1021-1065.
- [51] U. Lopez, J. Gautrais, I. D. Couzin, G. Theraulaz, From behavioural analyses to models of collective motion in fish schools, *Interface focus* 2(2012) 693-707.
- [52] M-C. Marchetti et al, Hydrodynamics of soft active matter, *Rev. Modern Phys.* 85(2013) 1143.
- [53] P. D. Miller, *Applied asymptotic analysis*, 75, American Mathematical Soc. 2006.
- [54] S. Motsch, E. Tadmor, A new model for self-organized dynamics and its flocking behavior, *J. Stat. Phys.* 144(2011) 923-947.
- [55] S. Motsch, E. Tadmor, Heterophilious dynamics enhances consensus, *SIAM Rev.* 56 (2014) 577–621.
- [56] H. Neunzert, The Vlasov equation as a limit of Hamiltonian classical mechanical systems of interacting particles, *Trans. Fluid Dynamics*, 18(1977) 663-678.
- [57] M. Reed, B. Simon, *Methods of modern mathematical physics, Vol. IV Analysis of Operators*, Academic Press, 1980.
- [58] D. Saintillan, M. J. Shelley, Instabilities, pattern formation, and mixing in active suspensions, *Phys. Fluids* 20(2008), 123304.
- [59] J. Toner, Y. Tu, Long-range order in a two-dimensional dynamical XY model: How birds fly together, *Phys. Rev. Lett.* 75(1995) 4326–4329.
- [60] T. Vicsek, A. Czirok, E. Ben-Jacob, I. Cohen, O. Shochet, Novel type of phase transition in a system of self-driven particles, *Phys. Rev. Lett.* 75 (1995) 1226-1229.
- [61] T. Vicsek, A. Zafeiris, Collective motion, *Phys. Rep.* 517(2012) 71–140.