

# Stationary compressible Stokes equations

T. Gallouët

joint works with

R. Eymard, A. Fettah, R. Herbin, J.-C. Latché, H. Lakehal

Bruxelles, mai 2013

# Stationary Compressible Stokes Equations

$\Omega$  is a connected bounded open set of  $\mathbb{R}^N$ ,  $N = 2$  or  $3$ , with a Lipschitz continuous boundary,

$$\operatorname{div}(\varphi(\rho)u) = 0 \text{ in } \Omega, \quad \rho \geq 0 \text{ in } \Omega, \quad \int_{\Omega} \rho(x) dx = M,$$

$$-\Delta u + \nabla p = f(\cdot, \rho) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

$$p = \eta(\rho) \text{ in } \Omega$$

$M > 0$ ,

$f$  is at most linear,  $|f(x, \rho)| \leq B(h(x) + |\rho|)$ ,  $h \in L^2(\Omega)$

$\varphi$  is Lipschitz continuous, increasing,  $\varphi(0) = 0$

$\eta$  is continuous superlinear,  $\liminf_{s \rightarrow +\infty} \eta(s)/s = +\infty$ , increasing  
( $\eta(0) = 0$ )

Functional spaces :  $u \in H_0^1(\Omega)^N$ ,  $p, \rho \in L^2(\Omega)$

## Main example

$$\varphi(\rho) = \rho$$

$$f(x, \rho) = \bar{f}(x) + g(x)\rho, \quad \bar{f} \in L^2(\Omega), \quad g \in L^\infty(\Omega)$$

$$\eta(\rho) = \rho^\gamma, \quad \gamma > 1.$$

$$\operatorname{div}(\rho u) = 0 \quad \text{in } \Omega, \quad \rho \geq 0 \quad \text{in } \Omega, \quad \int_{\Omega} \rho(x) dx = M,$$

$$-\Delta u + \nabla p = \bar{f} + g\rho \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

$$p = \rho^\gamma \quad \text{in } \Omega \quad (\text{Equation Of State})$$

$$M > 0$$

The main difficulty is the nonlinearity of the EOS (for passing to the limit with weak convergences)

# Existence of a solution

Existence of a (weak) solution can be proved passing to the limit on two different approximate solution

1. Approximate solution obtained using a convenient numerical scheme
2. Approximate solution obtained using a viscous regularization of the mass equation

The second way is essentially in previous works of P. L. Lions, E. Feireisl, A. Novotny...

No uniqueness result.

# Existence of a solution

Existence of a (weak) solution can be proved passing to the limit on two different approximate solution

1. Approximate solution obtained using a convenient numerical scheme
2. Approximate solution obtained using a viscous regularization of the mass equation. Convergence of the approximate solution proven with a simple proof which can be adapted in order to do the first method (in particular with schemes used in an industrial framework)

# Steps of the proof

1. Definition of the approximate problem and existence of an approximate solution
2. Estimates on the approximate solution
3. passing to the limit

## The approximate problem

$$T_n(s) = \min\{\max\{s, -n\}, n\}$$

Then, the regularized problem reads, for  $n, l, m \in \mathbb{N}^*$ ,

$$u \in H_0^1(\Omega)^N, \rho \in H^1(\Omega), p \in L^2(\Omega),$$

$$\int_{\Omega} \nabla u : \nabla v \, dx - \int_{\Omega} p \operatorname{div}(v) \, dx = \int_{\Omega} f_l(x, \rho) \cdot v \, dx, \quad \forall v \in H_0^1(\Omega)^N$$

$$\int_{\Omega} \varphi(\rho) u \cdot \nabla \psi \, dx - \frac{1}{n} \int_{\Omega} \nabla \rho(x) \cdot \nabla \psi(x) \, dx = 0, \quad \forall \psi \in H^1(\Omega)$$

$$\rho > 0 \text{ a.e. in } \Omega, \quad \int_{\Omega} \rho \, dx = M, \quad p = \eta_m(\rho) \text{ a.e. in } \Omega$$

where  $f_l(x, s) = T_l(f(x, s))$  and  $\eta_m(s) = T_m(\eta(s))$

# Existence of an approximate solution

Schauder fixed point for the application  $T$  from  $L^2(\Omega)$  to  $L^2(\Omega)$ ,  
 $T(\rho) = \rho$ .

$u \in H_0^1(\Omega)^N$ ,  $\rho \in H^1(\Omega)$ ,  $p \in L^2(\Omega)$ ,

$$\int_{\Omega} \nabla u : \nabla v \, dx - \int_{\Omega} p \operatorname{div}(v) \, dx = \int_{\Omega} f_I(x, \rho) \cdot v \, dx, \quad \forall v \in H_0^1(\Omega)^N$$

$$\int_{\Omega} \varphi(\rho) u \cdot \nabla \psi \, dx - \frac{1}{n} \int_{\Omega} \nabla \rho \cdot \nabla \psi \, dx = 0, \quad \forall \psi \in H^1(\Omega)$$

$\rho > 0$  a.e. in  $\Omega$ ,  $\int_{\Omega} \rho \, dx = M$ ,  $p = \eta_m(\rho^+)$  a.e. in  $\Omega$

where  $f_I(x, s) = T_I(f(x, s))$  and  $\eta_m(s) = T_m(\eta(s))$



## Intermediate problem

$u \in L^p(\Omega)^N$ ,  $p > N$  (true if  $u \in H_0^1(\Omega)^N$ ).  $M > 0$ .  $\varphi$  Lipschitz continuous and  $\varphi(0) = 0$ .

There exist a unique  $\rho$  solution of

$$\rho \in H^1(\Omega),$$
$$\int_{\Omega} \nabla \rho \cdot \nabla v \, dx - \int_{\Omega} \varphi(\rho) u \cdot \nabla v \, dx = 0, \quad \forall v \in H^1(\Omega)$$

with  $\int_{\Omega} \rho(x) \, dx = M$

Furthermore:

1.  $\rho > 0$  a.e. on  $\Omega$
2. For any  $A > 0$ , there exists  $C$  such that

$$\| |u| \|_{L^p(\Omega)} \leq A \Rightarrow \|\rho\|_{H^1(\Omega)} \leq C(A, p, M, \varphi, \Omega)$$

Proof using the Leray-Schauder topological degree

## Intermediate problem, main point

$u \in L^p(\Omega)^N$ ,  $p > N$ .  $M > 0$ .  $\varphi$  Lipschitz continuous and  $\varphi(0) = 0$ .

$$\rho \in H^1(\Omega),$$
$$\int_{\Omega} \nabla \rho \cdot \nabla v \, dx - \int_{\Omega} \varphi(\rho) u \cdot \nabla v \, dx = 0, \quad \forall v \in H^1(\Omega)$$

with  $\int_{\Omega} \rho(x) \, dx = M$

Proof of *a priori* positivity of  $\rho$  taking  $v = T_{\varepsilon}(\rho^+)$  and  $\varepsilon \rightarrow 0$

and uniqueness taking  $v = T_{\varepsilon}((\rho_1 - \rho_2)^+)$ , as in an old paper of Boccardo-G-Murat

# Intermediate problem, Leray-Schauder topological degree

$u \in L^p(\Omega)^N$ ,  $p > N$ .  $M > 0$ .  $\varphi$  Lipschitz continuous and  $\varphi(0) = 0$ .

$F$  from  $[0, 1] \times L^2(\Omega)$  to  $L^2(\Omega)$

$$F(t, \rho) = \rho$$

$$\rho \in H^1(\Omega),$$

$$\int_{\Omega} \nabla \rho \cdot \nabla v \, dx - \int_{\Omega} t \varphi(\rho) u \cdot \nabla v \, dx = 0, \quad \forall v \in H^1(\Omega)$$

with  $\int_{\Omega} \rho \, dx = tM$

$L^2(\Omega)$ -estimate on  $\rho$  if  $F(t, \rho) = \rho$

## Passing to the limit

$$n, l, m \in \mathbb{N}^*,$$

$$u \in H_0^1(\Omega)^N, \rho \in H^1(\Omega), p \in L^2(\Omega),$$

$$\int_{\Omega} \nabla u : \nabla v \, dx - \int_{\Omega} p \operatorname{div}(v) \, dx = \int_{\Omega} f_l(x, \rho) \cdot v \, dx, \forall v \in H_0^1(\Omega)^N$$

$$\int_{\Omega} \varphi(\rho) u \cdot \nabla \psi \, dx - \frac{1}{n} \int_{\Omega} \nabla \rho(x) \cdot \nabla \psi(x) \, dx = 0, \forall \psi \in H^1(\Omega)$$

$$\rho > 0 \text{ a.e. in } \Omega, \int_{\Omega} \rho \, dx = M, p = \eta_m(\rho) \text{ a.e. in } \Omega$$

$$m \rightarrow +\infty$$

$$l \rightarrow +\infty$$

$$n \rightarrow +\infty$$

$m \rightarrow +\infty$

$n, l$  are fixed

$u \in H_0^1(\Omega)^N$ ,  $\rho \in H^1(\Omega)$ ,  $p \in L^2(\Omega)$ ,

$$\int_{\Omega} \nabla u : \nabla v \, dx - \int_{\Omega} p \operatorname{div}(v) \, dx = \int_{\Omega} f_l(x, \rho) \cdot v \, dx, \quad \forall v \in H_0^1(\Omega)^N$$

$$\int_{\Omega} \varphi(\rho) u \cdot \nabla \psi \, dx - \frac{1}{n} \int_{\Omega} \nabla \rho(x) \cdot \nabla \psi(x) \, dx = 0, \quad \forall \psi \in H^1(\Omega)$$

$\rho > 0$  a.e. in  $\Omega$ ,  $\int_{\Omega} \rho \, dx = M$ ,  $p = \eta_m(\rho)$  a.e. in  $\Omega$

$H_0^1(\Omega)$ -estimate on  $u$  thanks to  $\int_{\Omega} \eta_m(\rho) \operatorname{div}(u) \, dx \leq 0$

$H^1(\Omega)$ -estimate on  $\rho$

$L^2(\Omega)$ -estimate on  $p$  taking  $\operatorname{div}(v) = p - m$  with  $v \in H_0^1(\Omega)^N$

and using  $\int_{\Omega} \rho \, dx = M$ .

$l \rightarrow +\infty$

$n$  is fixed

$u \in H_0^1(\Omega)^N$ ,  $\rho \in H^1(\Omega)$ ,  $p \in L^2(\Omega)$ ,

$$\int_{\Omega} \nabla u : \nabla v \, dx - \int_{\Omega} p \operatorname{div}(v) \, dx = \int_{\Omega} f_l(x, \rho) \cdot v \, dx, \quad \forall v \in H_0^1(\Omega)^N$$

$$\int_{\Omega} \varphi(\rho) u \cdot \nabla \psi \, dx - \frac{1}{n} \int_{\Omega} \nabla \rho(x) \cdot \nabla \psi(x) \, dx = 0, \quad \forall \psi \in H^1(\Omega)$$

$\rho > 0$  a.e. in  $\Omega$ ,  $\int_{\Omega} \rho \, dx = M$ ,  $p = \eta(\rho)$  a.e. in  $\Omega$

$H_0^1(\Omega)$ -estimate on  $u$  and  $L^2(\Omega)$ -estimate on  $p$  thanks to

$\int_{\Omega} \eta(\rho) \operatorname{div}(u) \, dx \leq 0$  and taking  $\operatorname{div}(v) = p - m$  with  $v \in H_0^1(\Omega)^N$   
and using  $\int_{\Omega} \rho \, dx = M$  and the superlinearity of  $\eta$

$H^1(\Omega)$ -estimate on  $\rho$

$n \rightarrow +\infty$

$$u_n \in H_0^1(\Omega)^N, \rho_n \in H^1(\Omega), p_n \in L^2(\Omega),$$

$$\int_{\Omega} \nabla u_n : \nabla v \, dx - \int_{\Omega} p_n \operatorname{div}(v) \, dx = \int_{\Omega} f(x, \rho_n) \cdot v \, dx, \forall v \in H_0^1(\Omega)^N$$

$$\int_{\Omega} \varphi(\rho_n) u_n \cdot \nabla \psi \, dx - \frac{1}{n} \int_{\Omega} \nabla \rho_n(x) \cdot \nabla \psi(x) \, dx = 0, \forall \psi \in H^1(\Omega)$$

$$\rho_n > 0 \text{ a.e. in } \Omega, \int_{\Omega} \rho_n \, dx = M, p_n = \eta(\rho_n) \text{ a.e. in } \Omega$$

$H_0^1(\Omega)$ -estimate on  $u_n$  and  $L^2(\Omega)$ -estimate on  $p_n$  thanks to

$\int_{\Omega} \eta(\rho_n) \operatorname{div}(u_n) \, dx \leq 0$  and taking  $\operatorname{div}(v_n) = p_n - m_n$  with  $v_n \in H_0^1(\Omega)^N$  and using  $\int_{\Omega} \rho_n \, dx = M$  and the superlinearity of  $\eta$

$L^2(\Omega)$ -estimate on  $\rho$

$$n \rightarrow +\infty$$

$$u_n \rightarrow u \text{ in } L^2(\Omega)^N \text{ and weakly in } H_0^1(\Omega)^N$$

$$\rho_n \rightarrow \rho \text{ weakly in } L^2(\Omega)$$

$$p_n \rightarrow p \text{ weakly in } L^2(\Omega)$$

But, we do not have an  $H^1(\Omega)$  estimate on  $\rho_n$

$$h_n = f(\cdot, \rho_n) \rightarrow h \text{ weakly in } L^2(\Omega)^N$$

$$q_n = \varphi(\rho_n) \rightarrow q \text{ weakly in } L^2(\Omega)$$

We need some additional tricks to prove  $h = f(\cdot, \rho)$ ,  $q = \varphi(\rho)$ ,  
 $p = \eta(\rho)$



## Passage to the limit in the momentum equation

$$v \in C_c^\infty(\Omega)^N,$$

$$\int_{\Omega} \nabla u_n : \nabla v \, dx - \int_{\Omega} p_n \operatorname{div}(v) \, dx = \int_{\Omega} h_n \cdot v \, dx.$$

$$n \rightarrow \infty$$

$$\int_{\Omega} \nabla u : \nabla v \, dx - \int_{\Omega} p \operatorname{div}(v) \, dx = \int_{\Omega} h \cdot v \, dx.$$

$$h = f(\cdot, \rho)?$$

## Passage to the limit in the mass equation

$$v \in C_c^\infty(\mathbb{R}^N)$$

$$\int_{\Omega} q_n u_n \cdot \nabla v \, dx - \frac{1}{n} \int_{\Omega} \nabla \rho_n \cdot \nabla v \, dx = 0,$$

$$n \rightarrow \infty$$

$$\int_{\Omega} q u \cdot \nabla v \, dx = 0$$

$$q \geq 0 \text{ a.e.}$$

$$q = \varphi(\rho)?$$

# Passage to the limit in the nonlinear functions of $\rho$

$h = f(\cdot, \rho)$ ? (easy if  $f(x, \rho) = \bar{f}(x) + g(x)\rho$ )

$q = \varphi(\rho)$ ? (easy if  $\varphi(\rho) = \rho$ )

$p = \eta(\rho)$

Idea: prove  $\int_{\Omega} p_n q_n \rightarrow \int_{\Omega} p q$  and deduce a.e. convergence (of  $p_n$  and  $\rho_n$ )

$\nabla : \nabla = \operatorname{div} \operatorname{div} + \operatorname{curl} \cdot \operatorname{curl}$

For all  $\bar{u}, \bar{v}$  in  $H_0^1(\Omega)^N$ ,

$$\int_{\Omega} \nabla \bar{u} : \nabla \bar{v} = \int_{\Omega} \operatorname{div}(\bar{u}) \operatorname{div}(\bar{v}) + \int_{\Omega} \operatorname{curl}(\bar{u}) \cdot \operatorname{curl}(\bar{v})$$

Then, for all  $\bar{v}$  in  $H_0^1(\Omega)^N$ , the momentum equation is

$$\begin{aligned} \int_{\Omega} \operatorname{div}(u_n) \operatorname{div}(\bar{v}) + \int_{\Omega} \operatorname{curl}(u_n) \cdot \operatorname{curl}(\bar{v}) \\ - \int_{\Omega} p_n \operatorname{div}(\bar{v}) = \int_{\Omega} h_n \cdot \bar{v} \end{aligned}$$

**Choice of  $\bar{v}$  ?**  $\bar{v} = \bar{v}_n$  with  $\operatorname{curl}(\bar{v}_n) = 0$ ,  $\operatorname{div}(\bar{v}_n) = q_n$  and  $\bar{v}_n$  bounded in  $H_0^1(\Omega)^N$  (unfortunately,  $0$  is impossible).

Then, up to a subsequence,

$\bar{v}_n \rightarrow v$  in  $L^2(\Omega)^N$  and weakly in  $H_0^1(\Omega)^N$ ,

$\operatorname{curl}(v) = 0$ ,  $\operatorname{div}(v) = q$ .

## Proof using $\bar{v}_n$ (1)

$$\begin{aligned} \int_{\Omega} \operatorname{div}(u_n) \operatorname{div}(\bar{v}_n) + \int_{\Omega} \operatorname{curl}(u_n) \cdot \operatorname{curl}(\bar{v}_n) - \int_{\Omega} p_n \operatorname{div}(\bar{v}_n) \\ = \int_{\Omega} h_n \cdot \bar{v}_n \end{aligned}$$

But,  $\operatorname{div}(\bar{v}_n) = q_n$  and  $\operatorname{curl}(\bar{v}_n) = 0$ . Then:

$$\int_{\Omega} (\operatorname{div}(u_n) - p_n) q_n = \int_{\Omega} h_n \cdot \bar{v}_n$$

$n \rightarrow \infty$

$$\lim_{n \rightarrow \infty} \int_{\Omega} (\operatorname{div}(u_n) - p_n) q_n = \int_{\Omega} h \cdot v$$

## Proof using $\bar{v}_n$ (2)

But, since  $-\Delta u + \nabla p = h$

$$\begin{aligned} \int_{\Omega} \operatorname{div}(u) \operatorname{div}(v) + \int_{\Omega} \operatorname{curl}(u) \cdot \operatorname{curl}(v) - \int_{\Omega} p \operatorname{div}(v) \\ = \int_{\Omega} h \cdot v \end{aligned}$$

which gives (using  $\operatorname{div}(v) = \rho$  and  $\operatorname{curl}(v) = 0$ )

$$\int_{\Omega} (\operatorname{div}(u) - \rho) q = \int_{\Omega} h \cdot v \quad \text{Then}$$

$$\lim_{n \rightarrow \infty} \int_{\Omega} (\rho_n - \operatorname{div}(u_n)) q_n = \int_{\Omega} (\rho - \operatorname{div}(u)) q$$

thanks to the mass equations,  $\int_{\Omega} q_n \operatorname{div}(u_n) \leq 0$  and  $\int_{\Omega} q \operatorname{div}(u) = 0$ . Then,

$$\limsup_{n \rightarrow \infty} \int_{\Omega} \rho_n q_n \leq \int_{\Omega} \rho q$$

## Error in the preceding proof

In the preceding proof, we used  $\bar{v}_n$  such that  $\operatorname{curl}(\bar{v}_n) = 0$ ,  $\operatorname{div}(\bar{v}_n) = \rho_n$  and  $\bar{v}_n$  bounded in  $H_0^1(\Omega)^N$ .

Unfortunately, it is impossible to have  $\bar{v}_n \in H_0^1(\Omega)^d$  but only  $\bar{v}_n \in H^1(\Omega)^N$ .

## Curl-free test function

Let  $w_n \in H_0^1(\Omega)$ ,  $-\Delta w_n = q_n$ ,

One has  $w_n \in H_{loc}^2(\Omega)$  since, for  $\psi \in C_c^\infty(\Omega)$ , one has  $\Delta(w_n\psi) \in L^2(\Omega)$  and

$$\begin{aligned} \sum_{i,j=1}^d \int_{\Omega} \partial_i \partial_j (w_n \psi) \partial_i \partial_j (w_n \psi) &= \sum_{i,j=1}^d \int_{\Omega} \partial_i \partial_i (w_n \psi) \partial_j \partial_j (w_n \psi) \\ &= \int_{\Omega} (\Delta(w_n \psi))^2 = C_\psi < \infty \end{aligned}$$

Then, taking  $v_n = \nabla w_n$

- ▶  $v_n \in (H_{loc}^1(\Omega))^N$ ,
- ▶  $\operatorname{div}(v_n) = q_n$  a.e. in  $\Omega$ ,
- ▶  $\operatorname{curl}(v_n) = 0$  a.e. in  $\Omega$ ,
- ▶  $H_{loc}^1(\Omega)$ -estimate on  $v_n$  with respect to  $\|q_n\|_{L^2(\Omega)}$ .

Then, up to a subsequence, as  $n \rightarrow \infty$ ,  $v_n \rightarrow v$  in  $L_{loc}^2(\Omega)^N$  and weakly in  $H_{loc}^1(\Omega)^N$ ,  $\operatorname{curl}(v) = 0$ ,  $\operatorname{div}(v) = q$ .



Proof of  $\int_{\Omega} (\rho_n - \operatorname{div}(u_n)) q_n \psi \rightarrow \int_{\Omega} (\rho - \operatorname{div}(u)) q \psi$

Let  $\psi \in C_c^\infty(\Omega)$  (so that  $v_n \psi \in H_0^1(\Omega)^N$ ). Taking  $\bar{v} = v_n \psi$ :

$$\begin{aligned} \int_{\Omega} \operatorname{div}(u_n) \operatorname{div}(v_n \psi) + \int_{\Omega} \operatorname{curl}(u_n) \cdot \operatorname{curl}(v_n \psi) - \int_{\Omega} \rho_n \operatorname{div}(v_n \psi) \\ = \int_{\Omega} h_n \cdot (v_n \psi). \end{aligned}$$

Using a proof similar to that given if  $\psi = 1$  (with additional terms involving  $\psi$ ), we obtain :

$$\lim_{n \rightarrow \infty} \int_{\Omega} (\rho_n - \operatorname{div}(u_n)) q_n \psi = \int_{\Omega} (\rho - \operatorname{div}(u)) q \psi$$

Proof of  $\int_{\Omega} (\rho_n - \operatorname{div}(u_n)) q_n \rightarrow \int_{\Omega} (\rho - \operatorname{div}(u)) q$

$$F_n = (\rho_n - \operatorname{div}(u_n)) q_n, \quad F = (\rho - \operatorname{div}(u)) q$$

$$F_n \rightarrow F \text{ in } D'(\Omega)$$

The sequence  $F_n$  is equiintegrable (since  $\rho_n - \operatorname{div}(u_n)$  is bounded in  $L^2$  and  $q_n^2$  is equiintegrable thanks to  $\rho_n$  bounded in  $L^2$  and  $\eta$  superlinear)

Then  $F_n \rightarrow F$  weakly in  $L^1(\Omega)$

Proving  $\int_{\Omega} p_n q_n \rightarrow \int_{\Omega} p q$

$$\int_{\Omega} (p_n - \operatorname{div}(u_n)) q_n \rightarrow \int_{\Omega} (p - \operatorname{div}(u)) q$$

But thanks to the mass equations:

$$\int_{\Omega} \operatorname{div}(u_n) q_n \leq 0, \quad \int_{\Omega} \operatorname{div}(u) q = 0;$$

Then:

$$\limsup_{n \rightarrow \infty} \int_{\Omega} p_n q_n \leq \int_{\Omega} p q$$

## a.e. convergence of $\rho_n$ and $p_n$ . Leray-Lions trick

Simple case:  $\varphi(\rho) = \rho$  and assuming  $\eta(\rho) \in L^2(\Omega)$ .

Let  $G_n = (\eta(\rho_n) - \eta(\rho))(\rho_n - \rho) \in L^1(\Omega)$  and  $G_n \geq 0$  a.e. in  $\Omega$ .

Futhermore

$G_n = (p_n - \eta(\rho))(\rho_n - \rho) = p_n\rho_n - p_n\rho - \eta(\rho)\rho_n + \eta(\rho)\rho$  and

$$\int_{\Omega} G_n = \int_{\Omega} p_n\rho_n - \int_{\Omega} p_n\rho - \int_{\Omega} \eta(\rho)\rho_n + \int_{\Omega} \eta(\rho)\rho$$

Using the weak convergence in  $L^2(\Omega)$  of  $p_n$  and  $\rho_n$  and

$$\lim_{n \rightarrow \infty} \int_{\Omega} p_n\rho_n \leq \int_{\Omega} p\rho$$

$$\lim_{n \rightarrow \infty} \int_{\Omega} G_n = 0,$$

Then (up to a subsequence),  $G_n \rightarrow 0$  a.e. and then  $\rho_n \rightarrow \rho$  a.e. (since  $\eta$  is increasing function). Finally

$p_n = \eta(\rho_n) \rightarrow \eta(\rho)$  in  $L^q(\Omega)$  for all  $1 \leq q < 2$ ,

$p_n \rightarrow p$  weakly in  $L^2(\Omega)$

then  $p = \eta(\rho)$ , similarly  $h = f(\cdot, \rho)$

## a.e. convergence of $\rho_n$ and $p_n$ , general case

The function  $\eta$  is a one-to-one function from  $\mathbb{R}_+$  onto  $\mathbb{R}_+$ . We denote by  $\bar{\eta}$  the reciprocal function of  $\eta$ . ( $\bar{\eta}$  sublinear)

Since  $p \in L^2(\Omega)$ , one has  $\bar{\eta}(p) \in L^2(\Omega)$  and we set  $\bar{\rho} = \bar{\eta}(p)$

$$G_n = (\varphi(\rho_n) - \varphi(\bar{\rho}))(\eta(\rho_n) - \eta(\bar{\rho}))$$

so that  $G_n \in L^1(\Omega)$ ,  $G_n \geq 0$  a.e.

$$0 \leq \int_{\Omega} G_n = \int_{\Omega} (q_n - \varphi(\bar{\rho}))(p_n - p)$$

Then

$$\lim_{n \rightarrow \infty} \int_{\Omega} G_n dx \leq \int_{\Omega} (q - \varphi(\bar{\rho}))(p - p) dx = 0.$$

This gives  $G_n \rightarrow 0$  in  $L^1(\Omega)$  and then, up to a subsequence,

$$G_n = (\varphi(\rho_n) - \varphi(\bar{\rho}))(\eta(\rho_n) - \eta(\bar{\rho})) \rightarrow 0 \text{ a.e. in } \Omega$$

We now use the fact that  $\varphi$  and  $\eta$  are increasing,  $\rho_n \rightarrow \bar{\rho}$  a.e. in  $\Omega$

Then  $\bar{\rho} = \rho$ ,  $q = \varphi(\rho)$ ,  $h = f(\cdot, \rho)$ ,  $p = \eta(\rho)$

## $\eta$ non decreasing instead of increasing

Simple case  $f(\cdot, \rho) = \bar{f} + g\rho$ ,  $\varphi(\rho) = \rho$

One proves  $\rho = \eta(\rho)$  with the Minty trick (but no a.e. convergence)

We set  $\eta(s) = 0$  for  $s < 0$  and  $\bar{\eta}$  is the reciprocal function of  $s \mapsto \eta(s) + s$  (which is a one-to-one function from  $\mathbb{R}$  onto  $\mathbb{R}$ )

Let  $\bar{\rho} \in L^2(\Omega)$  and  $\bar{\rho} = \bar{\eta}(\bar{\rho})$  so that  $\bar{\rho} \in L^2(\Omega)$

$$0 \leq \int_{\Omega} (\rho_n - \bar{\rho})(\eta(\rho_n) - \eta(\bar{\rho})) = \int_{\Omega} (\rho_n - \bar{\rho})(\rho_n - \bar{\rho} + \bar{\rho})$$

$$0 \leq \int_{\Omega} (\rho - \bar{\rho})(\rho - \bar{\rho} + \bar{\rho})$$

which gives also

$$0 \leq \int_{\Omega} (\rho - \bar{\rho})(\rho - \bar{\rho} + \rho) = \int_{\Omega} (\rho - \bar{\eta}(\bar{\rho}))(\rho - \bar{\rho} + \rho)$$

$\eta$  non decreasing instead of increasing

For all  $\bar{p} \in L^2(\Omega)$

$$0 \leq \int_{\Omega} (\rho - \bar{\eta}(\bar{p}))(\rho - \bar{p} + \rho) dx$$

Let  $\psi \in C_c^\infty(\Omega)$ ,  $\epsilon > 0$ . Taking  $\bar{p} = \rho + \rho + \epsilon\psi$ , letting  $\epsilon \rightarrow 0$  leads to, with the Dominated Convergence Theorem,

$$0 \leq - \int_{\Omega} (\rho - \bar{\eta}(\rho + \rho))\psi dx.$$

Since  $\psi$  is arbitrary in  $C_c^\infty(\Omega)$ , we then conclude that  $\rho = \bar{\eta}(\rho + \rho)$  which gives  $\eta(\rho) + \rho = \rho + \rho$  and then  $\rho = \eta(\rho)$ .

# Generalizations

- ▶ (Easy) Complete Diffusion term:  $-\mu\Delta u - \frac{\mu}{3}\nabla(\operatorname{div} u)$ , with  $\mu \in \mathbb{R}_+^*$  given, instead of  $-\Delta u$ .
- ▶ Stationary compressible Navier Stokes equation  $\eta(\rho) = \rho^\gamma$ ,  $\gamma > 3$  if  $N = 3$ .
- ▶ (Ongoing work) Navier-Stokes Equations with  $N = 3$  and  $\frac{3}{2} < \gamma \leq 3$ . (probably sharp result with respect to  $\gamma$  without changing the diffusion term or the EOS)
- ▶ (Ongoing work) Evolution equation (Stokes and Navier-Stokes)
- ▶ (Open question) Other boundary condition. Addition of an energy equation



# Stationary compressible Navier Stokes equations

$\Omega$  is a bounded open set of  $\mathbb{R}^N$ ,  $N = 2$  or  $3$ , with a Lipschitz continuous boundary,  $\gamma > 1$ ,  $f \in L^2(\Omega)^N$  and  $M > 0$

$$-\Delta u + (\rho u \cdot \nabla)u + \nabla p = f \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

$$\operatorname{div}(\rho u) = 0 \text{ in } \Omega, \quad \rho \geq 0 \text{ in } \Omega, \quad \int_{\Omega} \rho(x) dx = M,$$

$$p = \rho^\gamma \text{ in } \Omega$$

Functional spaces :  $u \in H_0^1(\Omega)^N$ ,  $p \in L^{\bar{q}}(\Omega)$ ,  $\rho \in L^{\gamma\bar{q}}(\Omega)$ .

If  $d = 2$  or if  $d = 3$  and  $\gamma \geq 3$  :  $\bar{q} = 2$ .

If  $d = 3$  and  $\frac{3}{2} < \gamma < 3$  :  $\bar{q} = \frac{3(\gamma-1)}{\gamma}$ .

$$\gamma = \frac{3}{2}, \quad \frac{3\gamma-1}{\gamma} = 1, \quad 3(\gamma-1) = \frac{3}{2}$$