

SURVIVAL OF A SINGLE MUTANT IN ONE DIMENSION

BY ENRIQUE ANDJEL[†], JUDITH MILLER^{*,§} AND ETIENNE PARDOUX^{†,‡}

Université de Provence[‡] and Georgetown University[§]

We study a one dimensional two-type contact process and give necessary and sufficient conditions on the initial configuration for both types to survive forever. These results are proved under the assumption that the rates of propagation (and death) of the two types are equal.

1. Introduction. The aim of this paper is to study the probability that a single mutant in an infinite population of residents will survive. We consider this problem in the framework of the one dimensional two-type contact process.

We will prove that if the mutant has no selective advantage nor disadvantage, compared with the individuals of the resident population, then, provided we are in the supercritical case (which means that a single individual's progeny may survive for ever), a single mutant with an empty half-line in front of him, and all sites behind him occupied by resident individuals, has a progeny which survives forever with positive probability, while any finite number of mutants, with infinitely many residents on both sides, have a progeny which goes extinct a. s. Note that we define the progeny at time t of a given ancestor at time 0 as the set of individuals alive at time t , who are the descendants of that ancestor at time 0.

We shall next discuss how far those results remain true or differ significantly, in case of a selective advantage or disadvantage.

Let us now explain what we mean by the contact process. Note that this process is often presented in the language of infection. We shall rather consider it here as a model of the spread of a population. Consider first the usual one-type contact process with birth parameter $\lambda > 0$. This process $\{\xi_t, t \geq 0\}$ is a $\{0, 1\}^{\mathbb{Z}}$ -valued Markov process, hence ξ_t is a random mapping which to each $x \in \mathbb{Z}$ associates $\xi_t(x) \in \{0, 1\}$. The statement $\xi_t(x) = 1$ means that the site x is occupied at time t , while $\xi_t(x) = 0$ means that site x is empty at time t . The process evolves as follows. Let x be such that $\xi_0(x) = 1$.

*Supported in part by National Science Foundation grants DMS-0201173 and DMS-0818727

[†]Supported in part by ANR MAEV under contract ANR-06-BLAN-0113

AMS 2000 subject classifications: Primary 60J25; Secondary 92D5

Keywords and phrases: two-type contact process, survival

We wait a random exponential time with parameter $1 + 2\lambda$. At that time, with probability $1/(1 + 2\lambda)$, the individual at site x dies; with probability $\lambda/(1 + 2\lambda)$, the individual, while continuing its own life at site x , gives birth to another individual; the newborn occupies site $x + 1$ if it is empty, and dies otherwise; and with probability $\lambda/(1 + 2\lambda)$, it gives birth to a newborn who occupies site $x - 1$ if this one is empty, and dies otherwise. Then the same operation repeats itself until site x becomes empty, independently of what happened so far. The same happens at any occupied site, and the exponential clocks at various sites are mutually independent. Note that we will use the same notation ξ_t to denote the random element of $\{0, 1\}^{\mathbb{Z}}$ defined above, and the random subset of \mathbb{Z} which contains all sites $x \in \mathbb{Z}$ where $\xi_t(x) = 1$.

The two-type contact process $\{\eta_t, t \geq 0\}$ is a $\{0, 1, 2\}^{\mathbb{Z}}$ -valued Markov process which starts from an initial condition (A, B) , where A and B are two nonintersecting subsets of \mathbb{Z} , A denoting the set of sites which are occupied by type 1 individuals and B the set of sites which are occupied by type 2 individuals at time $t = 0$. In other words,

$$\eta_0(x) = \begin{cases} 0, & \text{if } x \notin A \cup B; \\ 1, & \text{if } x \in A; \\ 2, & \text{if } x \in B. \end{cases}$$

The two-type contact process with equal birth rates λ evolves exactly like the one-type process, with each individual possibly giving birth to individuals of the same type. We shall consider in section 4 the case where the birth rate of the mutants (i. e. type 2 individuals) differs from that of the residents (i. e. type 1 individuals).

The (one-type) contact process has been extensively studied and plays a central role in the theory of interacting particle systems (see [3], [4] and references therein) but there are very few papers on the two-type contact process (see [2] and [5]).

Let us now present a useful construction of the contact process, called the graphical representation, which is valid in both the one-type and the two-type cases (at least in the case of equal birth rates), the two-type case with different birth rates requiring an additional refinement, which we will explain in section 4 below. The important feature of this construction is that processes corresponding to different initial conditions are coupled through it. Indeed, $\{\xi_t, t \geq 0\}$ (resp. $\{\eta_t, t \geq 0\}$) is a fixed function of both the initial condition, and the set of Poisson point processes, which code all the randomness, which we now introduce.

Consider a collection $\{P_t^x, P_t^{x,+}, P_t^{x,-}, t \geq 0; x \in \mathbb{Z}\}$ of mutually independent Poisson point processes, such that the P^x 's have intensity 1 while

both the $P^{x,+}$'s and the $P^{x,-}$'s have intensity λ , all defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. At each time of the point process P^x , we place a δ on the line $\{y = x, t \geq 0\}$ in the set $\{(y, t); y \in \mathbb{Z}, t \geq 0\}$. At each time t of the point process $P^{x,+}$ we place an arrow from (x, t) to $(x+1, t)$. Finally, at each time t of the point process $P^{x,-}$ we place an arrow from (x, t) to $(x-1, t)$.

The process $\{\xi_t^A, : t \geq 0\}$ is defined as follows. An open path in $\mathbb{Z} \times [0, +\infty)$ is a connected oriented path which moves along the time lines in the increasing t direction without passing through a δ symbol, and along birth arrows, in the direction of the arrow. Now

$$\{y; \xi_t^A(y) = 1\} = \{y \in \mathbb{Z}; \exists x \in A \text{ with an open path from } (0, x) \text{ to } (t, y)\}.$$

In the case of the two-type contact process, we need to put a restriction on the definition of an open path. An open path follows only those arrows which reach a site where no individual is alive at that time. For A, B two disjoint subsets of \mathbb{Z} , we define $\{\eta_t^{A,B}, t \geq 0\}$ as the $\{0, 1, 2\}^{\mathbb{Z}}$ -valued process whose value at time t is given by

$$\begin{aligned} \{y; \eta_t^{A,B}(y) = 1\} &= \{y \in \mathbb{Z}; \exists x \in A \text{ with an open path from } (0, x) \text{ to } (t, y)\} \\ \{y; \eta_t^{A,B}(y) = 2\} &= \{y \in \mathbb{Z}; \exists x \in B \text{ with an open path from } (0, x) \text{ to } (t, y)\} \end{aligned}$$

Given a finite subset $B \subset \mathbb{Z}$, write $B^+ = \{x, x > y, \forall y \in B\}$ and $B^- = \{x, x < y, \forall y \in B\}$.

The aim of this paper is to prove

THEOREM 1.1. *Suppose that $0 < |B| < \infty$. Then*

$$\mathbb{P}(\{x, \eta_t^{A,B}(x) = 2\} \neq \emptyset, \forall t > 0) > 0$$

if and only if at least one of the two sets $A \cap B^+$ and $A \cap B^-$ is finite.

From the results needed to prove Theorem 1.1 we can also deduce:

THEOREM 1.2. *Suppose that $0 < |A| < \infty$ and $0 < |B| < \infty$. Then*

$$\mathbb{P}(\{x, \eta_t^{A,B}(x) = 1\} \neq \emptyset, \{x, \eta_t^{A,B}(x) = 2\} \neq \emptyset, \forall t > 0) > 0.$$

We conjecture that Theorem 1.2 holds for the two-type contact process on \mathbb{Z}^d for all $d \geq 1$. In [5] it is proved that for $d \leq 2$ and all initial configurations $\lim_{t \rightarrow \infty} \mathbb{P}(\eta_t(x) = 1, \eta_t(y) = 2) = 0$ for all x, y , while for $d \geq 3$ the process admits invariant measures μ such that for all $x \neq y$, $\mu(\{\eta : \eta(x) = 1, \eta(y) =$

2}) > 0. Although this last result may be seen as evidence favoring our conjecture (when $d \geq 3$) it does not imply it nor is it implied by it.

The paper is organized as follows. In section 2, we recall and prove several results on the one-type contact process which are needed in further sections. In section 3, we study the case of a single or a finite number of mutants confronted with an infinite number of residents, in the case of equal birth rates. Theorems 1.1 and 1.2 are proved in subsections 3.3 and 3.4 respectively. Finally, in section 4, we conclude with a discussion of the case of unequal birth rates (i. e. when one of the two species has a selective advantage). We formulate one result and two conjectures.

2. Some results on the one-type contact process. Let $\{\xi_t^A, t \geq 0\}$ denote the contact process starting from the configuration whose set of occupied sites is A . We will write ξ_t^x for $\xi_t^{\{x\}}$. We shall use the notation

$$(2.1) \quad \rho = \mathbb{P}(\xi_t^0 \neq \emptyset, \forall t > 0) = \lim_{s \rightarrow \infty} \mathbb{P}(\xi_s^0 \neq \emptyset).$$

It follows from well-known results on the contact process, see e. g. Liggett [3], that there exists $\lambda_c < \infty$ such that $\rho > 0$ whenever $\lambda > \lambda_c$, which we shall suppose from now on.

Let \mathbb{Z}^- be the set of integers smaller than or equal to 0 and let \mathbb{Z}^+ be the set of integers greater than or equal to 0.

Let $r_t = \sup \{x : \xi_t^{\mathbb{Z}^-}(x) = 1\}$ and let $\ell_t = \inf \{x : \xi_t^{\mathbb{Z}^+}(x) = 1\}$.

It is known that since $\lambda > \lambda_c$, there exists $v = v(\lambda) > 0$ such that

$$\lim_{t \rightarrow \infty} \frac{r_t}{t} = - \lim_{t \rightarrow \infty} \frac{\ell_t}{t} = v \text{ a.s. and in } L^1(\Omega, \mathcal{F}, \mathbb{P}).$$

For a proof of these results the reader is referred to Theorems VI.2.19 and VI.2.24 in [3].

Let $R_t = \sup_{s \leq t} r_s$.

LEMMA 2.1. $\mathbb{P}(r_t \geq a) \geq \frac{a}{2} \mathbb{P}(R_t \geq a), \forall t, a.$

PROOF: Let $\tau_a = \inf \{s : r_s \geq a\}$. Then

$$\mathbb{P}(r_t \geq a | R_t \geq a) = \mathbb{P}(r_t \geq a | \tau_a \leq t).$$

By the strong Markov property this is bounded below by

$$\inf_{s \geq 0} \mathbb{P}(\xi_s^0 \cap [0, \infty) \neq \emptyset),$$

which by symmetry is at least

$$\inf_{s \geq 0} \frac{1}{2} \mathbb{P}(\xi_s^0 \neq \emptyset) = \frac{\rho}{2}.$$

□

LEMMA 2.2. $\lim_{t \rightarrow \infty} \frac{R_t}{t} = v$ a.s. and in L^1 .

PROOF: The a. s. convergence follows from the a.s convergence of $\frac{r_t}{t}$ and the fact that $v > 0$.

For the L^1 convergence note first that since $\frac{R_t}{t} \geq \frac{r_t}{t}$ and $\frac{r_t}{t}$ converges to v in L^1 , it suffices to show that

$$\lim_{t \rightarrow \infty} \mathbb{E} \left[\frac{R_t}{t} - v \right]^+ = 0$$

To do so fix $\varepsilon > 0$ and let $c = \frac{2}{\rho}$. Then write

$$\begin{aligned} \lim_{t \rightarrow \infty} \sum_{n=1}^{\infty} \mathbb{P} \left(\frac{R_t}{t} - v \geq \varepsilon n \right) &\leq \lim_{t \rightarrow \infty} c \sum_{n=1}^{\infty} \mathbb{P} \left(\frac{r_t}{t} - v \geq \varepsilon n \right) \\ &\leq \lim_{t \rightarrow \infty} \frac{c}{\varepsilon} \mathbb{E} \left[\frac{r_t}{t} - v \right]^+ \\ &= 0, \end{aligned}$$

where for the first inequality we used Lemma 2.1. Hence

$$\limsup_{t \rightarrow \infty} \mathbb{E} \left[\frac{R_t}{t} - v \right]^+ \leq \limsup_{t \rightarrow \infty} \varepsilon \sum_{n=0}^{\infty} \mathbb{P} \left(\frac{R_t}{t} - v \geq \varepsilon n \right) \leq \varepsilon.$$

Since ε is arbitrary the lemma is proved. □

The following lemma is an easy consequence of the fact that $v(\lambda) > 0$ whenever λ is strictly bigger than the critical value of the contact process on \mathbb{Z} .

LEMMA 2.3. *The critical values of λ for the contact processes on \mathbb{N} and \mathbb{Z} are equal.*

Let μ^+ denote the upper invariant measure for the contact process on \mathbb{N} . This is defined as follows. Denote by $\{\chi_t, t \geq 0\}$ the one-type contact process on \mathbb{N} . This process takes its values in $\{0, 1\}^{\mathbb{N}}$. In accordance with

the above conventions, for $A \subset \mathbb{N}$, we write χ_t^A for the contact process on \mathbb{N} starting with the initial condition $\chi_0^A(x) = 1$ iff $x \in A$. Then μ^+ is the weak limit, as $t \rightarrow \infty$, of the law of $\chi_t^{\mathbb{N}}$.

Define moreover for $\eta \in \{0, 1\}^{\mathbb{N}}$, $Y(\eta) = \inf \{x > 0 : \eta(x) = 1\}$.

LEMMA 2.4. $\alpha := \mathbb{E}_{\mu^+}(Y) < \infty$.

For the proof of this result, we will need the following

LEMMA 2.5. Denoting again $r_t = \sup\{x, \xi_t^{\mathbb{Z}^-}(x) = 1\}$, we have

$$\mu^+(Y > n) = \mathbb{P}(\inf_{t>0} r_t \leq -n).$$

PROOF: We first exploit the well-known self-duality of the contact process. Since there is a one to one correspondance between the open paths from some $(y, 0)$, $y \in \mathbb{N}$, to some (x, t) , $x \in (0, n]$ and the open paths from some $(x, 0)$, $x \in (0, n]$ to some (y, t) , $y \in \mathbb{N}$ obtained by reversing the directions of the arrows,

$$\mathbb{P}(\exists x \in (0, n] : \chi_t^{\mathbb{N}}(x) = 1) = \mathbb{P}(\exists x \in (0, n] : \chi_t^x \neq \emptyset).$$

Letting $t \rightarrow \infty$ in the above identity yields

$$\mu^+(Y \leq n) = \mathbb{P}(\exists x \in (0, n], \chi_t^x \neq \emptyset, \forall t > 0).$$

The last right hand side is the probability that there is an infinite open path starting from some $(x, 0)$, $x \in (0, n]$, which visits only points located at the right of the vertical line $\{1\} \times \mathbb{R}_+$. This has the same probability as the event that there is in $(-n, \infty) \times \mathbb{R}_+$ an infinite open path starting in $(-n, 0] \times \{0\}$, i. e. it equals $\mathbb{P}(\inf_{t>0} r_t > -n)$. The result follows. \square

PROOF OF LEMMA 2.4 In view of Lemma 2.5, it suffices to show that for the contact process on \mathbb{Z} there exist constants $K, c > 0$ such that for

$$\mathbb{P}(\inf_{t>0} r_t \leq -n) \leq K e^{-cn} \quad \forall n \geq 1.$$

It follows from Corollary VI.3.22 in [3] that for some $K_1, c > 0$ we have:

$$(2.2) \quad \mathbb{P}\left(r_t \leq \frac{v}{2}t\right) \leq K_1 e^{-ct} \quad \forall t \geq 1.$$

We first deduce that (here and below $[t]$ stands for the integer part of t)

$$\begin{aligned} \mathbb{P}\left(\inf_{n \geq [t], n \in \mathbb{N}} r_n \leq \frac{v}{2}t\right) &\leq \mathbb{P}\left(\bigcup_{n \geq [t], n \in \mathbb{N}} \left\{r_n \leq \frac{v}{2}n\right\}\right) \\ &\leq \sum_{n \geq [t]} \mathbb{P}\left(r_n \leq \frac{v}{2}n\right) \\ &\leq K_2 e^{-ct}. \end{aligned}$$

Next note that defining $\tau = \inf\{n < s \leq n+1, r_s \leq \frac{v}{2}t\}$ (with the convention that $\tau = n+1$ on the set $\{\inf_{n < s \leq n+1} r_s > \frac{v}{2}t\}$), we have

$$\left\{\inf_{n < s \leq n+1} r_s \leq \frac{v}{2}t\right\} \cap \{r_{n+1} - r_\tau = 0\} \subset \{r_{n+1} \leq \frac{v}{2}t\}$$

where the two sets on the left are mutually independent, and

$$\mathbb{P}(r_{n+1} - r_\tau = 0) \geq \mathbb{P}(X = 0),$$

the law of X being Poisson(λ). In other words $\mathbb{P}(r_{n+1} - r_\tau = 0) \geq e^{-\lambda}$, and we have

$$\begin{aligned} \mathbb{P}\left(\inf_{s \geq t} r_s \leq \frac{v}{2}t\right) &\leq \sum_{n \geq [t]} \mathbb{P}\left(\inf_{n < s \leq n+1} r_s \leq \frac{v}{2}t\right) \\ &\leq e^\lambda \sum_{n \geq [t]} \mathbb{P}\left(r_{n+1} \leq \frac{v}{2}t\right) \\ &\leq K e^{-ct}. \end{aligned}$$

We have shown in particular that

$$(2.3) \quad \mathbb{P}(\inf_{s \geq t} r_s \leq 0) \leq K e^{-ct}.$$

Fix $\beta > 0$ such that $2\lambda\beta < 1 + v\beta$. Now, write

$$\mathbb{P}(\inf_{t \geq 0} r_t \leq -n) \leq \mathbb{P}(\inf_{0 \leq t \leq \beta n} r_t \leq -n) + \mathbb{P}(\inf_{t \geq \beta n} r_t \leq 0).$$

It follows from (2.3) that the second term of the right hand side decays exponentially in n . Hence, the lemma will be proved if we show that the first term also decays exponentially in n . To do so, let $\tau = \inf\{t : r_t \leq -n\}$ and let Y_n be a Poisson random variable of parameter $2\lambda\beta n$. It now follows from the Strong Markov property applied at the stopping time τ that:

$$\mathbb{P}(r_{2\beta n} \leq v\beta n) \geq \mathbb{P}(\tau \leq \beta n)\mathbb{P}(Y_n \leq (1 + v\beta)n).$$

Since $\lim_{n \rightarrow \infty} \mathbb{P}(Y_n \leq (1 + v\beta)n) = 1$ and from (2.2) $\mathbb{P}(r_{2\beta n} \leq v\beta n)$ decays exponentially in n , it follows that the same happens to

$$\mathbb{P}(\tau \leq \beta n) = \mathbb{P}\left(\inf_{0 \leq t \leq \beta n} r_t \leq -n\right).$$

□

Let T^{-1} be the operator on the set of probability measures on $\{0, 1\}^{\mathbb{N}}$ defined by

$$T^{-1}(\nu)(\eta(x_1) = \gamma_1, \dots, \eta(x_n) = \gamma_n) = \nu(\eta(x_1 + 1) = \gamma_1, \dots, \eta(x_n + 1) = \gamma_n),$$

for any $n \geq 1$, $\gamma_1, \dots, \gamma_n \in \{0, 1\}$.

The natural partial order on $\{0, 1\}^{\mathbb{N}}$ induces a partial order on the set of probability measures on $\{0, 1\}^{\mathbb{N}}$ which we denote by \leq .

We have

$$\text{LEMMA 2.6. } T^{-1}(\mu^+) \geq \mu^+.$$

PROOF: Consider the contact process $\{\chi_t, t \geq 0\}$ this time on $\mathbb{N} \cup \{0\}$, starting again from $\chi_0 \equiv 1$. Let now $\{\bar{\chi}_t, t \geq 0\}$ denote the same process, with the same initial condition and the same realization of the graphical representation, except that we delete all arrows between states 0 and 1. The restriction to \mathbb{N} of the asymptotic (as $t \rightarrow \infty$) law of $\bar{\chi}_t$ coincides with μ^+ , while the same law associated with χ_t coincides with $T^{-1}(\mu^+)$. The result follows from the fact that for all $t > 0$, $x \geq 1$, $\mathbb{P}(\chi_t(x) \geq \bar{\chi}_t(x)) = 1$. □

To prove our next lemma we will use Theorem 2 from [1]. This theorem is stated for the contact process in \mathbb{Z} but its proof is also valid for the contact process on \mathbb{N} . In this lemma the space $\{0, 1\}^{\mathbb{N}}$ is endowed with its natural partial order, and μ^+ denotes the upper invariant measure for the contact process on \mathbb{N} .

LEMMA 2.7. *Let f be a continuous increasing real valued function on $\{0, 1\}^{\mathbb{N}}$ which depends only upon coordinates which are greater than or equal to $x + 1$ (for some $x \in \mathbb{N}$). Then*

$$\int f d\mu^+(\cdot | \eta(1) = 0, \dots, \eta(x-1) = 0, \eta(x) = 1) \geq \int f d\mu^+.$$

PROOF: Consider the contact process $\{\chi_t, t \geq 0\}$ on \mathbb{N} , starting from $\chi_0 \equiv 1$. Applying the above mentioned result of [1] we obtain:

$$\mathbb{E}(f(\chi_t) | \chi_t(1) = 0, \dots, \chi_t(x-1) = 0, \chi_t(x) = 1) \geq \mathbb{E}(f(\chi_t) | \chi_t(x) = 1),$$

for any deterministic initial configuration η . It then follows from Lemma 2.8 below that

$$\mathbb{E}(f(\chi_t) \mid \chi_t(1) = 0, \dots, \chi_t(x-1) = 0, \chi_t(x) = 1) \geq \mathbb{E}(f(\chi_t)).$$

It remains to let $t \rightarrow \infty$. \square

LEMMA 2.8. *Let $\{\chi_t, t \geq 0\}$ denote the contact process on \mathbb{N} , starting from any deterministic initial condition. For any $t > 0$, the law of χ_t has positive correlations.*

PROOF: For the contact process on $[1, \dots, n]$, the result follows from Theorem 2.14 on page 80 of Liggett [3]. Our result then follows by letting $n \rightarrow \infty$. \square

Note that Lemma 2.8 applies as well to the contact process $\{\xi_t, t \geq 0\}$ on \mathbb{Z} .

Let $S^{x,y} = \{\xi_t^x \neq \emptyset, \forall t > 0; \xi_t^y \neq \emptyset, \forall t > 0\}$. Recall that both processes $\{\xi_t^x, t \geq 0\}$ and $\{\xi_t^y, t \geq 0\}$ are constructed with the same set of Poisson processes $\{P_t^x, P_t^{x,+}, P_t^{x,-}, x \in \mathbb{Z}\}$ as explained above. Note that on the event $S^{x,y}$ the process starting from $\{x, y\}$ survives but this does not mean that under that initial condition the progeny of (say) x lives forever. We now show that (recall the definition of ρ in (2.1))

LEMMA 2.9. *For all $x, y \in \mathbb{Z}$*

$$\mathbb{P}(S^{x,y}) \geq \rho^2.$$

PROOF: Denoting by μ the upper invariant measure of the contact process $\{\xi_t, t \geq 0\}$ on \mathbb{Z} , i. e. μ is the limit as $t \rightarrow \infty$ of the law of $\xi_t^{\mathbb{Z}}$, we have by the same duality argument already used in the proof of Lemma 2.5 the identities

$$\begin{aligned} \mathbb{P}(\xi_t^x \neq \emptyset, \forall t > 0) &= \mu(\eta(x) = 1), \\ \mathbb{P}(\xi_t^y \neq \emptyset, \forall t > 0) &= \mu(\eta(y) = 1), \\ \mathbb{P}(S^{x,y}) &= \mu(\eta(x) = 1, \eta(y) = 1). \end{aligned}$$

Letting $t \rightarrow \infty$ in the result of Lemma 2.8 applied to the contact process on \mathbb{Z} implies that μ has positive correlations, which implies that

$$\mu(\eta(x) = 1, \eta(y) = 1) \geq \mu(\eta(x) = 1) \times \mu(\eta(y) = 1).$$

The result follows from this inequality and the three above identities. \square

We now fix some $\lambda > \lambda_c$ and let $v = v(\lambda)$. We pick

$$0 < \varepsilon < \frac{v}{2} \wedge \frac{\rho^2}{4}.$$

From now on t_0 will be a large enough multiple of $\frac{2}{v}$ so that the following holds :

$$(2.4) \quad \mathbb{P}(B(t_0, \varepsilon)) \geq 1 - \varepsilon,$$

where

$$B(t_0, \varepsilon) = \left\{ v - \varepsilon \leq \frac{r_{t_0}}{t_0} \leq v + \varepsilon, v - \varepsilon \leq -\frac{\ell_{t_0}}{t_0} \leq v + \varepsilon \right\}.$$

Let us define new processes. For any $z \in \mathbb{Z}$, we write

$$\begin{aligned} r_t^z &= \sup \{x : \xi_t^z(x) = 1\} - z, \\ \ell_t^z &= \inf \{x : \xi_t^z(x) = 1\} - z, \end{aligned}$$

where as usual the sup (resp. the inf) over an empty set is $+\infty$ (resp. $-\infty$).

Now we define the event

$$\begin{aligned} C(v, t_0, \varepsilon) &= \left\{ v - \varepsilon \leq \frac{r_{t_0}^0}{t_0} \leq v + \varepsilon, v - \varepsilon \leq -\frac{\ell_{t_0}^0}{t_0} \leq v + \varepsilon \right\} \\ &\cap \left\{ v - \varepsilon \leq \frac{r_{t_0}^{vt_0}}{t_0} \leq v + \varepsilon, v - \varepsilon \leq -\frac{\ell_{t_0}^{vt_0}}{t_0} \leq v + \varepsilon \right\}, \end{aligned}$$

and prove:

LEMMA 2.10. *Let ε be as above. Then, for any large enough t_0 , we have:*

$$\mathbb{P}(C(v, t_0, \varepsilon)) \geq \rho^2 - 2\varepsilon.$$

PROOF: First note that on the event $\{\xi_t^0 \neq \emptyset, \forall t > 0\}$ we have: $r_t^0 = r_t$ and $\ell_t^0 = \ell_t$ and a similar result holds for $r_t^{vt_0}$ and $\ell_t^{vt_0}$. Hence the result follows from translation invariance, Lemma 2.9 and (2.4). \square

From now on, t_0 will be a large enough multiple of $\frac{2}{v}$ such that both the inequality (2.4) and Lemma 2.10 hold.

3. The two-type contact process with equal birth rates. Let η_t denote the contact process with two types. For $A, B \subset \mathbb{Z}$ with $A \cap B = \emptyset$, $\{\eta_t^{A,B}, t \geq 0\}$ now denotes the contact process where at time zero A is the set of sites occupied by individuals of type 1, and B is the set of sites occupied by individuals of type 2. The dynamics is the same as before, using the same construction with the same collection of Poisson processes, except that now an individual of type $\alpha \in \{1, 2\}$ located at site z gives birth at time t to an individual of the same type at site $z + 1$ (resp. at site $z - 1$), if t is a point of the Poisson process $P^{x,2}$ (resp. $P^{x,3}$) and the site $z + 1$ (resp. $z - 1$) is not occupied at time t .

3.1. *A single mutant in front of an infinite number of residents may survive.* In this subsection, we consider the process $\{\eta_t^{A,B}, t \geq 0\}$ only in the case where $A < B$, meaning that all points in A are located on the left of each point of B . A consequence of the definition of our process is that a.s., for all $t > 0$, $\eta_t(x) = 1$ and $\eta_t(y) = 2$ imply that $x < y$. Whenever the process starts from one of these configurations then it remains in that set with probability 1.

For a configuration η in that set we define

$$\begin{aligned} br(\eta) &= \sup \{x : \eta(x) = 1\} \text{ and} \\ bl(\eta) &= \inf \{x : \eta(x) = 2\}. \end{aligned}$$

We now have the following consequence of Lemma 2.10 (here $\mathbb{P}^{A,B}$ denotes the law of $\{\eta_t^{A,B}, t \geq 0\}$):

COROLLARY 3.1. *For t_0 large enough, we have*

$$\mathbb{P}^{(-\infty, 0], \{vt_0\}} \left(\left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \right) \geq \frac{\rho^2}{2} - \varepsilon.$$

PROOF:

By Lemma 2.10 and symmetry arguments we have:

$$\mathbb{P}^{\{0\}, \{vt_0\}} \left(\left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \right) \geq \frac{\rho^2}{2} - \varepsilon$$

For t_0 large enough, on $C(v, t_0, \varepsilon)$ there is an open path from $(0, 0)$ to some point in $[-(v + \varepsilon)t_0, -(v - \varepsilon)t_0] \times \{t_0\}$. Any open path starting from $(vt_0, 0)$ remains strictly to the right of the previous path, since otherwise there would be an open path from $(vt_0, 0)$ to $[-(v + \varepsilon)t_0, -(v - \varepsilon)t_0] \times \{t_0\}$, which cannot occur on the event $C(v, t_0, \varepsilon)$. Consequently adding to the initial

configuration extra 1-type particles to the left of the origin does not alter the process to the right of that open path. Therefore

$$\begin{aligned} & \mathbb{P}^{(-\infty, 0], \{vt_0\}} \left(\left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \right) \\ &= \mathbb{P}^{\{0\}, \{vt_0\}} \left(\left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \right) \\ &\geq \frac{\rho^2}{2} - \varepsilon. \end{aligned}$$

□

To show that a similar result holds for the two type contact process on $(-\infty, \frac{3}{2}vt_0]$, we start with another lemma concerning the two type contact process on \mathbb{Z} :

LEMMA 3.2. *As $t_0 \rightarrow \infty$,*

$$\begin{aligned} & \mathbb{P}^{\{0\}, \{vt_0\}} \left(\left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \right) - \\ & \mathbb{P}^{\{0\}, \{vt_0\}} \left(\left\{ \exists x \in \left[\frac{vt_0}{2}, \frac{3vt_0}{4} \right]; \eta_{t_0}(x) = 2 \right\} \cap \left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \right) \end{aligned}$$

converges to 0.

PROOF: It suffices to show that

$$\mathbb{P}^{\{0\}, \{vt_0\}} \left(\left\{ \forall x \in \left(\frac{vt_0}{2}, \frac{3vt_0}{4} \right]; \eta_{t_0}(x) \neq 2 \right\} \cap \left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \right),$$

converges to 0 as t_0 goes to infinity. But on the event $\{br(\eta_{t_0}) \leq \frac{vt_0}{2}\}$ there are no 1's at time t_0 on the interval $[\frac{vt_0}{2} + 1, \frac{3vt_0}{4}]$, hence we only need to prove that for the one type contact process

$$\mathbb{P}^{\{vt_0\}} \left(\left\{ \forall x \in \left(\frac{vt_0}{2}, \frac{3vt_0}{4} \right]; \eta_{t_0}(x) = 0 \right\} \cap C(v, t_0, \varepsilon) \right)$$

converges to 0 as t_0 goes to infinity. But on the event $C(v, t_0, \varepsilon)$ the set of occupied points in the interval $(\frac{vt_0}{2}, \frac{3vt_0}{4}]$ is the same whether the initial condition of the process is \mathbb{Z} or $\{vt_0\}$. Since starting from \mathbb{Z} we have more occupied points than under the invariant measure the result follows from the fact that under the invariant measure the probability of having an empty interval of length n tends to 0 as n tends to infinity. □

Now we can prove:

COROLLARY 3.3. *Consider the two type contact process ζ_t on $(-\infty, \frac{3}{2}vt_0]$ and let $A = \left\{ \zeta : br(\zeta) \leq \frac{1}{2}vt_0 \right\}$. Then, taking t_0 large enough we have*

$$\mathbb{P}^{(-\infty, 0], \{vt_0\}} \left(\left\{ br(\zeta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap \left\{ \exists x : \frac{vt_0}{2} < x < \frac{3vt_0}{2}, \zeta_{t_0}(x) = 2 \right\} \right) \geq \frac{\rho^2}{2} - 2\varepsilon.$$

PROOF: In this proof we will consider the two type contact process on both \mathbb{Z} and $(-\infty, \frac{3}{2}vt_0]$. These two processes are constructed on the same probability space with the same Poisson processes. For the second of these processes $\{P_t^x; x > \frac{3}{2}vt_0\}$, $\{P_t^{x,-}; x > \frac{3}{2}vt_0\}$ and $\{P_t^{x,+}; x \geq \frac{3}{2}vt_0\}$ play no role.

On the set $\{\exists x \in [\frac{vt_0}{2}, \frac{3vt_0}{4}]; \eta_{t_0}(x) = 2\}$ there is an open path from $(vt_0, 0)$ to $[\frac{vt_0}{2}, \frac{3vt_0}{4}] \times \{t_0\}$. We now show that the probability that this path ever reaches the vertical line $\{x = \frac{3vt_0}{2}\}$ between time 0 and time t_0 converges to 0 as t_0 goes to infinity. Indeed, if that happened, there would be either an open path from $(vt_0, 0)$ to $\{\frac{3}{2}vt_0\} \times [0, \frac{3}{8}t_0]$ or an open path from $\{\frac{3}{2}vt_0\} \times [\frac{3}{8}t_0, t_0]$ to $[\frac{1}{2}vt_0, \frac{3}{4}vt_0] \times \{t_0\}$. The existence of the first of these paths has a probability which converges to 0 as t_0 goes to infinity by Lemma 2.2. By reversing the arrows and using symmetry and again Lemma 2.2, we see that the same happens to the second path.

Hence, if we define

$$B = \left\{ \exists \text{ an open path from } (vt_0, 0) \text{ to } \left[\frac{vt_0}{2}, \frac{3vt_0}{4} \right] \times \{t_0\} \right. \\ \left. \text{which remains to the left of the line } x = \frac{3vt_0}{2} \right\},$$

we deduce from Lemma 3.2 that

$$\mathbb{P}^{\{0\}\{vt_0\}} \left(C(v, t_0, \varepsilon) \cap \left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \right) - \mathbb{P}^{\{0\}\{vt_0\}} \left(B \cap C(v, t_0, \varepsilon) \cap \left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \right)$$

converges to 0 as t_0 goes to infinity. The result follows from Corollary 3.1 and the fact that starting both η_t and ζ_t from $(\{0\}\{vt_0\})$ we have:

$$B \cap \left\{ br(\eta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap C(v, t_0, \varepsilon) \\ \subset \left\{ br(\zeta_{t_0}) \leq \frac{vt_0}{2} \right\} \cap \left\{ \exists \frac{vt_0}{2} < x < \frac{3vt_0}{2}, \zeta_{t_0}(x) = 2 \right\}.$$

□

We now introduce the following partial order :

$\eta_1 \succeq \eta_2$ whenever both

$$\{x : \eta_1(x) = 2\} \subset \{x : \eta_2(x) = 2\} \text{ and } \{x : \eta_2(x) = 1\} \subset \{x : \eta_1(x) = 1\}.$$

Intuitively \succeq means “more 1’s” and “fewer 2’s”. Note that $\eta_1 \succeq \eta_2$ implies $br(\eta_1) \geq br(\eta_2)$ and that if $\gamma \succeq \zeta$, then we can couple two versions of our process in such a way that

$$(3.1) \quad \mathbb{P}(\eta_t^\gamma \succeq \eta_t^\zeta \ \forall t \geq 0) = 1.$$

This partial order extends to probability measures on the set of configurations: $\mu_1 \succeq \mu_2$ means that there exists a probability measure ν on $(\{0, 1, 2\}^{\mathbb{Z}})^2$ with marginals μ_1 and μ_2 such that $\nu(\{(\eta, \zeta) : \eta \succeq \zeta\}) = 1$.

In the sequel for any probability measure μ on $\{0, 1, 2\}^{\mathbb{Z}}$ and any $i \in \mathbb{N}$, $T^i(\mu)$ will denote the measure μ translated by i . That is the measure such that for all $n \in \mathbb{N}$, all $x_1 < x_2 < \dots < x_n$ and all possible values of a_1, \dots, a_n we have:

$$\begin{aligned} T^i(\mu)(\{\eta : \eta(x_1) = a_1, \dots, \eta(x_n) = a_n\}) = \\ \mu(\{\eta : \eta(x_1 - i) = a_1, \dots, \eta(x_n - i) = a_n\}) \quad (*). \end{aligned}$$

Moreover, if μ is a measure on $A^{[n, \infty)}$ where A is any subset of $\{0, 1, 2\}$, then $T^i(\mu)$ will be the measure on $A^{[n+i, \infty)}$ satisfying (*).

As before μ^+ denotes the upper invariant measure for the contact process on \mathbb{N} and μ_2^+ will be the measure obtained from μ^+ by means of the map: $F : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 2\}^{\mathbb{N}}$ given by $F(\eta)(x) = 2\eta(x)$. With a slight abuse of notation the measures μ^+ and μ_2^+ will also be seen as measures on $\{0, 1, 2\}^{\mathbb{N}}$ and a similar abuse of notation will be used for the translates of these measures.

We start the process $\{\eta_t, t \geq 0\}$ from the initial distribution $\bar{\mu}$ determined by

- (i) The projection of $\bar{\mu}$ on $\{0, 1, 2\}^{(-\infty, vt_0]}$ is the point mass on the configuration

$$\eta(x) = \begin{cases} 1, & \text{if } x \leq 0, \\ 0, & \text{if } 0 < x < vt_0, \\ 2, & \text{if } x = vt_0, \end{cases}$$

- (ii) the projection of $\bar{\mu}$ on $\{0, 1, 2\}^{[vt_0+1, \infty)}$ is $T^{vt_0}(\mu_2^+)$.

In the sequel η^0 will denote a random initial configuration distributed according to $\bar{\mu}$.

We now proceed as follows. We partition the probability space into a countable number of events: C, D_0, D_1, \dots and let the process run on a time interval of length t_0 . Then we show that the distribution of η_{t_0} conditioned on any event of the partition is \preceq than a convex combination of translations of $\bar{\mu}$. Hence the unconditioned distribution of η_{t_0} is also \preceq such a convex combination. Then we replace η_{t_0} by a random configuration η^1 whose distribution is this convex combination and let the process run on another time interval of length t_0 and so on.

For each $n \in \{\frac{3vt_0}{2}\} \cup \{2vt_0, 2vt_0 + 1, \dots\}$ we define two new processes: ${}_n\zeta_s$ on $\{0, 1, 2\}^{(-\infty, n]}$ and ${}_n\xi_s$ on $\{0, 2\}^{[n+1, \infty)}$. These evolve like the process η_t and are constructed with the same Poisson processes $P_t^{x,-}, P_T^{x,+}$ and P_t^x . For the first of these processes the Poisson processes $\{P_t^x : x > n\}$, $\{P_t^{x,+} : x \geq n\}$ and $\{P_t^{x,-} : x > n\}$ play no role. A similar statement holds for the second process. The initial distribution of these processes are the projections of $\bar{\mu}$ on $\{0, 1, 2\}^{(-\infty, n]}$ and $\{0, 1, 2\}^{[n+1, \infty)}$ respectively. In all cases considered later the second projection will concentrate on $\{0, 2\}^{[n+1, \infty)}$.

Our partition of the probability space is given by :

$$C = \left\{ br\left(\frac{3vt_0}{2}\zeta_{t_0}\right) \leq \frac{vt_0}{2}, \exists \frac{vt_0}{2} < x < \frac{3vt_0}{2} : \frac{3vt_0}{2}\zeta_{t_0}(x) = 2 \right\},$$

$$D_m = \{Q_{t_0} = vt_0 + m\} \cap C^C \quad \text{for } m = 0, 1, \dots,$$

where $Q_{t_0} = \max\{R_{t_0}, vt_0\}$ (recall that $R_t = \sup_{s \leq t} r_s$).

Note that on C

1. The set $\{x : \eta_{t_0}(x) = 1\}$ is contained in $(-\infty, \frac{vt_0}{2}]$ (indeed since $\{x, \frac{3vt_0}{2}\zeta_{t_0}(x) = 2\} \neq \emptyset, \{x, \eta_{t_0}(x) = 1\} = \{x, \frac{3vt_0}{2}\zeta_{t_0}(x) = 1\}$)
2. the set $\{x : \eta_{t_0}(x) = 2\}$ contains $\{x : \frac{3vt_0}{2}\xi_{t_0}(x) = 2\}$
3. The distribution of $\frac{3vt_0}{2}\xi_{t_0}$ is $\geq T^{\frac{3vt_0}{2}}\mu_2^+$ (this follows from Lemma 2.6).

Therefore, the distribution of η_{t_0} conditioned on C is $\preceq \nu$ where ν is determined by:

1. The projection of ν on $\{0, 1, 2\}^{(-\infty, \frac{3vt_0}{2}]}$ is the point mass on the configuration

$$\eta(x) = \begin{cases} 1, & \text{if } x \leq \frac{vt_0}{2}, \\ 0, & \text{if } \frac{vt_0}{2} < x \leq \frac{3vt_0}{2} \end{cases}$$

and

2. the projection of ν on $\{0, 1, 2\}^{[\frac{3vt_0}{2}+1, \infty)}$ is $T^{\frac{3vt_0}{2}}(\mu_2^+)$.

It follows from Lemma 2.7 (applied to μ_2^+ instead of μ^+) that if Y is a random variable such that

$$\mathbb{P}(Y = n) = \mu^+(\{\eta : \eta(x) = 0, x = 1, \dots, n-1, \eta(n) = 1\}),$$

then

$$\nu \preceq \sum_{n=1}^{\infty} \mathbb{P}(Y = n) T^{\frac{vt_0}{2}+n} \bar{\mu}.$$

Hence the distribution of η_{t_0} given C is $\preceq \sum_{n=1}^{\infty} \mathbb{P}(Y = n) T^{\frac{vt_0}{2}+n} \bar{\mu}$.

A similar argument shows that the conditional distribution of η_{t_0} given D_m is \preceq

$$\sum_{n=1}^{\infty} \mathbb{P}(Y = n) T^{vt_0+n+m} \bar{\mu},$$

where Y is distributed as above.

Putting our results together we get :

PROPOSITION 3.4. *Let $\bar{\mu}$ be the initial distribution of our process and let η^1 be a random configuration distributed as*

$$\mathbb{P}(C) \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \mathbb{P}(Y = n) T^{\frac{vt_0}{2}+n} \bar{\mu} + \sum_{m=0}^{\infty} \mathbb{P}(D_m) \sum_{n=1}^{\infty} \mathbb{P}(Y = n) T^{vt_0+m+n} \bar{\mu}.$$

Then $\eta_{t_0} \preceq \eta^1$.

PROPOSITION 3.5. *If t_0 is large enough and η^1 is distributed as in Proposition 3.4, then $w := (t_0)^{-1} \mathbb{E}(br(\eta^1) - br(\eta_0)) < v$.*

PROOF: Write

$$\begin{aligned}
\mathbb{E}[br(\eta^1) - br(\eta^0)] &= \sum_{n=1}^{\infty} \mathbb{P}(C) \mathbb{P}(Y = n) \left(\frac{vt_0}{2} + n \right) \\
&\quad + \sum_{m=0}^{\infty} \sum_{n=1}^{\infty} \mathbb{P}(C^c, Q_{t_0} = vt_0 + m) \mathbb{P}(Y = n) (vt_0 + n + m) \\
&\leq \mathbb{P}(C) \left[\frac{vt_0}{2} + \mathbb{E}(Y) \right] + \mathbb{P}(C^c) \mathbb{E}(Y) \\
&\quad + \sum_{m=0}^{\infty} \mathbb{P}(C^c, Q_{t_0} = vt_0 + m) (vt_0 + m) \\
&= \mathbb{P}(C) \frac{vt_0}{2} + \mathbb{E}(Y) + \mathbb{E}(Q_{t_0}) - \mathbb{E}(Q_{t_0}; C) \\
&\leq \mathbb{P}(C) \frac{vt_0}{2} + \mathbb{E}(Y) + \mathbb{E}(Q_{t_0}) - \mathbb{P}(C)vt_0 \\
&= \mathbb{E}(Q_{t_0}) + \mathbb{E}(Y) - \mathbb{P}(C) \frac{vt_0}{2}.
\end{aligned}$$

Hence it follows from Lemma 2.2 that

$$\limsup_{t_0 \rightarrow \infty} \frac{\mathbb{E}(br(\eta_{t_0}) - br(\eta^0))}{t_0} \leq v \left(1 - \frac{\mathbb{P}(C)}{2} \right).$$

□

We can now prove:

COROLLARY 3.6. *Let $\bar{\mu}$ be the initial distribution of the process. Then*

$$\limsup_{t \rightarrow \infty} \frac{br(\eta_t)}{t} \leq w \quad a.s.$$

PROOF: It follows from the last two propositions that for t_0 large enough, there exists an i.i.d. sequence of random variables $(X_n)_{n \geq 1}$ such that $\mathbb{E}(X_1) < vt_0$ and for all $n \geq 1$ $br(\eta_{nt_0}) - br(\eta_0) \leq \sum_{i=1}^n X_i$. The strong law of large numbers implies that the conclusion of the corollary holds along the sequence nt_0 , and the gaps are easy to control since for any initial configuration, the process $br(\eta_t) - br(\eta_0)$ is bounded above by a Poisson process of parameter λ .

□

It follows readily from this result that

COROLLARY 3.7.

$$\gamma := \mathbb{P}^{\mathbb{Z}^-, \{1\}}(\text{the type 2 population survives for ever}) > 0.$$

PROOF: First suppose that the initial distribution of the process is $\bar{\mu}$ and call η_0 the initial random configuration. It then follows from the above corollary that there exists $x > 0$ such that $\eta_0(x) = 2$ and there is an infinite open path starting at $(x, 0)$ such that for any (y, t) in this path we have $\eta_t(y) = 2$. This conclusion remains true if we suppress all the initial "2's" to the right of x . The corollary then follows from the Markov property and (3.1). \square

3.2. *A finite number of mutants do not survive in between a double infinity of residents.* The aim of this subsection is to prove

THEOREM 3.8. *Consider the two type contact process $\{\eta_t^{A,B}, t \geq 0\}$, where $|B| < \infty$, and the set A contains an infinite number of points located both to the left and to the right of B ,*

Then a. s. there exists $t > 0$ such that

$$\{x; \eta_s^{A,B}(x) = 2\} = \emptyset, \quad \forall s \geq t.$$

Let us first prove the following weaker statement. We shall then verify that the Theorem follows from it.

PROPOSITION 3.9. *For any $n, m \in \mathbb{N}$ let $A_{n,m} = \{x \in \mathbb{Z} : x \leq -m \text{ or } x \geq n\}$ and $B = \{0\}$, then a. s. there exists $t > 0$ such that*

$$\{x; \eta_s^{A_{n,m},B}(x) = 2\} = \emptyset, \quad \forall s \geq t.$$

PROOF: By the Markov property and (3.1) it suffices to prove the result for $n = m = 1$. Indeed starting from that configuration, for any $n, m > 1$, with positive probability we find ourselves at time one with the same unique type 2 individual located at $x = 0$, sites $-m + 1, \dots, -1$ empty, sites $1, \dots, n - 1$ empty, and some of the other sites occupied by type 1 individuals.

Let α_t denote the number of descendants at time t of the unique initial type 2 individual (hence α_t denotes also the number of type 2 individuals at time t). On the event that the lineage of the unique type 2 individual survives for ever we have $\alpha_t \rightarrow \infty$ as $t \rightarrow \infty$ a. s. Hence if that event has positive probability, $\mathbb{E}(\alpha_t) \rightarrow \infty$ as $t \rightarrow \infty$.

Denote by r'_t the rightmost descendant of the ancestor located at site 0 at time $t = 0$. If that individual has no descendants at time t , then $r'_t = -\infty$. In any case, $r'_t \leq r_t$, where r_t denotes the rightmost individual at time t , with all sites of \mathbb{Z}_- occupied and all sites of \mathbb{N} empty at time 0. Again we consider those various configurations with the same graphical construction. It is known (see e. g. Liggett [3] Theorem 2.19 page 281) that $r_t/t \rightarrow v$ a. s. and in $L^1(\Omega)$. Let now fix t large enough so that

1. $\mathbb{E}\left(\frac{r_t}{t}\right) \leq v + 1$;
2. $\mathbb{E}(\alpha_t) \geq 1 + \delta$, where $\delta > 0$.

Now by stationarity and 2., the expectation of the number of descendants at time t of the n (we assume that n is odd) ancestors located at points $-(n-1)/2, \dots, (n-1)/2$ at time 0 equals

$$n\mathbb{E}(\alpha_t) \geq n(1 + \delta).$$

On the other hand, from 1. and symmetry,

$$n\mathbb{E}(\alpha_t) \leq n + 2t(v + 1).$$

Choosing $n > 2t(v + 1)/\delta$, the last two inequalities yield a contradiction. \square

In order to deduce Theorem 3.8 from Proposition 3.9, we shall need the following Lemma.

LEMMA 3.10. *Let $(x_n)_{n \geq 0}$ be a strictly increasing sequence of strictly positive integers and let $(y_m)_{m \geq 0}$ be a strictly decreasing sequence of strictly negative integers. Then,*

$$\mathbb{P}(\exists n : \forall t > 0, \exists x : \eta^{\{x_n\}, \{x_{n-1}, x_{n-2}, \dots\}}(x) = 1) = 1,$$

and

$$\mathbb{P}(\exists m : \forall t > 0, \exists x : \eta^{\{y_m\}, \{y_{m+1}, y_{m+2}, \dots\}}(x) = 1) = 1.$$

PROOF: Define for $n, m \geq 0$ the events

$$\begin{aligned} C_n &= \{\forall t > 0, \exists x : \eta^{\{x_n\}, \{x_{n-1}, x_{n-2}, \dots\}}(x) = 1\}, \\ D_m &= \{\forall t > 0, \exists x : \eta^{\{y_m\}, \{y_{m+1}, y_{m+2}, \dots\}}(x) = 1\}. \end{aligned}$$

From Corollary 3.7, symmetry, translation invariance and (3.1),

$$\mathbb{P}(C_n) = \mathbb{P}(D_m) \geq \gamma \quad \forall n, m \geq 0.$$

On the set $\{(x, t) : x \in \mathbb{Z}, t \geq 0\}$ the Poisson processes used in the construction are n -fold mixing with respect to translations on \mathbb{Z} for any $n \in \mathbb{N}$. Since $x_{n+1} \geq x_n + 1$, this implies that for all $k \geq 1$

$$\limsup_{N \rightarrow \infty} \mathbb{P}\left(\bigcap_{j=0}^k C_{Nj}^c\right) \leq (1 - \gamma)^k.$$

Consequently

$$\mathbb{P}\left(\bigcap_{n \geq 0} C_n^c\right) \leq (1 - \gamma)^k$$

for all $k \geq 1$. This shows that

$$\mathbb{P}(\cup_{n \geq 0} C_n) = 1.$$

The result for the D_m 's is proved similarly. \square

PROOF OF THEOREM 3.8 By the Markov property, it suffices to consider the case where $A = \{y_n : n \in \mathbb{N}\} \cup \{x_n : n \in \mathbb{N}\}$, $B = \{0\}$ and the sequences (x_n) and (y_n) are as in the previous lemma.

For all $n, m \geq 1$, we define

$$E_{n,m} = \left\{ \forall t > 0, \exists x : \eta_t^{\{x_n\}, \{x_{n-1}, x_{n-2}, \dots\}}(x) = 1 \right\} \cap \left\{ \forall t > 0, \exists x : \eta_t^{\{y_m\}, \{y_{m+1}, y_{m+2}, \dots\}}(x) = 1 \right\}.$$

From the last lemma we know that $\mathbb{P}(\cup_{n,m} E_{n,m}) = 1$. Hence, it suffices to show that for all $n, m \in \mathbb{N}$, we have:

$$\mathbb{P}(\forall t > 0 \exists x : \eta_t^{A, \{0\}}(x) = 2, E_{n,m}) = 0.$$

But on the event $E_{n,m}$ the evolution of "2"'s is not altered by adding "1"'s to the left of y_m or to the right of x_n . Therefore the result follows from Proposition 3.9. \square

3.3. *Proof of Theorem 1.1.* The only if part follows from Theorem 3.8. Let us prove the if part.

We consider the case where $|A \cap B^+| < \infty$. The other case is treated similarly.

Define the set of configurations

$$\Lambda = \{\eta; \text{ s. t. } \eta(x) = 2, \text{ and } \eta(y) = 1 \text{ rightarrow } < x\}.$$

We let

$$T = \inf\{t \geq 0, \eta_t \in \Lambda\}.$$

Clearly $|A \cap B^+| < \infty$ implies that

$$\mathbb{P}^{A,B}(T < \infty) > 0.$$

Hence from the strong Markov property it remains to show that whenever $A \cap B^+ = \emptyset$,

$$\mathbb{P}^{A,B}(\text{the type 2 population survives for ever}) > 0.$$

This last statement follows from translation invariance, (3.1) and Corollary 3.7.

3.4. *Proof of Theorem 1.2.* By the Markov property and symmetry it suffices to show that the theorem holds for some A and B . To prove this, let $(x_n)_{n \geq 0}$ and $(y_m)_{m \geq 0}$ be as in the statement of Lemma 3.10 and let C_n and D_m be as in the proof of that lemma. It follows from that same lemma that there exist n and m such that $\mathbb{P}(C_n \cap D_m) > 0$. This implies that

$$\mathbb{P}\{y_m\}^{\{x_n\}}(\forall t > 0 \exists x, y : \eta_t(x) = 1, \eta_t(y) = 2) > 0.$$

Hence, the theorem holds when $A = \{x_n\}$ and $B = \{y_m\}$.

3.5. *Corollary for the one-type contact process.* The following is an immediate consequence of the above results.

COROLLARY 3.11. *Let $\{\xi_t^A, t \geq 0\}$ denote the one-type contact process starting from the configuration ξ_0 and let $A = \{x, \xi_0(x) = 1\}$. It follows from our results that*

1. *if A contains both a sequence which converges to $+\infty$ and a sequence which converges to $-\infty$, then no individual has a progeny which survives for ever;*
2. *if $|A| = +\infty$ but $\sup A < \infty$, then exactly one individual has a progeny which survives for ever.*

PROOF: The first statement is a consequence of Theorem 3.8. For the second statement first note that it follows from (3.1) and Corollary 3.7 that for any initial condition having a rightmost individual, the probability that this individual has a progeny which survives forever is bounded below by $\gamma > 0$. We then define an increasing sequence of stopping times: τ_1 is the smallest time at which the progeny of the rightmost initial individual dies out, τ_2 is the smallest time at which the progeny of the rightmost individual at time τ_1 dies out and so on. It then follows from a repeated application of the Strong Markov Property that $\mathbb{P}(\tau_n < \infty) \leq \gamma^n$. Hence, with probability 1 for some k , $\tau_k = \infty$ which implies that at least one individual has a progeny which survives forever. Suppose now that two individuals, say $x < y$, have a progeny which survives for ever with positive probability. Adding infinitely many individuals at time $t = 0$ on the right of y cannot possibly modify the fate of the progeny of x . This would mean that the progeny of x would survive for ever with positive probability, in the presence of infinitely many individuals at time $t = 0$ on both of its sides. This contradicts Theorem 3.8.

4. Remarks about the case of unequal birth rates. Let us first indicate the modification of the graphical representation which is required

in order to cover the case where the two types of individuals have distinct birth rates. Assume e. g. that type 1 individuals have a birth rate $\lambda > 0$, while type 2 individuals have a birth rate μ , where $\mu > \lambda$. Consider a collection $\{P_t^x, P_t^{x,+}, P_t^{x,-}, P_t^{x,+,*}, P_t^{x,-,*}, t \geq 0; x \in \mathbb{Z}\}$ of mutually independent Poisson point processes, such that the P^x 's have intensity 1, both the $P^{x,+}$'s and the $P^{x,-}$'s have intensity λ , and both the $P^{x,+,*}$'s and the $P^{x,-,*}$'s have intensity $\mu - \lambda$. At each time of the point process P^x , we place a δ on the line $\{y = x, t \geq 0\}$ in the half plane $\{(y, t); y \in \mathbb{R}, t \geq 0\}$. At each time t of the point process $P^{x,+}$ we place an arrow from (x, t) to $(x + 1, t)$. At each time t of the point process $P^{x,-}$ we place an arrow from (x, t) to $(x - 1, t)$. At each time t of the point process $P^{x,+,*}$ we place an arrow from (x, t) to $(x + 1, t)$ and mark it with a $*$. Finally at each time t of the point process $P^{x,-,*}$ we place an arrow from (x, t) to $(x - 1, t)$, and again mark it with a $*$.

We now need to define what is an open path in this framework.

An open path for the type 1 individuals in $[0, +\infty) \times \mathbb{Z}$ is a connected oriented path which moves along the time lines in the increasing t direction without passing through a δ symbol, and along birth arrows, in the direction of the arrow, but only following those arrows which are not marked, and which lead to a site where no individual is alive at that time. The $*$ -marked arrows are forbidden to type 1 individuals.

An open path for the type 2 individuals in $[0, +\infty) \times \mathbb{Z}$ is a connected oriented path which moves along the time lines in the increasing t direction without passing through a δ symbol, and along birth arrows, in the direction of the arrow. This time both types of arrows must be followed, provided they lead to a site where no individual is alive at that time.

We continue to denote by λ the birth rate of the type 1 population and denote now by μ the birth rate of the type 2 population. It is not hard to deduce from our argument that for any $\lambda > \lambda_c$, there exists $\varepsilon > 0$ such that for all $\lambda - \varepsilon < \mu < \lambda$, the if part of Theorem 1.1 remains true. This is true in particular if $\mu > \lambda_{cc}$, the critical birth rate for survival of the contact process allowed to give births only on one side, and any λ . In that case $\lambda - \mu$ can be arbitrarily large.

We conjecture that on the other hand, for any $\lambda > \lambda_c$, the if part of Theorem 1.1 fails when μ is too close to λ_c .

Concerning the only if part, we conjecture that whenever $\mu > \lambda$, a single mutant's progeny can survive for ever, even if all other sites are initially occupied by residents. This belief is based on the conjecture that again if $\mu > \lambda$ and the initial condition is the pair $(\mathbb{Z}_-, \mathbb{Z}_+)$, then the front between

both population moves asymptotically with a negative speed.

Acknowledgment We thank T.M. Liggett and J. Steif for useful discussions concerning the proof of Lemma 2.7.

References.

- [1] J van den Berg, O. Haggstrom and J.Kahn, Proof of a conjecture of N. Konno for the 1D contact process. In *Dynamics and Stochastics, Festschrift in Honor of Michael Keane*, MS Lecture notes-Monograph Series, Institute of Mathematical Statistics 2006
- [2] J. T. Cox, R. B. Schinazi, Survival and coexistence for a multitype contact process, *The Annals of Probability*, to appear.
- [3] Thomas M. Liggett, *Interacting particle systems*, Springer 1985.
- [4] Thomas M. Liggett, *Stochastic interacting systems: contact, voter and exclusion processes*, Springer 1999.
- [5] Claudia Neuhauser, Ergodic theorems for the multitype contact process, *Probab. Theory Related Fields* **91**, 467–506, 1992.

LATP, EBM
39, RUE F. JOLIOT CURIE
F 13453 MARSEILLE CEDEX 13
E-MAIL: enrique.andjel@cmi.univ-mrs.fr
E-MAIL: etienne.pardoux@cmi.univ-mrs.fr

DEPARTMENT OF MATHEMATICS
GEORGETOWN UNIVERSITY
WASHINGTON DC 20057 USA
E-MAIL: jrm32@georgetown.edu