	Equation to be solved			Perspectives
	000 0	00 00000 00000		

A Finite Volume Method for a Distributed Order Diffusion Equation

Fayssal Benkhaldoun¹ and Abdallah Bradji^{1,2}

¹ LAGA (Laboratoire d'Analyse, Géométrie, et Applications)-Paris 13–France
² Department of Mathematics, University of Annaba–Algeria

ICNTAM 2022 (International Conference on New Trends in Applied Mathematics) May 19–21, 2022, Béni Mellal- Morocco On-Line Presentation



< ロ > < 同 > < 三 > < 三 >

Aim		Equation to be solved			Perspectives
•		000 0	00 00000 00000		

Aim of the presentation

The aim of this talk is to establish a Finite Volume Scheme for a Distributed Order Diffusion Equation and prove its convergence.



Plan	Equation to be solved			Perspectives
•	000 0	00 00000 00000		

Plan of this presentation

- Some References
- 2 Equation to be solved
- Introduction: Finite Volume methods from Admissible to Nonconforming meshes (SUSHI scheme)

- 4 Overview on the approximations of Distributed Caputo derivative
- 5 Formulation of a Finite Volume Scheme
- 6 Statement of the Convergence of the numerical scheme
- 7 Perspectives

	Some references	Equation to be solved 000 0	Introduction OO OOOOO OOOOO	Formulation of the scheme	

Some references

- A. Bradji and J. Fuhrmann. Convergence order of a finite volume scheme for the time-fractional diffusion equation. Numerical analysis and its applications, 33–45, Lecture Notes in Comput. Sci., 10187, Springer, Cham, 2017.
- K. Diethelm and N. Ford. Numerical analysis for distributed-order differential equations. J. Comput. Appl. Math. 225 (2009), no. 1, 96–104.
- R. Eymard, T. Gallouët, R. Herbin. Discretization of heterogeneous and anisotropic diffusion problems on general nonconforming meshes. IMA J. Numer. Anal. 30/4, 1009–1043 (2010).
- G. H. Gao, H. W. Sun, and Z. Z. Sun. Some high-order difference schemes for the distributed-order differential equations. J. Comput. Phys. 298 (2015), 337–359.
- H. Ye, F. Liu, V. Anh, and I. Turner. Numerical analysis for the time distributed-order and Riesz space fractional diffusions on bounded domains. IMA J. Appl. Math. 80 (2015), no. 3, 825–838.



	Equation to be solved			Perspectives
	• 00 0	00 00000 00000		

Problem to be solved

Equation

We consider the following time fractional diffusion equation:

$$\mathbb{D}_t^{\alpha} u(\mathbf{x}, t) - \Delta u(\mathbf{x}, t) = f(\mathbf{x}, t), \quad (\mathbf{x}, t) \in \Omega \times (0, T),$$
(1)

(2)

E

(日)

where $\Omega \subset \mathbb{R}^d$ is an open domain of \mathbb{R}^d , T > 0, and f is a given function with:

Distributed Caputo derivative

$$\mathbb{D}^{lpha}_t arphi(t) = \int_0^1 \omega(lpha) \partial^{lpha}_t arphi(t) dlpha$$

with
$$\partial_t^{\alpha} \varphi(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-s)^{-\alpha} \varphi' ds$$
 for $0 \le \alpha < 1$ and $\partial_t^{\alpha} \varphi(t) = \varphi'(t)$
when $\alpha = 1$.

	Equation to be solved			Perspectives
	0 ● 0 ○	00 00000 00000		

Problem to be solved: Suite

Hypotheses on the weight function ω

$$\omega(\alpha) \ge 0$$

and

$$\int_0^1 \omega(\alpha) d\alpha = c_0 > 0.$$

(日)

(3)

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

	Equation to be solved			Perspectives
	00 ● 0	00 00000 00000		

Initial and boundary conditions

Initial condition

$$u(\boldsymbol{x},0) = u^0(\boldsymbol{x}), \quad \boldsymbol{x} \in \Omega.$$

Homogeneous Dirichlet boundary

$$u(\mathbf{x},t) = 0, \quad (\mathbf{x},t) \in \partial \Omega \times (0,T).$$

(日)

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

			Equation to be solved				Perspectives			
			000 •	00 00000 00000						
What abou	What about time fractional diffusion equation?									

What about time fractional diffusion equation?

Some physics

Fractional differential equations have been successfully used in the modeling of many different processes and systems. They are used, for instance, to describe anomalous transport in disordered semiconductors, penetration of light beam through a turbulent medium, transport of resonance radiation in plasma, blinking fluorescence of quantum dots, penetration and acceleration of cosmic ray in the Galaxy, and large-scale statistical Cosmography. We refer to the monograph Uchaikin (Fractional Derivatives for Physicists and Engineers, Springer-Verlag Heidelberg, 2013) where we find many details.



< < >> < <</p>

	Equation to be solved	Introduction		Perspectives
	000 0	• 0 00000 00000		

Introduction: Finite Volume from Admissible meshes to Nonconforming meshes

Finite volume methods are numerical methods approximating different types of Partial Differential Equations (PDEs). They are based on three principle ideas:

- Subdivision of the spatial domain into subsets called Control Volumes.
- Integration of the equation to be solved over the Control Volumes.
- Approximation of the derivatives appearing after integration.



¹We mean here the "pure" finite volume methods and not finite volume-element methods > < = >

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

		Introduction		
	000 0	00 00000 00000		

Introduction (suite): Finite Volume from Admissible meshes to Nonconforming meshes

Finite Volume methods passed by three steps:

First step

Finite Volume methods using Admissible meshes.

Second step

SUSHI method.



イロト イ団ト イヨト イヨト

		Introduction		
	000 0	00 00000 00000		

Introduction (suite): Finite Volume from Admissible meshes to Nonconforming meshes

Finite Volume methods passed by three steps:

First step

Finite Volume methods using Admissible meshes.

Second step

SUSHI method.



<ロト <回 > < 回 > < 回 > < 回 >

		Introduction		
	000 0	00 00000 00000		

Introduction (suite): Finite Volume from Admissible meshes to Nonconforming meshes

Finite Volume methods passed by three steps:

First step

Finite Volume methods using Admissible meshes.

Second step

SUSHI method.

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022 Finite Volumes for Distributed Fractional



		Equation to be solved 000 0	Introduction	Formulation of the scheme	
Einite Mele	 an adminible machine		00000		

Introduction (suite): Finite Volume methods on admissible meshes

Definition

Let \mathcal{T} be an Admissible Mesh in the sense of Eymard et al. (Handbook, 2000).

 $K \in \mathcal{T}$ are the control volumes and σ are the edges of the control volumes K.



 $T_{K,L} = \frac{m_{K,L}}{d_{K,L}}$

Figure: transmissivity between K and L: $\mathcal{T}_{\sigma} = \mathcal{T}_{K|L} = \frac{m_{K,L}}{d_{K,L}}$



			Equation to be solved	Introduction			Perspectives
			000 0	00 00000 00000			
Finite Volume methods on admissible meshes							

Introduction (suite): Finite Volume methods on admissible meshes

Main properties of Admissible mesh:

- Convexity of the Control Volumes.
- **2** The orthogonality property: the $(x_K x_L)$ is orthogonal to the common edge σ between the control volumes *K* and *L*.



			Equation to be solved	Introduction			Perspectives				
			000	00 00000 00000							
Einite Volu	Cinita Valuma mathada an admissibla machaa										

Introduction (suite): Finite Volume methods on admissible meshes

Model to be solved:

$$-\Delta u(\mathbf{x}) = f(\mathbf{x}), \quad \mathbf{x} \in \Omega \quad \text{and} \quad u(\mathbf{x}) = 0, \quad \mathbf{x} \in \partial \Omega.$$
(5)

Principles of Finite Volume scheme:

Integration on each control volume
$$K := \int_{K} \Delta u(\mathbf{x}) d\mathbf{x} = \int_{K} f(\mathbf{x}) d\mathbf{x}$$
,

2 Integration by Parts gives :-
$$\int_{\partial K} \nabla u(\mathbf{x}) \cdot \mathbf{n}(\mathbf{x}) d\gamma(\mathbf{x}) = \int_{K} f(\mathbf{x}) d\mathbf{x}$$

Summing on the lines of
$$K$$
: $-\sum_{\sigma \in \mathcal{E}_K} \int_{\sigma} \nabla u(\mathbf{x}) \cdot \mathbf{n}(\mathbf{x}) d\gamma(\mathbf{x}) = \int_K f(\mathbf{x}) d\mathbf{x}$



(日)

	Equation to be solved	Introduction		Perspectives
	000 0	00 00000 00000		

Introduction (suite): Finite Volume methods on admissible meshes

Approximate Finite Volume Solution $u_{\mathcal{T}} = (u_K)_K$

$$-\sum_{\sigma\in\mathcal{E}_{K}}\frac{\mathbf{m}(\sigma)}{d_{K|L}}(u_{L}-u_{K})=\int_{K}f(\mathbf{x})d\mathbf{x}.$$

Matrix Form

$$\mathcal{A}^{\mathcal{T}} u_{\mathcal{T}} = f_{\mathcal{T}}.$$



(日)

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

	Equation to be solved	Introduction		Perspectives
	000 0	00 00000 00000		

Introduction (suite): Finite Volume methods on admissible meshes

Approximate Finite Volume Solution $u_{\mathcal{T}} = (u_K)_K$

$$-\sum_{\sigma\in\mathcal{E}_K}\frac{\mathbf{m}(\sigma)}{d_{K|L}}(u_L-u_K)=\int_K f(\mathbf{x})d\mathbf{x}.$$

Matrix Form

$$\mathcal{A}^{\mathcal{T}} u_{\mathcal{T}} = f_{\mathcal{T}}.$$



(日)

(6)

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

	Equation to be solved	Introduction		Perspectives
	000 0	00 00000 00000		

Introduction (suite): Finite Volume methods on admissible meshes

Approximate Finite Volume Solution $u_{\mathcal{T}} = (u_K)_K$

$$-\sum_{\sigma\in\mathcal{E}_K}\frac{\mathbf{m}(\sigma)}{d_{K|L}}(u_L-u_K)=\int_K f(\mathbf{x})d\mathbf{x}.$$

Matrix Form

$$\mathcal{A}^{\mathcal{T}} u_{\mathcal{T}} = f_{\mathcal{T}}.$$



(日)

(6)

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

	Equation to be solved	Introduction		Perspectives
	000 0	00 00000 00000		

Introduction (suite): Finite Volume methods on admissible meshes

Theorem

Let $\mathcal{X}(\mathcal{T})$: functions which are constant on each control volume K. Let $e_{\mathcal{T}} \in \mathcal{X}(\mathcal{T})$ be defined by $e_K = u(\mathbf{x}_K) - u_K$ for any $K \in \mathcal{T}$. Assume that the exact solution usatisfies $u \in C^2(\overline{\Omega})$. Then the following convergence results hold:

 \blacksquare H_0^1 -error estimate

$$\|e_{\mathcal{T}}\|_{1,\mathcal{T}} \le Ch \|u\|_{2,\overline{\Omega}},\tag{7}$$

where
$$\|\cdot\|_{1,\mathcal{T}}$$
 is the H_1^0 -norm $\|e_{\mathcal{T}}\|_{1,\mathcal{T}}^2 = \sum_{\sigma=K|L\in\mathcal{E}} \frac{\mathrm{m}(\sigma)}{d_{\sigma}} (u_L - u_K)^2$.

2 L^2 -error estimate:

$$\|e_{\mathcal{T}}\|_{L^2(\Omega)} \leq Ch \|u\|_{2,\overline{\Omega}}.$$

	Equation to be solved	Introduction		Perspectives
	000 0	00 00000 00000		

Introduction (suite): Finite Volume methods using nonconforming grids, SUSHI scheme

Definition (New mesh of Eymard et al., IMAJNA 2010)



Figure: Notations for two neighbouring control volumes in d = 2



(a)

	Equation to be solved	Introduction		Perspectives
	000 0	00 00000 00000		

Introduction (suite): Finite Volume methods using nonconforming grids, SUSHI scheme

Main properties of this new mesh:

- I (mesh defined at any space dimension): $\Omega \subset \mathbb{R}^d$, $d \in \mathbb{N}$
- (orthogonality property is not required): the orthogonality property is not required in this new mesh. But, additional discrete unknowns are required.
- (convexity): the classical admissible mesh should satisfy that the control volumes are convex, whereas the convexity property is not required in this new mesh.



< < >> < <</p>

	Equation to be solved	Introduction		Perspectives
	000	00 00000 00000		

Introduction (suite): Finite Volume methods using nonconforming grids, SUSHI scheme

Principles of discretization for Poisson's equation:

Discrete unknowns: the space of solution as well as the space of test functions are in

$$\mathcal{X}_{\mathcal{D},0} = \{ \left(\left(v_K \right)_{K \in \mathcal{M}}, \left(v_\sigma \right)_{\sigma \in \mathcal{E}} \right), \ v_K, v_\sigma \in \mathbb{R}, \ v_\sigma = 0, \, \forall \sigma \in \mathcal{E}_{\text{ext}} \}$$

- **2** Discretization of the gradient: the discretization of ∇ can be performed using a stabilized discrete gradient denoted by $\nabla_{\mathcal{D}}$, see Eymard et *al.* (IMAJNA, 2010):
 - 1 The discrete gradient $\nabla_{\mathcal{D}}$ is stable
 - 2 The discrete gradient $\nabla_{\mathcal{D}}$ is consistent.



	Equation to be solved	Introduction		Perspectives
	000	00 00000 00000		

Introduction (suite): Finite Volume methods using nonconforming grids, SUSHI

Weak formulation for Poisson's equation: Find $u \in H_0^1(\Omega)$ such that

$$\int_{\Omega} \nabla u(\mathbf{x}) \cdot \nabla v(\mathbf{x}) d\mathbf{x} = \int_{\Omega} f(\mathbf{x}) v(\mathbf{x}) d\mathbf{x}, \quad \forall v \in H_0^1(\Omega).$$
(9)

SUSHI (Scheme Using stabilized Hybrid Interfaces) for Poisson's equation: Find $u_{\mathcal{D}} \in \mathcal{X}_{\mathcal{D},0}$ such that

$$\int_{\Omega} \nabla_{\mathcal{D}} u_{\mathcal{D}}(\boldsymbol{x}) \cdot \nabla_{\mathcal{D}} v(\boldsymbol{x}) d\boldsymbol{x} = \int_{\Omega} f(\boldsymbol{x}) v(\boldsymbol{x}) d\boldsymbol{x}, \quad \forall v \in \mathcal{X}_{\mathcal{D},0}.$$



ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

	Equation to be solved	Introduction		Perspectives
	000 0	00 00000 00000		

Introduction (suite): Finite Volume methods using nonconforming grids, SUSHI

Theorem

Assume that the exact solution u satisfies $u \in C^2(\overline{\Omega})$. Then the following convergence result hold:

1 H_0^1 -error estimate

$$\|\nabla u - \nabla_{\mathcal{D}} u_{\mathcal{D}}\|_{L^{2}(\Omega)^{d}} \le Ch \|u\|_{2,\overline{\Omega}}.$$
(11)

(12)

2 L^2 -error estimate:

$$\|u - \Pi_{\mathcal{M}} u_{\mathcal{D}}\|_{L^2(\Omega)} \le Ch \|u\|_{2,\overline{\Omega}}.$$

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

			Formulation of the scheme	
	000 0	00 00000 00000	0000	

Principles of the time- discretization

Step 1: Discretize the integral (2).

We discretize [0, 1] with uniform mesh $\{\xi = i/q; i = 0, ..., q\}$ with constant time step 1/q where $q \in \mathbb{N} \setminus \{0\}$.

$$\mathbb{D}_{t}^{\alpha}\varphi(t) \approx \frac{1}{q}\sum_{i=0}^{q-1}\lambda_{i}\omega(\alpha_{i})\partial_{t}^{\alpha_{i}}\varphi(t),$$
(13)

(日)

with $\lambda_i > 0$ and $\alpha_0 < \alpha_1 < \ldots < \alpha_q$.

Nice remark

The RHS of (13) is a Multi-Term Fractional Operator.

	Equation to be solved		Formulation of the scheme	Perspectives
	000 0	00 00000 00000	0000	

Examples of quadratures

Examples of quadratures

- Composite Mid point Formula
- Composite Trapezoid Formula
- Composite Simpson Formula



<ロト <回 > < 回 > < 回 > < 回 >

			Formulation of the scheme	
	000 0	00 00000 00000	00000	

Principles of the time-discretization: Suite

Step 2: Discretize Caputo derivatives.

We define k = T/(M + 1) and $t_n = nk, n = 0, ..., M + 1$.

$$\partial_t^{\alpha_i}\varphi(t_{n+1}) = \frac{1}{\Gamma(2-\alpha)} \sum_{j=0}^n d_{j,\alpha_i} \overline{\partial}^{\alpha_i}\varphi(t_{n-j+1}) + \mathbb{T}_{\alpha_i}^{n+1},$$
(14)

where $\overline{\partial}^{\alpha_i}$ is the following discrete derivative of order α

$$\overline{\partial}^{\alpha_i} v^{j+1} = \frac{v^{j+1} - v^j}{k^{\alpha_i}} \quad \text{and} \quad d_{j,\alpha_i} = (j+1)^{1-\alpha_i} - j^{1-\alpha_i} \tag{15}$$

(16)

・ロト ・四ト ・ヨト ・ヨト

and

$$\begin{split} |\mathbb{T}^{n+1}_{lpha_i}| &\leq Ck^{2-lpha_i} \ &\leq Ck^{2-lpha_q}, \quad ext{for} \quad k\leq 1. \end{split}$$

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

			Formulation of the scheme	
	000 0	00 00000 00000	00000	

Principles of the discretization (suite)

Discretization in space

We use SUSHI scheme



(日)

	Equation to be solved 000 0	Introduction 00 00000 00000	Formulation of the scheme 0000●	

Formulation of scheme

The finite volume scheme can then be defined as:

Discretization of initial condition: Find $u_{\mathcal{D}}^0 \in \mathcal{X}_{\mathcal{D},0}$ such that

$$\left(\nabla_{\mathcal{D}} u_{\mathcal{D}}^{0}, \nabla_{\mathcal{D}} v\right)_{\left(\mathbb{L}^{2}(\Omega)\right)^{d}} = -\left(\Delta u^{0}, \Pi_{\mathcal{M}} v\right)_{\mathbb{L}^{2}(\Omega)}, \quad \forall v \in \mathcal{X}_{\mathcal{D},0}, \qquad (17)$$

Discretization of the factional heat equation: and for any $n \in [\![0,M]\!]$, find $u_{\mathcal{D}}^{n+1} \in \mathcal{X}_{\mathcal{D},0}$ such that, for all $v \in \mathcal{X}_{\mathcal{D},0}$

$$\frac{1}{q\Gamma(2-\alpha)}\sum_{i=0}^{q-1}\lambda_{i}\omega(\alpha_{i})\sum_{j=0}^{n}d_{j,\alpha_{i}}\left(\overline{\partial}^{\alpha_{i}}\Pi_{\mathcal{M}}u_{\mathcal{D}}^{n+1},\Pi_{\mathcal{M}}v\right)_{\mathbb{L}^{2}(\Omega)} + \left(\nabla_{\mathcal{D}}u_{\mathcal{D}}^{n+1},\nabla_{\mathcal{D}}v\right)_{\left(\mathbb{L}^{2}(\Omega)\right)^{d}} = (f(t_{n+1}),\Pi_{\mathcal{M}}v)_{\mathbb{L}^{2}(\Omega)}.$$

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

			Convergence result	
	000 0	00 00000 00000	•0	

Convergence result

Theorem (An $L^{\infty}(L^2)$ -error estimate)

For smooth solution u and weight function ω

$$\|u(t_n) - \Pi_{\mathcal{M}} u_{\mathcal{D}}^n\|_{\mathbb{L}^2(\Omega)} \le C\left(h_{\mathcal{D}} + k^{2-\alpha_q} + \mathbb{T}(q)\right),\tag{19}$$

where $\mathbb{T}(q)$ is quadrature error in the approximation:

$$\int_0^1 \psi(\alpha) d\alpha \approx \frac{1}{q} \sum_{i=0}^{q-1} \lambda_i \psi(\alpha_i).$$

<ロト <回 > < 回 > < 回 > < 回 >

ICNTAM Talk of Benkhaldoun, Bradji, May 21st-2022

			Convergence result	
	000 0	00 00000 00000	00	

Convergence result in a particular case: Mid Point formula

Corollary (An $L^{\infty}(L^2)$ -error estimate)

For Composite Mid Point formula and smooth solution u and weight function ω

$$\|u(t_n) - \Pi_{\mathcal{M}} u_{\mathcal{D}}^n\|_{\mathbb{L}^2(\Omega)} \le C \left(h_{\mathcal{D}} + k^{1+1/(2q)} + q^{-2}\right).$$
(20)



(a)

				Perspectives
	000 0	00 00000 00000		•

Perspectives

First perspective

Convergence analysis in other discrete norms

Second perspective

Gradient schemes framework for Distributed Time Fractional Diffusion Equations



(日)

				Perspectives
	000 0	00 00000 00000		•

Perspectives

First perspective

Convergence analysis in other discrete norms

Second perspective

Gradient schemes framework for Distributed Time Fractional Diffusion Equations



(日)

				Perspectives
	000 0	00 00000 00000		•

Perspectives

First perspective

Convergence analysis in other discrete norms

Second perspective

Gradient schemes framework for Distributed Time Fractional Diffusion Equations



<ロト <回 > < 回 > < 回 > < 回 >