

# Quasiadditivity and measure property of capacity and the tangential boundary behavior of harmonic functions

*H. Aikawa and A. A. Borichev*

**Abstract.** A certain quasiadditive property of capacity is shown. Namely, it is proved that if a set  $E$  is dispersely decomposed into subsets, then the capacity of  $E$  is comparable to the summation of the capacities of the subsets. From the quasiadditivity it is derived that the Lebesgue measure of a certain expanded set is estimated by the capacity of the original set. The estimation has an application to the boundary behavior of harmonic functions.

1991 Mathematics Subject Classification: Primary 31B15, 31B25

## 1. Introduction

Throughout this article we denote by  $A$  a positive constant whose value is unimportant and may change from line to line. We say that two positive quantities  $f$  and  $g$  are comparable, written  $f \approx g$ , if there exists a constant  $A$  such that  $A^{-1}g \leq f \leq Ag$ . We say that a capacity  $C$  is quasiadditive if

$$C(E) \approx \sum C(E_j)$$

for some decomposition  $E = \bigcup E_j$ . In [3] and [4] we considered the quasiadditivity for certain capacities with respect to the Whitney decomposition. Here we give a different type of quasiadditivity.

Let  $K(r) \not\equiv 0$  be a nonnegative nonincreasing lower semicontinuous (l. s. c.) function for  $r > 0$ . We assume that

$$\lim_{r \rightarrow 0} K(r) = \infty \text{ and } \lim_{r \rightarrow \infty} K(r) = 0.$$

For  $x \in \mathbb{R}^N$  we define  $K(x) = K(|x|)$ , and assume that  $K(x)$  is integrable over  $\mathbb{R}^N$ . We define the capacity  $C_K$  by

$$C_K(E) = \inf\{\|\mu\| : K * \mu \geq 1 \text{ on } E\}.$$

Define a positive function  $\eta(r)$  by  $|B(0, \eta(r))| = C_K(B(0, r))$  and put  $\eta^*(r) = \max\{\eta(r), 2r\}$ .

**Theorem 1.** *Suppose  $\{B(x_j, \eta_p^*(r_j))\}$  is disjoint and  $E$  is an analytic subset of  $\bigcup B(x_j, r_j)$ . Then*

$$C_K(E) \approx \sum C_K(E \cap B(x_j, r_j)).$$

Theorem 1 has a counterpart in  $L^p$ -capacity and in energy capacity. Let  $1 < p < \infty$ . We define

$$C_{K,p}(E) = \inf\{\|f\|_p^p : K * f \geq 1 \text{ on } E, f \geq 0\}.$$

Define a positive function  $\eta_p(r)$  by  $|B(0, \eta_p(r))| = C_{K,p}(B(0, r))$  and put  $\eta_p^*(r) = \max\{\eta_p(r), 2r\}$ .

**Theorem 2.** *Suppose  $\{B(x_j, \eta_p^*(r_j))\}$  is disjoint and  $E$  is an analytic subset of  $\bigcup B(x_j, r_j)$ . Then*

$$C_{K,p}(E) \approx \sum C_{K,p}(E \cap B(x_j, r_j)).$$

The energy capacity is defined by

$$e_K(E) = \sup\{\|\mu\|^2 : \mu \text{ is concentrated on } E, \int K * \mu d\mu \leq 1\}.$$

We define  $\eta_e(r)$  by  $|B(0, \eta_e(r))| = e_K(B(0, r))$  and put  $\eta_e^*(r) = \max\{\eta_e(r), 2r\}$ .

**Theorem 3.** *Suppose  $\{B(x_j, \eta_e^*(r_j))\}$  is disjoint and  $E$  is an analytic subset of  $\bigcup B(x_j, r_j)$ . Then*

$$e_K(E) \approx \sum e_K(E \cap B(x_j, r_j)).$$

From Theorems 1–3 we can deduce the following measure property of  $C_K$ ,  $C_{K,p}$  and  $e_K$ . For notational convenience we extend  $C_{K,p}$ ,  $\eta_p$  and  $\eta_p^*$  for  $p \geq 1$ . Thus,  $C_{K,1}$ ,  $\eta_1$  and  $\eta_1^*$  mean  $C_K$ ,  $\eta$  and  $\eta^*$ , respectively. We put

$$\tilde{E}_{K,p} = \bigcup_{x \in E} B(x, \eta_p^*(\delta_E(x))), \quad \tilde{E}_{K,e} = \bigcup_{x \in E} B(x, \eta_e^*(\delta_E(x))).$$

**Theorem 4.** *Let  $p \geq 1$ . There is a positive constant  $A$  depending only on  $N$ ,  $K$  and  $p$  such that*

$$\begin{aligned} |\tilde{E}_{K,p}| &\leq AC_{K,p}(E), \\ |\tilde{E}_{K,e}| &\leq Ae_K(E). \end{aligned}$$

Theorem 4 has an application to the tangential boundary behavior of harmonic functions. We shall later give Theorem 5, a generalization of Theorem 4, in connection with Nagel-Stein approach regions ([10]). We shall introduce a notion of

“thin sets” to obtain precise description of the tangential boundary behavior of harmonic functions given as the Poisson integral of certain potentials. We shall combine it with Theorem 5 and observe that [9, Theorem 2.9] follows.

The plan of this article is as follows: We prove Theorems 1–3 in the next section. Since the proofs are similar, we shall give a complete proof only for Theorem 3. For Theorems 1 and 2 we refer the reader [6]. In Section 3 we shall prove Theorem 4 by using the usual covering lemma. We shall also indicate that if we invoke the covering lemma due to Nagel-Stein [10], then we obtain Theorem 5, a generalization of Theorem 4. In Section 4 we shall give some applications of Theorem 5 to the boundary behavior of harmonic functions.

## 2. Proof of Theorems 1–3

We prepare an elementary lemma.

**Lemma 1.** *Let  $0 < 2r \leq R$ . Suppose  $x \notin B(x_0, R)$  and let  $\rho = \text{dist}(x, B(x_0, r))$ . Then*

$$|B(x, \rho) \cap B(x_0, R)| \geq A|B(x_0, R)|,$$

where  $A$  depends only on the dimension.

*Proof.* Let  $x_1$  be the point on the line segment connecting  $x_0$  and  $x$  such that  $|x_1 - x_0| = \frac{3}{4}R$ . It is easy to see that

$$\rho = |x - x_0| - r = |x - x_1| + \frac{3}{4}R - r \geq |x - x_1| + \frac{1}{4}R.$$

We observe that

$$B(x_1, \frac{1}{4}R) \subset B(x, \rho) \cap B(x_0, R),$$

since if  $y \in (x_1, \frac{1}{4}R)$ , then

$$|x - y| < |x - x_1| + \frac{1}{4}R \leq \rho \text{ and } |x_0 - y| < |x_0 - x_1| + \frac{1}{4}R = R.$$

Thus the required inequality follows.  $\square$

*Proof of Theorem 3.* For simplicity we write  $E_j$  for  $E \cap B(x_j, r_j)$ . It is sufficient to show that  $\sum e_K(E_j) \leq A e_K(E)$ . Let  $\mu_j$  be the  $e_K$ -equilibrium measure for  $E_j$ , i.e.

$$\begin{aligned} \mu_j &\text{ is concentrated on } E_j, \\ \int K * \mu_j d\mu_j &= 1, \\ \|\mu_j\|^2 &= e_K(E_j). \end{aligned}$$

Let  $\mu_j^* = e_K(E_j)^{1/2}\mu_j$  and let  $\mu^* = \sum \mu_j^*$ . We observe that

$$\int K * \mu_j^* d\mu_j^* = \|\mu_j^*\| = e_K(E_j) \text{ and } \|\mu^*\| = \sum e_K(E_j). \quad (1)$$

We put

$$f_j = \frac{\|\mu_j^*\|}{|B(x_j, \eta_e^*(r_j))|} \chi_{B(x_j, \eta_e^*(r_j))},$$

$d\mu'_j = f_j dx$  and  $\mu' = \sum \mu'_j$ . Then  $\|\mu'_j\| = \|\mu_j^*\| = e_K(E_j)$  and  $f_j \leq \chi_{B(x_j, \eta_e^*(r_j))}$ , since  $|B(x_j, \eta_e^*(r_j))| \geq |B(x_j, \eta_e(r_j))| = e_K(B(x_j, r_j)) \geq e_K(E_j) = \|\mu_j^*\|$ . We observe from the disjointness of  $B(x_j, \eta_e^*(r_j))$  that  $\sum f_j \leq \sum \chi_{B(x_j, \eta_e^*(r_j))} \leq 1$ . Hence

$$K * \mu' = K * \left( \sum f_j \right) \leq K * 1 = \int K dx < \infty. \quad (2)$$

Let us compare  $K * \mu^*$  and  $K * \mu'$ . Suppose first  $x \notin B(x_j, \eta_e^*(r_j))$ . We apply Lemma 1 with  $x_0 = x_j$ ,  $R = \eta_e^*(r_j)$  and  $r = r_j$ . Let  $\rho_j = \text{dist}(x, B(x_j, r_j))$ . We have

$$\begin{aligned} K * \mu'_j(x) &\geq \int_{B(x, \rho_j) \cap B(x_j, \eta_e^*(r_j))} K(x-y) d\mu'_j(y) \\ &\geq K(\rho_j) \mu'_j(B(x, \rho_j) \cap B(x_j, \eta_e^*(r_j))) \\ &= K(\rho_j) \|\mu_j^*\| \frac{|B(x, \rho_j) \cap B(x_j, \eta_e^*(r_j))|}{|B(x_j, \eta_e^*(r_j))|} \\ &\geq AK(\rho_j) \|\mu_j^*\|. \end{aligned}$$

Obviously,  $K * \mu_j^*(x) \leq K(\rho_j) \|\mu_j^*\|$ , whence

$$K * \mu_j^*(x) \leq AK * \mu'_j(x). \quad (3)$$

Now suppose  $x \notin \bigcup B(x_j, \eta_e^*(r_j))$ . Then (3) holds for all  $j$ , so that by (2)

$$K * \mu^*(x) = \sum K * \mu_j^*(x) \leq A \sum K * \mu'_j(x) = AK * \mu'(x) \leq A.$$

Suppose  $x \in B(x_j, \eta_e^*(r_j))$ . Then, by the disjointness of  $\{B(x_j, \eta_e^*(r_j))\}$ , we have  $x \notin \bigcup_{i \neq j} B(x_i, \eta_e^*(r_i))$ . Hence (2) and (3) yield

$$K * \mu^*(x) = K * \mu_j^*(x) + \sum_{i \neq j} K * \mu_i^*(x) \leq K * \mu'_j(x) + A.$$

Therefore

$$\begin{aligned}
\int K * \mu^* d\mu^* &= \int_{\mathbb{R}^N \setminus \cup B(x_j, \eta_e^*(r_j))} K * \mu^* d\mu^* + \sum \int_{B(x_j, \eta_e^*(r_j))} K * \mu^* d\mu^* \\
&\leq A \|\mu^*\| + \sum \int_{B(x_j, \eta_e^*(r_j))} K * \mu_j^* d\mu^* \\
&= A \|\mu^*\| + \sum \int_{B(x_j, \eta_e^*(r_j))} K * \mu_j^* d\mu_j^* \\
&= (A + 1) \sum e_K(E_j),
\end{aligned}$$

where the last equality follows from (1).

Now the proof is easy. Let

$$\tilde{\mu} = \left( \sum e_K(E_j) \right)^{-1/2} \mu^*.$$

We have

$$\begin{aligned}
\int K * \tilde{\mu} d\tilde{\mu} &\leq A, \\
\|\tilde{\mu}\| &= \left( \sum e_K(E_j) \right)^{1/2}.
\end{aligned}$$

Obviously  $\tilde{\mu}$  is concentrated on  $E$ . Hence by definition

$$e_K(E) \geq A \|\tilde{\mu}\|^2 = A \sum e_K(E_j).$$

Thus the required inequality follows. The theorem is proved.  $\square$

The proofs of Theorems 1 and 2 can be carried out in a similar way with the help of Lemma 1 and the following dual definition of  $C_K$  and  $C_{K,p}$  (cf. [8, Theorem 14]). For details we refer to [6].

**Theorem A.** *Let  $E$  be an analytic set. Then*

$$C_K(E) = \sup\{\|\mu\| : \mu \text{ is concentrated on } E, K * \mu \leq 1 \text{ on } \mathbb{R}^N\}.$$

**Theorem B.** *Let  $E$  be an analytic set. Then*

$$C_{K,p}(E) = \sup\{\|\mu\|^p : \mu \text{ is concentrated on } E, \|K * \mu\|_q \leq 1\},$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ .

### 3. Proof of Theorem 4

*Proof of Theorem 4.* Since the proof is same, we shall prove only the first inequality. Take an arbitrary compact subset  $F$  of  $\tilde{E}_{K,p}$ . By the usual covering lemma we can

find  $x_j \in E$  such that

$$\begin{aligned} F &\subset \bigcup B(x_j, 5\eta_p^*(r_j)), \\ \{B(x_j, \eta_p^*(r_j))\} &\text{ is disjoint,} \\ r_j &= \delta_E(x_j). \end{aligned}$$

Let  $E' = \bigcup B(x_j, r_j)$ . By definition this is a subset of  $E$ . We apply Theorems 1 and 2 for  $B(x_j, r_j)$  and  $E'$ . We obtain

$$\sum C_{K,p}(B(x_j, r_j)) \leq AC_{K,p}(E') \leq AC_{K,p}(E).$$

On the other hand we have

$$|F| \leq \sum |B(x_j, 5\eta_p^*(r_j))| = A \sum |B(x_j, \eta_p^*(r_j))|.$$

It is easy to see that

$$|B(0, \eta_p^*(r))| \approx |B(0, \eta_p(r))| = C_{K,p}(B(0, r)).$$

Hence

$$|F| \leq AC_{K,p}(E).$$

Since  $F$  is an arbitrary compact subset of  $\tilde{E}_{K,p}$ , the required inequality follows. The theorem is proved.  $\square$

In Theorem 4 we have considered the enlargement based on balls. We can replace balls by the so-called Nagel-Stein region. Let  $\Omega$  be a set in  $\mathbb{R}_+^{N+1}$  with  $\overline{\Omega} \cap \partial\mathbb{R}_+^{N+1} = \{0\}$ . Put  $\Omega(y) = \{x : (x, y) \in \Omega\}$ . We say that  $\Omega$  satisfies the Nagel-Stein condition (abbreviated to (NS)), if

- (i)  $|\Omega(y)| \leq Ay^N$  with  $A = A(\Omega)$ ;
- (ii) there is  $\alpha > 0$  such that

$$(x_1, y_1) \in \Omega \text{ and } |x - x_1| < \alpha(y - y_1) \implies (x, y) \in \Omega.$$

Obviously, the nontangential cone  $\Gamma = \{(x, y) : |x| < y\}$  satisfies (NS). The section  $\Gamma(y)$  is the open ball with center at 0 and radius  $y$ . So,  $\Omega(y)$  may be regarded as an extension of a ball. For  $E$  we put

$$\tilde{E}_{K,p;\Omega} = \bigcup_{x \in E} (x - \Omega(\eta_p^*(\delta_E(x)))) .$$

This is a generalization of  $\tilde{E}_{K,p}$

**Theorem 5.** *Let  $\Omega$  satisfy (NS). Then*

$$|\tilde{E}_{K,p;\Omega}| \leq AC_{K,p}(E),$$

where  $A > 0$  depends only on  $N, K, p$  and  $\Omega$ .

Theorem 5 can be proved in a similar way with the help of the covering lemma due to Nagel-Stein [10, pp.90–92]. For details we refer to [6].

## 4. Boundary behavior of harmonic functions

In what follows we are interested in the boundary behavior of harmonic functions in  $\mathbb{R}_+^{N+1}$ . Hereafter we let  $1 < p < \infty$ . Following the idea in [2] and [7] we introduce the notion of thinness at the boundary. For a set  $E \subset \mathbb{R}_+^{N+1}$  we put  $E_t = \{(x, y) : 0 < y < t\}$  and  $E^* = \bigcup_{(x,y) \in E} B(x, y)$ . We recall that  $B(x, y)$  is the  $N$ -dimensional ball with center at  $x$  and radius  $y$ , so that the set  $E^*$  is a set on the boundary  $\mathbb{R}^N = \partial\mathbb{R}_+^{N+1}$ . We shall combine the above notation and write

$$E_t^* = \bigcup_{(x,y) \in E, 0 < y < t} B(x, y).$$

**Definition.** Let  $E \subset \mathbb{R}_+^{N+1}$ . We say that  $E$  is  $C_{K,p}$ -thin at  $\partial\mathbb{R}_+^{N+1}$  if

$$\lim_{t \rightarrow 0} C_{K,p}(E_t^*) = 0.$$

**Remark.** If  $E$  is  $C_{K,p}$ -thin at  $\partial\mathbb{R}_+^{N+1}$ , then the essential projection of  $E$

$$\{x : \text{for any } t > 0 \text{ there is a positive number } y < t \text{ such that } (x, y) \in E\}$$

is of  $C_{K,p}$ -capacity 0, and hence of measure 0.

From Theorem 5 we have

**Theorem 6.** Suppose  $\Omega$  satisfies (NS). Let  $\Omega_{K,p} = \{(x, y) : x \in \Omega(\eta_p^*(y))\}$ . If  $E$  is  $C_{K,p}$ -thin at  $\partial\mathbb{R}_+^{N+1}$ , then

$$\left| \bigcap_{t>0} \{x : (x + \Omega_{K,p}) \cap E_t \neq \emptyset\} \right| = 0.$$

In other words, for almost all  $x \in \partial\mathbb{R}_+^{N+1}$ ,  $x + \Omega_{K,p}$  lies eventually outside  $E$ , i.e., there is  $t = t_x > 0$  such that  $E_t \cap (x + \Omega_{K,p}) = \emptyset$ .

*Proof.* It is not so difficult to see that

$$\{x \in \mathbb{R}^N : (x + \Omega_{K,p}) \cap E \neq \emptyset\} \subset \bigcup_{x \in E^*} (x - \Omega_{K,p}(\delta_{E^*}(x))) = \bigcup_{x \in E^*} (x - \Omega(\eta_p^*(\delta_{E^*}(x))))$$

(see [6, Lemma 2]). Hence Theorem 5 yields

$$|\{x \in \mathbb{R}^N : (x + \Omega_{K,p}) \cap E \neq \emptyset\}| \leq AC_{K,p}(E^*).$$

Apply this inequality for  $E_t$  replacing  $E$ . Then the definition of thinness implies that

$$|\{x : (x + \Omega_{K,p}) \cap E_t \neq \emptyset\}| \leq AC_{K,p}(E_t^*) \rightarrow 0 \text{ as } t \rightarrow 0.$$

Thus the theorem follows.  $\square$

**Remark.** It is not so difficult to see that  $\eta_p^*(r)/r \rightarrow \infty$  as  $r \rightarrow 0$  (cf. [1]). Hence  $\Omega_{K,p}$  is a tangential region.

For a function  $f$  on  $\partial\mathbb{R}_+^{N+1}$  we denote by  $PI(f)$  its Poisson integral. In [6] we have proved

**Theorem 7.** *Let  $\Omega \subset \mathbb{R}_+^{N+1}$  and suppose  $\overline{\Omega} \cap \partial\mathbb{R}_+^{N+1} = \{0\}$ . Suppose  $f \in L^p(\mathbb{R}^N)$ . Then there is a set  $E \subset \mathbb{R}_+^{N+1}$  such that  $E$  is  $C_{K,p}$ -thin at  $\partial\mathbb{R}_+^{N+1}$  and that*

$$\lim_{P \rightarrow x, P \in (x+\Omega) \setminus E} PI(K * f)(P) = K * f(x) \quad (4)$$

for  $C_{K,p}$ -a.e.  $x \in \partial\mathbb{R}_+^{N+1}$ , i.e. there is a set  $F \subset \partial\mathbb{R}_+^{N+1}$  such that  $C_{K,p}(F) = 0$  and (4) holds for  $x \in \partial\mathbb{R}_+^{N+1} \setminus F$ .

As a corollary to Theorems 6 and 7 we have the following theorem. This is a generalization of [9, Theorem 2.9].

**Corollary.** *Let  $\Omega \subset \mathbb{R}_+^{N+1}$  and suppose  $\Omega$  satisfies (NS). Suppose  $f \in L^p(\mathbb{R}^N)$ . Then*

$$\lim_{P \rightarrow x, P \in x + \Omega_{K,p}} PI(K * f)(P) = K * f(x)$$

for almost all  $x \in \partial\mathbb{R}_+^{N+1}$ .

**Remark.** In the proof of Theorem 7 we use the nontangential maximal function, which is of type  $(p, p)$  for  $p > 1$  but not of type  $(1, 1)$ . Hence the assumption  $1 < p < \infty$  is necessary. However, we can show similar results even for  $p = 1$  under an additional assumption on the kernel  $k$ . In fact, if

$$K(r) \approx r^{-N} \int_0^r K(t) t^{N-1} dt \text{ for small } r > 0,$$

then the same conclusions of Theorem 7 and Corollary hold. For details we refer to [6].

**Remark.** The approach region  $\Omega_{K,p}$  in Theorem 7 is in some sense most tangential. If one consider less tangential approach regions, then one may obtain boundary limit theorems with smaller boundary exceptional sets (corresponding to  $F$  in Theorem 7) which can be measured by the Hausdorff measure. This problem was considered in [5].



## References

- [1] H. Aikawa, Comparison of  $L^p$ -capacity and Hausdorff measure, *Complex Variables*, 15, (1990), 223–232
- [2] H. Aikawa, Thin sets at the boundary, *Proc. London Math. Soc.* (3), 65, (1992), 357–382
- [3] H. Aikawa, Quasiadditivity of Riesz capacity, *Math. Scand.*, 69, (1991), 15–30
- [4] H. Aikawa, Quasiadditivity of capacity and minimal thinness, *Ann. Acad. Sci. Fenn. Ser. A. I. Mathematica*, 18, (1993), 65–75
- [5] H. Aikawa, Bessel capacity, Hausdorff content and the tangential boundary behavior of harmonic functions, preprint
- [6] H. Aikawa and A. A. Borichev, Quasiadditivity and measure property of capacity and the tangential boundary behavior of harmonic functions, preprint
- [7] A. A. Borichev, Boundary behavior in Dirichlet-type spaces, Uppsala University Department of Mathematics Report, 3, (1994)
- [8] N. G. Meyers, A theory of capacities for potentials of functions in Lebesgue, classes *Math. Scand.*, 26, (1970), 255–292
- [9] A. Nagel, W. Rudin and J. H. Shapiro, Tangential boundary behavior of functions in Dirichlet-type spaces, *Ann. of Math.*, 116, (1982), 331–360
- [10] A. Nagel and E. M. Stein, On certain maximal functions and approach regions, *Adv. in Math.*, 54, (1984), 83–106