

Bose condensates in interaction with excitations - a two-component space-dependent model close to equilibrium.

Leif ARKERYD and Anne NOURI

Mathematical Sciences, 41296 Göteborg, Sweden,
arkeryd@chalmers.se

Aix-Marseille University, CNRS, Centrale Marseille, I2M UMR 7373, 13453 Marseille, France,
anne.nouri@univ-amu.fr

Abstract. The paper considers a model for Bose gases in the so-called 'high-temperature range' below the temperature where Bose-Einstein condensation sets in. The model is of non-linear two-component type, consisting of a kinetic equation with periodic boundary conditions for the distribution function of a gas of excitations interacting with a Bose condensate, which is described by a Gross-Pitaevskii equation. Results on well-posedness and long time behaviour are proved in a Sobolev space setting close to equilibrium.

1 Preliminaries and main results.

1.1 Physics motivations.

The phenomenon of Bose-Einstein condensation occurs when a large number of particles of a Bose gas enter the same lowest accessible quantum state. Predicted by Bose and Einstein in 1924 [4] [6], it was first unambiguously produced in 1995 by E. Cornell and C. Wieman. This paper studies a Bose condensate below the transition temperature T_c for condensation, and in interaction with a non-condensates component. The setting is a two-component space-dependent model well established in physics (see the monograph [10] and its references) of pair-collision interactions involving a gas of thermally excited (quasi-)particles and a condensate. The two-component model consists of a kinetic equation for the distribution function of the gas, and a Gross-Pitaevskii equation (cf [21]) for the condensate. A rather general form of the kinetic equation in the superfluid frame is (cf [22], [26])

$$\partial_t f + (\nabla_p(E(p)) + v_c) \cdot \nabla_x f - \nabla_x(E(p) + v_c \cdot p) \cdot \nabla_p f = C_{22}(f) + C_{12}(f, n_c). \quad (1.1)$$

Here f is the quasi-particle phase space density, n_c (resp. v_c) is the mass density (resp. the velocity) of the condensate, and $E(p)$ denotes the (Bogoliubov) quasi-particle energy. The Nordheim-Uehling-Uhlenbeck term C_{22} for collisions between (quasi-)particles is given by

$$C_{22}(f)(p) = \frac{g^2}{\hbar} \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} B \delta(p + p_* = p' + p'_*) \delta(E(p) + E(p_*) = E(p') + E(p'_*)) \\ \left(f' f'_* (1 + f)(1 + f_*) - f f_* (1 + f')(1 + f'_*) \right) dp_* dp' dp'_*, \quad (1.2)$$

¹2010 Mathematics Subject Classification. 82C10, 82C22, 82C40.

²Key words; low temperature kinetics, Bose condensate, two component model, $0.7 T_c$.

where $g = \frac{4\pi a\hbar^2}{m}$, a is the scattering length of the interaction potential, \hbar the Planck constant, m the atomic mass, B a collision kernel, and

$$f = f(p), \quad f_* = f(p_*), \quad f' = f(p'), \quad f'_* = f(p'_*).$$

The collision term C_{12} for collisions between (quasi-)particles and condensate is

$$C_{12}(f, n_c)(p) = \frac{g^2 n_c}{\hbar} \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} A \delta(p_1 - p_2 - p_3) \delta(E_1 - E_2 - E_3) [\delta(p - p_1) - \delta(p - p_2) - \delta(p - p_3)] ((1 + f_1) f_2 f_3 - f_1 (1 + f_2) (1 + f_3)) dp_1 dp_2 dp_3, \quad (1.3)$$

where A is a collision kernel and

$$f_j = f(p_j), \quad E_j = E(p_j), \quad 1 \leq j \leq 3.$$

The usual Gross-Pitaevskii equation for the wave function ψ (the order parameter) associated with a Bose condensate is

$$i\hbar \partial_t \psi = -\frac{\hbar^2}{2m} \Delta_x \psi + (g|\psi|^2 + U_{ext})\psi,$$

where U_{ext} is an external potential, i.e. a Schrödinger equation complemented by a non-linear term accounting for two-body interactions.

In the present context, the Gross-Pitaevskii equation is further generalized by letting the condensate move in a self-consistent Hartree-Fock mean field $2 \int_{\mathbb{R}^3} f(p) dp$ produced by the thermally excited atoms, together with a dissipative coupling term associated with the collisions. The generalized Gross-Pitaevskii equation derived in e.g. [15], [16], [22] and [10], is of the type

$$i\hbar \partial_t \psi = -\frac{\hbar^2}{2m} \Delta_x \psi + \left(g|\psi|^2 + U_{ext} + 2g \int_{\mathbb{R}^3} f dp + i \frac{g^2}{2\hbar} \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} A \delta(p_1 - p_2 - p_3) \delta(E_1 - E_2 - E_3) ((1 + f_1) f_2 f_3 - f_1 (1 + f_2) (1 + f_3)) dp_1 dp_2 dp_3 \right) \psi. \quad (1.4)$$

The two component problem (1.1), (1.4) is extensively discussed in the physics literature (see [5], [11], [12], [13], [14], [15], [16], [17], [22], [24], [26]). It is proven in [1] that these equations as given in [22], conserve the total energy. That is not so in some of the other settings, in particular not for (1.6)-(1.8) below.

1.2 The model under study.

We restrict to the 'high temperature range', and more particularly consider the temperature range close to $0.7T_c$. As discussed in [5], [15], [16], [26] and more in details in [13], then $|p| \gg \sqrt{2mgn_c}$, the approximation $E(p) = \frac{|p|^2}{2m} + gn_c$ of the quasi-particle energy is commonly used, $A = 1$, the operator C_{22} is negligible, and the mass of the condensate exceeds that of the excitations, i.e. $n_c > \int P(p) dp$. In equilibrium, the right hand side of (1.1) vanishes. Multiplying the collision term by $\log \frac{f}{1+f}$ and integrating in p , it follows that in equilibrium

$$\frac{f_1}{1+f_1} = \frac{f_2}{1+f_2} \frac{f_3}{1+f_3}, \quad \text{when } p_1 = p_2 + p_3, \quad |p_1|^2 = |p_2|^2 + |p_3|^2 + 2mgn_c. \quad (1.5)$$

Equation (1.5) implies that $\frac{f}{1+f}$ is a Maxwellian, hence the phase space density f of the excitations is a Planckian, which is of the type

$$\frac{1}{e^{\alpha(|p|^2 + 2mgn_c) + \beta \cdot p} - 1}, \quad \alpha > 0, \quad \beta \in \mathbb{R}^3, \quad p \in \mathbb{R}^3.$$

In the equilibrium Planckian distribution function, fix the condensate as identically equal to a constant $n_0 > 0$. Set $\alpha = 1$, take the x -component of β as zero, $|\beta| = 2\sqrt{2mgn_0}$ and write the Planckian as $\frac{1}{e^{|p-p_0|^2} - 1}$ with $p_0 = -\frac{\beta}{2}$. Changing variables $p \rightarrow p - p_0$ gives

$$P(p) := \frac{1}{e^{|p|^2} - 1}, \quad p \in \mathbb{R}^3,$$

as equilibrium Planckian distribution function.

The present paper studies the stability of the equilibrium $(P, \sqrt{n_0})$ of the system under small deviations, that respect the conservation laws. Although we are not deriving hydrodynamic limits, we take into account that the system is close to equilibrium and introduce a mean free path ϵ , so that C_{12} becomes $\frac{1}{\epsilon}C_{12}$. The factor g is proportional to the scattering length a , which is smaller than the mean free path ϵ . Take λ of magnitude bounded by $(\frac{g}{\epsilon})^2 (< 1)$. The functions $(f(t, x, p), \psi(t, x))$ are considered in the slab $\Omega = [0, 2\pi]$ in the x -direction with periodic boundary conditions, and taken as

$$(f, \psi) = (P(1 + \lambda R), \sqrt{n_0} + \lambda \Phi).$$

In this paper the external potential U_{ext} is assumed to be a constant that will be further discussed. We could alternatively have left out the external potential in (1.4) but replaced ψ by $e^{itU_{ext}}\psi$ in the proofs. The atomic mass m (resp. the Planck constant \hbar) will be taken as $\frac{1}{2}$ (resp. one) for simplicity. Contrary to the classical Boltzmann operator in velocity space, $f \in L^1(\mathbb{R}^3)$ does not imply $C_{12}(f) \in L^1(\mathbb{R}^3)$. This paper is restricted to distribution functions, cylindrically symmetric in $p = (p_x, p_r) \in \mathbb{R} \times \mathbb{R}^2$. That changes the linear moment conservation Dirac measure in the collision term to $\delta(p_{1x} - p_{2x} - p_{3x})$. Since the collective excitations play no role within the present temperature range, the domain of integration is here taken as the set of $p \in \mathbb{R}^3$ such that $|p|^2 > 2\Lambda^2$ with $\Lambda > 2\sqrt{gn_0}$. Denote by $\tilde{\chi}$ the characteristic function of the set

$$\{(p, p_1, p_2, p_3) \in \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3; |p|^2, |p_1|^2, |p_2|^2, |p_3|^2 > 2\Lambda^2\}.$$

The restriction $|p|^2 > 2\Lambda^2$ will be implicitly assumed below, and $\int dp$ will stand for $\int_{|p|^2 > 2\Lambda^2} dp$. Set

$$\delta_3 = \delta(p - p_1) - \delta(p - p_2) - \delta(p - p_3) \quad \text{and} \quad \delta_0 = \delta(p_{1x} = p_{2x} + p_{3x}, |p_1|^2 = |p_2|^2 + |p_3|^2 + n_0).$$

The system of equations to be satisfied by (f, ψ) is

$$\partial_t f + p_x \partial_x f = g\sqrt{\lambda}n_c \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} \tilde{\chi} \delta_0 \delta_3 (f_2 f_3 - f_1 (1 + f_2 + f_3)) dp_1 dp_2 dp_3, \quad (1.6)$$

$$f(0, x, p) = f_i(x, p), \quad (1.7)$$

and

$$\partial_t \psi - i \partial_x^2 \psi = \left(\frac{\sqrt{\lambda}}{2} \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} \tilde{\chi} \delta_0 (f_2 f_3 - f_1 (1 + f_2 + f_3)) dp_1 dp_2 dp_3 - i \left(n_c + \frac{U_{ext}}{g} + 2 \int f dp \right) \right) g \psi, \quad (1.8)$$

$$\psi(0, x) = \psi_i(x). \quad (1.9)$$

Here, the function n_c is defined by $n_c = n_c(t, x) := |\psi|^2(t, x)$. The approximate energy $|p|^2 + gn_c$ used in (1.5), at this range of temperature is replaced by $|p|^2 + gn_0$ as an approximation of order λ . The total initial mass is

$$2\pi \mathcal{M}_0 := \int_{\Omega} |\psi_i(x)|^2 dx + \int_{\Omega \times \mathbb{R}^3} f_i(x, p) dx dp,$$

which is formally conserved by the equations (1.6) and (1.8). The initial data f_i and ψ_i are taken as

$$f_i := P(1 + \lambda R_i), \quad \psi_i := \sqrt{n_0} + \lambda \Phi_i,$$

for some functions $R_i(x, p)$ and $\Phi_i(x)$ with

$$\int (|\psi_i|^2 - n_0 + \lambda \int_{\mathbb{R}^3} P R_i dp) dx = 0.$$

This is consistent with the asymptotic behavior proven in the paper, i.e. $(f - P, |\psi|^2 - n_0)$ tending to zero when time tends to infinity. It implies that (up to the multiplicative constant $\frac{1}{2\pi}$) the initial (and conserved) total mass equals the mass of (P, n_0) , i.e.

$$\mathcal{M}_0 = \int P(p) dp + n_0. \tag{1.10}$$

The separate masses of condensate and excitation may, however, not be conserved. The constant U_{ext} will be taken as $g(n_0 - 2\mathcal{M}_0)$. For a discussion of general modeling aspects, see also our paper [2] and its references.

1.3 The main mathematical result.

The main results of the paper concern the well-posedness and long time behaviour of the problem (1.6-9).

For an initial perturbation of an equilibrium $(P, \sqrt{n_0})$ of order $(\frac{g}{\epsilon})^2$ and conserving the total mass, the axial momentum and the kinetic energy of the excitations, the problem is well posed and the asymptotic limit when $t \rightarrow +\infty$ of the quasi-particle phase space density and the condensate mass are P resp. $2\pi n_0$. The mass of the excitations together with the mass, the kinetic energy and the internal energy of the condensate converge exponentially to their equilibrium values when $t \rightarrow +\infty$. Let $\| \cdot \|_2$ denote the norm in $L^2([0, 2\pi])$, and set $\| \psi \|_{H^1} := \| \psi \|_2 + \| \partial_x \psi \|_2$, let $\| \cdot \|_{2,2}$ denote the norm in $L^2_{\frac{P}{1+P}}([0, 2\pi] \times \mathbb{R}^3)$, i.e.

$$\| h \|_{2,2} := \left(\int h^2(x, p) \frac{P}{1+P} dp dx \right)^{\frac{1}{2}},$$

and let $L^2_{\frac{1}{P(1+P)}}$ denotes the L^2 -space of functions h with norm $(\int \frac{h^2(x,p)}{P(1+P)} dp dx)^{\frac{1}{2}}$.

The solutions of (1.6-7) will be strong solutions, i.e. such that the collision operator $C_{12}(f, n_c)$ belongs to $C_b(\mathbb{R}^+; L^2_{\nu^{-\frac{1}{2}} \sqrt{\frac{P}{1+P}}}(\mathbb{R}^3; H^1(0, 2\pi)))$, ν being the collision frequency defined in (2.5). The

solutions of (1.8-9) are H^1 -solutions in the following sense. A function $\psi \in C_b(\mathbb{R}^+; H^1_{\text{per}}(0, 2\pi))$ is an H^1 -solution to (1.8-9), if for all $\phi \in C(\mathbb{R}^+; H^1_{\text{per}}(0, 2\pi))$ and all $t > 0$,

$$\begin{aligned} & \int \psi(t, x) \bar{\phi}(t, x) dx - \int \psi_i(x) \bar{\phi}(0, x) dx + i \int_0^t \int \partial_x \psi(s, x) \partial_x \bar{\phi}(s, x) dx ds \\ & = \int_0^t \int \left(\frac{\sqrt{\lambda}}{2} \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3} \tilde{\chi} \delta_0(f_2 f_3 - f_1(1 + f_2 + f_3)) dp_1 dp_2 dp_3 - i(n_c + n_0 - 2\mathcal{M}_0 + 2 \int f dp) \right) g \psi \bar{\phi} dx ds. \end{aligned}$$

Theorem 1.1

There are λ_1 , c_ζ and $\eta_1 > 0$, such that for $\lambda < \lambda_1$ and $(R_i, \Phi_i) \in L^2_{\frac{1+|p|}{1+P}}(\mathbb{R}^3; H^1_{\text{per}}(0, 2\pi)) \times H^1_{\text{per}}(0, 2\pi)$ with

$$\int R_i(x, p) p_x P dx dp = \int R_i(x, p) (|p|^2 + gn_0) P dx dp = 0, \quad (1.11)$$

$$\int (|\psi_i|^2 - n_0 + \lambda \int_{\mathbb{R}^3} P R_i dp) dx = 0, \quad (1.12)$$

and

$$\| \Phi_i \|_{H^1} \leq \eta_1, \quad \| R_i \|_{2,2} + \| \partial_x R_i \|_{2,2} \leq \eta_1, \quad (1.13)$$

there is a unique solution

$$(f, \psi) = (P(1 + \lambda R), \sqrt{n_0} + \lambda \Phi) \in \mathcal{C}_b(\mathbb{R}^+; L^2_{\frac{1}{P(1+P)}}(\mathbb{R}^3; H^1_{\text{per}}(0, 2\pi))) \times \mathcal{C}_b(\mathbb{R}^+; H^1_{\text{per}}(0, 2\pi))$$

to (1.6-9) with $f > 0$. For all $t \in \mathbb{R}^+$, the solution satisfies,

$$\begin{aligned} f &\in L^2_{\frac{(1+|p|)^3}{P(1+P)}}([0, t] \times \mathbb{R}^3; H^1_{\text{per}}(0, 2\pi)), \\ \| R(t, \cdot, \cdot) \|_{2,2} + \| \partial_x R(t, \cdot, \cdot) \|_{2,2} &\leq c_\zeta \eta_1 e^{-\zeta t}, \\ \int (|\partial_x \psi|^2 + \frac{g}{2} (|\psi|^2 - n_0)^2)(t, x) dx &\leq 2\lambda, \end{aligned} \quad (1.14)$$

where $\zeta = c_\zeta \sqrt{\lambda}$.

Moreover, $n_c(t) = \int |\psi(t, x)|^2 dx$ converges exponentially of order ζ to n_0 , when $t \rightarrow +\infty$,

$$\lim_{t \rightarrow +\infty} \int (|\partial_x \psi|^2 + \frac{g}{2} (|\psi|^2 - n_0)^2)(t, x) dx \quad (1.15)$$

exists, and the convergence to its limit is exponential of order ζ .

Whereas non-linear systems of the type (1.6-9) and its generalizations have been much studied in mathematical physics below T_c , there are so far only few papers with their focus mainly on the non-linear mathematical questions. Starting from a similar Gross-Pitaevskii and kinetic frame, two-fluid models are derived in [1]. The space homogeneous initial value problem for this system is treated in [2] for a large data setting. A Milne problem related to the present set-up is studied in [3]. The paper [8] considers a related setting, and has its focus on linearized space homogeneous problems. Validation aspects in the space-homogeneous case are discussed in [23]. There has also been a considerable interest recently (see ee.g. [7], [18] and references therein) in the bosonic Nordheim-Uehling-Uhlenbeck equation as a model above and around T_c for blow-ups and for condensation in space-homogeneous boson gases.

A classical approach to study kinetic equations in a perturbative setting, is to use a spectral inequality (resp. Fourier techniques and the $\| \cdot \|_{T,2,2}$ norm) for controlling the non-hydrodynamic (resp. hydrodynamic) part of a solution. An additional problem here is the coupling with the generalized Gross-Pitaevskii equation. The general approach, together with a Fourier based analysis of the generalized Gross-Pitaevskii equation, provide local in time solutions to the present coupled system. Since the condensate and the normal gas are coupled by the collision interaction, the exponential

decrease of the deviation of the kinetic distribution function from the equilibrium Planckian P , helps to control the long-term evolution of the condensate. This is an important ingredient in the passage from local to global solutions, which leads to exponential decreases of the deviation of the condensate mass from its equilibrium state n_0 , and of the energy (1.15) from its limit value.

Within this frame the kinetic equation (1.6) differs from earlier classical ones. The collision operator in space-homogeneous bosonic Nordheim-Uehling-Uhlenbeck papers has so far been taken isotropic, but is here, due to the space-dependent slab-context, cylindric. Mass density does not belong to the kernel of the present linearized collision operator. The scaling at infinity in its collision frequency is stronger than in the classical case.

The one-dimensional spatial frame induces simplifications of the functional analysis, mainly in the control of the condensate. The \mathbb{T}^d spatial frame, for $d \geq 2$, is an open problem.

The conservation properties of the model (1.6-9), as well as some properties of the collision operator $\frac{C_{12}}{n_c}$ and its linearized operator L around the Planckian P , are discussed in Section 2, including a spectral estimate for L . This is used in Section 3, which is devoted to a priori estimates for some linear equations related to (1.6) and (1.8). They are then employed in the proof of the main theorem in Section 4. The proof starts with a contractive iteration scheme to obtain local solutions. A key point in the global in time analysis is the exponential convergence to equilibrium for f when $t \rightarrow +\infty$. The analysis of ψ differs from the classical Gross-Pitaevskii case. It uses the exponential convergence to equilibrium of f to control the behaviour of the kinetic energy $\int |\partial_x \psi|^2 dx$ and the internal energy $\frac{g}{2} \int |\psi|^4 dx$ of ψ .

2 Some properties of the model and the collision operator.

The model induces total mass conservation as well as axial momentum and kinetic energy conservations for the excitations, as stated in the following lemma.

Lemma 2.1 *It holds that*

$$\frac{d}{dt} \left(\int_{\Omega \times \mathbb{R}^3} f(t, x, p) dx dp + \int_{\Omega} |\psi(t, x)|^2 dx \right) = 0, \quad (2.1)$$

$$\frac{d}{dt} \int_{\Omega \times \mathbb{R}^3} p_x f(t, x, p) dx dp = 0, \quad (2.2)$$

$$\frac{d}{dt} \int_{\Omega \times \mathbb{R}^3} (|p|^2 + gn_0) f(t, x, p) dx dp = 0. \quad (2.3)$$

Proof of Lemma 2.1.

Integrate (1.6) with respect to space and momentum. Add it to (1.8) multiplied by $\bar{\psi}$ (resp. the conjugate of (1.8)) multiplied by ψ integrated with respect to space. One obtains (2.1). Multiplying (1.6) by p_x (resp. $(|p|^2 + gn_0)$) and integrating it w.r.t. space and momentum leads to (2.2) (resp. (2.3)). ■

Since the solutions will remain close to an equilibrium $(P, \sqrt{n_0})$, the linearized operator of C_{12} around P is of interest. For $\gamma := \sqrt{\lambda}$, consider the decomposition

$$f = P(1 + \gamma\tilde{R}), \quad \psi = \sqrt{n_0} + \gamma\tilde{\Phi}.$$

It holds

$$|\psi|^2 = n_c = n_0 + \gamma\sqrt{n_0}(\tilde{\Phi} + \bar{\tilde{\Phi}}) + \gamma^2|\tilde{\Phi}|^2,$$

and the collision term can be written

$$\int \tilde{\chi}\delta_0\delta_3(f_2f_3 - f_1(1 + f_2 + f_3))dp_1dp_2dp_3 = \gamma\left(PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R})\right),$$

where

$$L\tilde{R} := \frac{1}{P} \int \tilde{\chi}\delta(p_{1x} = p_{2x} + p_{3x})\delta(|p_1|^2 = |p_2|^2 + |p_3|^2 + gn_0)(\delta(p - p_1) - \delta(p - p_2) - \delta(p - p_3)) \\ \left[-(1 + P_2 + P_3)P_1\tilde{R}_1 + (P_3 - P_1)P_2\tilde{R}_2 + (P_2 - P_1)P_3\tilde{R}_3 \right] dp_1dp_2dp_3,$$

and

$$2Q(\tilde{R}, \tilde{S}) := \int \tilde{\chi}\delta_0\delta_3\left(P_2P_3(\tilde{R}_2\tilde{S}_3 + \tilde{R}_3\tilde{S}_2) - P_1\tilde{R}_1(P_2\tilde{S}_2 + P_3\tilde{S}_3) - P_1\tilde{S}_1(P_2\tilde{R}_2 + P_3\tilde{R}_3)\right) dp_1dp_2dp_3. \quad (2.4)$$

We recall some properties about L proved in [3].

Lemma 2.2 *L is a self-adjoint operator in $L^2_{\frac{P}{1+P}}$. Within the space of rotationally invariant distribution functions, its kernel is the subspace spanned by $(|p|^2 + gn_0)(1 + P)$ and $p_x(1 + P)$.*

The operator L splits into $K - \nu$, where

$$\nu(p) := \int \tilde{\chi}\delta_0(1 + P_2 + P_3)dp_2dp_3 + 2 \int \tilde{\chi}\delta_0(P_3 - P_1)dp_1dp_3 \quad (2.5)$$

and

$$Kh(p) := \frac{2}{P(p)} \left(\int \tilde{\chi}\delta_0(P_3 - P)P_2h_2dp_2dp_3 + \int \tilde{\chi}\delta_0(1 + P + P_3)P_1h_1dp_1dp_3 \right. \\ \left. + \int \tilde{\chi}\delta_0(P_1 - P)P_3h_3dp_1dp_3 \right). \quad (2.6)$$

Lemma 2.3 *The collision frequency ν satisfies*

$$\nu_0(1 + |p|)^3 \leq \nu(p) \leq \nu_1(1 + |p|)^3, \quad p = (p_x, p_r) \in \mathbb{R} \times \mathbb{R}^+, \quad (2.7)$$

for some positive constants ν_0 and ν_1 . The operator K is compact from $L^2_{\nu\frac{P}{1+P}}$ into $L^2_{\nu^{-1}\frac{P}{1+P}}$.

Denote by (\cdot, \cdot) the scalar product in $L^2_{\frac{P}{1+P}}$, and by \tilde{P} the orthonormal projection on the kernel of L . Set $h_{\parallel} := \tilde{P}h$ and $h_{\perp} := (I - \tilde{P})h$.

Lemma 2.4 L satisfies the spectral inequality,

$$-(Lh, h) \geq c_0(\nu h_\perp, h_\perp), \quad h \in L^2_{(1+|p|)^3 \frac{P}{1+P}}, \quad (2.8)$$

with $c_0 > 0$.

We will also need an estimate for the quadratic collision operator Q .

Lemma 2.5 For cylindrically symmetric functions $(g, h) \in L^2_{\nu \frac{P}{1+P}} \times L^2_{\frac{P}{1+P}}$ (resp. $(g, h) \in L^2_{\frac{P}{1+P}} \times L^2_{\nu \frac{P}{1+P}}$), it holds

$$\left(\int \nu^{-1} \frac{P}{1+P} \left(\frac{Q(g, h)}{P} \right)^2 dp \right)^{\frac{1}{2}} \leq c \left(\int \nu \frac{P}{1+P} g^2(p) dp \int \frac{P}{1+P} h^2(p) dp \right)^{\frac{1}{2}},$$

(resp.

$$\left(\int \nu^{-1} \frac{P}{1+P} \left(\frac{Q(g, h)}{P} \right)^2 dp \right)^{\frac{1}{2}} \leq c \left(\int \nu \frac{P}{1+P} h^2(p) dp \int \frac{P}{1+P} g^2(p) dp \right)^{\frac{1}{2}}).$$

Proof. Considering cylindrically symmetric functions, we will use $g = g(p_x, p_r^2)$, $h = h(p_x, p_r^2)$. The theorem is a consequence of the following estimates for each of the terms of $\frac{Q(h, h)}{P}$. They are of the type

$$Q_1(g, h)(p) := 2 \int k_1(p, p_2) g_2 dp_2 \text{ where } k_1(p, p_2) := P_2 \int \delta(p_x = p_{2x} + p_{3x}, |p|^2 = |p_2|^2 + |p_3|^2 + gn_0) \frac{P_3 h_3}{P} dp_3,$$

or

$$Q_2(g, h)(p) := 2 \int k_2(p, p_2) g_2 dp_2 \text{ where } k_2(p, p_2) := P_2 h \int \delta(p_{1x} = p_{2x} + p_x, |p_1|^2 = |p_2|^2 + |p|^2 + gn_0) dp_1,$$

or

$$Q_3(g, h)(p) := 2 \int k_3(p, p_2) g_2 dp_2 \text{ where } k_3(p, p_2) := P_2 h \int \delta(p_x = p_{2x} + p_{3x}, |p|^2 = |p_2|^2 + |p_3|^2 + gn_0) dp_3,$$

or

$$Q_4(g, h)(p) := 2 \int k_4(p, p_1) g_1 dp_1 \text{ where } k_4(p, p_1) := P_1 h \int \delta(p_{1x} = p_x + p_{3x}, |p_1|^2 = |p|^2 + |p_3|^2 + gn_0) dp_3,$$

or

$$Q_5(g, h)(p) := 2 \int k_5(p, p_3) h_3 dp_3 \text{ where } k_5(p, p_3) := \frac{P_3}{P} \int \delta(p_{1x} = p_x + p_{3x}, |p_1|^2 = |p|^2 + |p_3|^2 + gn_0) P_1 g_1 dp_1.$$

Let $(g, h) \in L^2_{\nu \frac{P}{1+P}} \times L^2_{\frac{P}{1+P}}$. Consider first the term $(\int \nu^{-1} \frac{P}{1+P} \left(\frac{Q_1(g, h)}{P} \right)^2 dp)^{\frac{1}{2}}$. P is uniformly bounded by M from above and below in the domain of integration, so in the estimates below it is

enough to use M instead of P . It holds

$$\begin{aligned}
& \left(\int \nu^{-1} M \left(\int k_1(p, p_2) g_2 dp_2 \right)^2 dp \right)^{\frac{1}{2}} \\
& \leq \int \left(\int \nu^{-1} M k_1^2(p, p_2) dp \right)^{\frac{1}{2}} g_2 dp_2 \\
& \leq c \int M_2 g_2 \left(\int \nu^{-1} M^{-1} (Mh)^2 (p_x - p_{2x}, |p|^2 - |p_2|^2 - gn_0 - |p_x - p_{2x}|^2) dp \right)^{\frac{1}{2}} dp_2 \\
& \leq c \int M_2 g_2 \left(\int \nu^{-1} (\sqrt{|p_2|^2 + |p_3|^2 + gn_0}) M_2^{-1} M_3^{-1} (M_3 h_3)^2 dp_3 \right)^{\frac{1}{2}} dp_2 \\
& \leq c \left(\int \frac{1}{(1 + |p_2|)^{\frac{3}{2}}} M_2^{\frac{1}{2}} g_2 dp_2 \right) \left(\int M_3 h_3^2 dp_3 \right)^{\frac{1}{2}} \\
& \leq c \left(\int \nu_2 M_2 g_2^2 dp_2 \right)^{\frac{1}{2}} \left(\int M_3 h_3^2 dp_3 \right)^{\frac{1}{2}},
\end{aligned}$$

by the Cauchy-Schwartz inequality. For the $(\int \nu^{-1} \frac{P}{1+P} (\frac{Q_2(g,h)}{P})^2 dp)^{\frac{1}{2}}$ term,

$$\begin{aligned}
\left(\int \nu^{-1} M \left(\int k_2(p, p_2) g_2 dp_2 \right)^2 dp \right)^{\frac{1}{2}} & \leq \int \left(\int \nu^{-1} M k_2^2(p, p_2) dp \right)^{\frac{1}{2}} g_2 dp_2 \\
& \leq c \left(\int M_2 g_2 dp_2 \right) \left(\int \nu^{-1} M h^2 dp \right)^{\frac{1}{2}} \\
& \leq c \left(\int \nu_2 M_2 g_2^2 dp_2 \right)^{\frac{1}{2}} \left(\int M h^2 dp \right)^{\frac{1}{2}}.
\end{aligned}$$

The $(\int \nu^{-1} \frac{P}{1+P} (\frac{Q_i(g,h)}{P})^2 dp)^{\frac{1}{2}}$ terms can be handled similarly. Finally,

$$\begin{aligned}
\left(\int \nu^{-1} M \left(\int \tilde{k}_5(p, p_3) h_3 dp_3 \right)^2 dp \right)^{\frac{1}{2}} & \leq \int \left(\int \nu^{-1} M \tilde{k}_5^2(p, p_3) dp \right)^{\frac{1}{2}} h_3 dp_3 \\
& \leq c \int M_3^{\frac{3}{2}} h_3 dp_3 \left(\int M_1 g_1^2 dp_1 \right)^{\frac{1}{2}} \\
& \leq c \left(\int M_3 h_3^2 dp_3 \right)^{\frac{1}{2}} \left(\int M_1 g_1^2 dp_1 \right)^{\frac{1}{2}}.
\end{aligned}$$

This completes the proof of the lemma. ■

Lemma 2.6

There is a constant $c > 0$ such that for any cylindrically symmetric function $f \in L^2(\mathbb{R}^3)$,

$$\left| \int P(Lf)(p) dp \right| \leq c \left(\int \frac{P}{1+P} f_{\perp}^2(p) dp \right)^{\frac{1}{2}}.$$

Proof. Using the Cauchy-Schwartz inequality,

$$\begin{aligned}
\left| \int P(Lf)(p) dp \right| & = \left| \int (\sqrt{P}\nu) \left(\frac{\sqrt{P}}{\nu} Lf \right) dp \right| \\
& \leq c \left(\int P f_{\perp}^2(p) dp \right)^{\frac{1}{2}} \\
& \leq c \left(\int \frac{P}{1+P} f_{\perp}^2(p) dp \right)^{\frac{1}{2}}.
\end{aligned}$$

3 Rest term estimates.

Consider the decomposition

$$f = P(1 + \gamma\tilde{R}), \quad \psi = \sqrt{n_0} + \gamma\tilde{\Phi}.$$

The equations (1.8)-(1.9) written for $\tilde{\Phi}$ with periodic boundary conditions when \tilde{R} is given, are

$$\partial_t \tilde{\Phi} - i\partial_x^2 \tilde{\Phi} = S_1 \tilde{\Phi} + S_2 \bar{\tilde{\Phi}} + U, \quad \tilde{\Phi}(0, \cdot) = \tilde{\Phi}_i. \quad (3.1)$$

Here S_1 and S_2 are the coefficients of the linear terms in $\tilde{\Phi}$ resp. $\bar{\tilde{\Phi}}$, and U contains the inhomogeneous terms and the non-linear terms in $\tilde{\Phi}$, $\bar{\tilde{\Phi}}$. In the following lemmas the dependence of U on $\tilde{\Phi}$ is not taken into account.

Lemma 3.1

Let $\tilde{\Phi}_i$ (resp. S_1, S_2, U) be a given function in $H_{per}^1(0, 2\pi)$ (resp. $L^\infty(\mathbb{R}^+; H_{per}^1(0, 2\pi))$).

There is a unique solution $\tilde{\Phi}$ to (3.1) in $\mathcal{C}(\mathbb{R}^+; H_{per}^1(0, 2\pi))$. Moreover,

$$\|\tilde{\Phi}(t, \cdot)\|_{H^1}^2 \leq (2\|\tilde{\Phi}_i\|_{H^1}^2 + 6t \int_0^t \|U(s, \cdot)\|_{H^1}^2 ds) e^{6t \int_0^t (\|S_1(r, \cdot)\|_{H^1}^2 + \|S_2(r, \cdot)\|_{H^1}^2) dr}, \quad t > 0. \quad (3.2)$$

Proof of Lemma 3.1

Consider first the equations

$$\partial_t \tilde{\Phi} - i\partial_x^2 \tilde{\Phi} = W, \quad \tilde{\Phi}(0, \cdot) = \tilde{\Phi}_i, \quad (3.3)$$

for a given $W \in L^\infty(\mathbb{R}^+; H_{per}^1(0, 2\pi))$. Writing W and $\tilde{\Phi}$ in Fourier series, gives

$$\hat{\Phi}'_n(t) + in^2 \hat{\Phi}_n = \hat{W}_n,$$

and so

$$\hat{\Phi}_n(t) = \hat{\Phi}_n(0)e^{-in^2 t} + \int_0^t \hat{W}_n(s)e^{in^2(s-t)} ds. \quad (3.4)$$

Hence,

$$\begin{aligned} |\hat{\Phi}_n(t)| &\leq |\hat{\Phi}_n(0)| + \int_0^t |\hat{W}_n(s)| ds, \\ \sum_{n \in \mathbb{N}} (1+n^2) |\hat{\Phi}_n(t)|^2 &\leq 2 \sum_{n \in \mathbb{N}} [(1+n^2) |\hat{\Phi}_n(0)|^2 + t \int_0^t (1+n^2) |\hat{W}_n(s)|^2 ds]. \end{aligned}$$

And so the function $\tilde{\Phi}$ defined by (3.4) belongs to $L_{loc}^\infty(\mathbb{R}^+; H_{per}^1(0, 2\pi))$. Moreover,

$$\|\tilde{\Phi}(t, \cdot)\|_{H^1}^2 \leq 2 \left(\|\tilde{\Phi}_i\|_{H^1}^2 + t \int_0^t \|W(s, \cdot)\|_{H^1}^2 ds \right).$$

We conclude that given $\tilde{\Phi}_i \in H_{per}^1(0, 2\pi)$ and $W \in L^\infty(\mathbb{R}^+; H_{per}^1(0, 2\pi))$, there exists a unique solution $\tilde{\Phi} \in L_{loc}^\infty(\mathbb{R}^+; H_{per}^1(0, 2\pi))$ to (3.3). It also follows from (3.4) that the solution is a continuous function of $t \in \mathbb{R}^+$ into $H_{per}^1(0, 2\pi)$. For $W = W(\tilde{\Phi}) := S_1\tilde{\Phi} + S_2\tilde{\Phi} + U$ it holds,

$$\begin{aligned} \|W(s, \cdot)\|_{H^1}^2 &\leq 3 \|S_1\tilde{\Phi}(s, \cdot)\|_{H^1}^2 + 3 \|S_2\tilde{\Phi}(s, \cdot)\|_{H^1}^2 + 3 \|U(s, \cdot)\|_{H^1}^2 \\ &\leq 3 \|S_1(s, \cdot)\|_{H^1}^2 \|\tilde{\Phi}(s, \cdot)\|_{H^1}^2 + 3 \|S_2(s, \cdot)\|_{H^1}^2 \|\tilde{\Phi}(s, \cdot)\|_{H^1}^2 + 3 \|U(s, \cdot)\|_{H^1}^2. \end{aligned}$$

With $\tilde{\Phi}_0 = 0$, an iterative sequence of solutions $\tilde{\Phi}_j$ of (3.3) for $j \geq 1$ with the right hand side $W(\tilde{\Phi}_{j-1})$, gives

$$\|\tilde{\Phi}_j(t, \cdot)\|_{H^1}^2 \leq 2 \|\tilde{\Phi}_i\|_{H^1}^2 + 6t \int_0^t \left((\|S_1(s, \cdot)\|_{H^1}^2 + \|S_2(s, \cdot)\|_{H^1}^2) \|\tilde{\Phi}_{j-1}(s, \cdot)\|_{H^1}^2 + \|U(s, \cdot)\|_{H^1}^2 \right) ds, \quad (3.5)$$

and with $\delta\tilde{\Phi}_j = \tilde{\Phi}_j - \tilde{\Phi}_{j-1}$,

$$\|\delta\tilde{\Phi}_j(t, \cdot)\|_{H^1}^2 \leq 6t \int_0^t \left((\|S_1(s, \cdot)\|_{H^1}^2 + \|S_2(s, \cdot)\|_{H^1}^2) \|\delta\tilde{\Phi}_{j-1}(s, \cdot)\|_{H^1}^2 \right) ds.$$

It follows that the sequence converges on some interval $t \in [0, T]$, and that (3.4) and (3.5) hold for the limit $\tilde{\Phi}$, a unique solution of (3.1). By an iteration of the argument the existence and the continuity of $\tilde{\Phi}$ hold for $t > 0$. Using Gronwall on (3.5) for $\tilde{\Phi}$ gives (3.2). \blacksquare

The rest of this section prepares for the control of the excitation distribution function f around the equilibrium P . With

$$f = P(1 + \gamma\tilde{R}), \quad f_i = P(1 + \gamma\tilde{R}_i),$$

the equations (1.6)-(1.7) written for \tilde{R} , are

$$\partial_t \tilde{R} + p_x \partial_x \tilde{R} = g\gamma(n_0 L\tilde{R} + \gamma(L_1\tilde{R} + Q_1(\tilde{R}, \tilde{R}))), \quad \tilde{R}(0, \cdot, \cdot) = \tilde{R}_i, \quad (3.6)$$

where L_1 (resp. Q_1) is a linear (resp. quadratic) operator.

The following norms are used. For $1 \leq q \leq \infty$,

$$\begin{aligned} \|f\|_{2,q} &= \left(\int_{\mathbb{R}^3} \frac{P}{1+P} \left(\int_{[0,2\pi]} |f(x,p)|^q dx \right)^{\frac{2}{q}} dp \right)^{\frac{1}{2}}, \\ \|f\|_{2,H^1} &= \left(\int_{[0,2\pi] \times \mathbb{R}^3} \frac{P}{1+P} (|f(x,p)|^2 + |\partial_x f(x,p)|^2) dx dp \right)^{\frac{1}{2}}, \\ \|f\|_{T,2,2} &= \left(\int_{[0,T] \times [0,2\pi] \times \mathbb{R}^3} \frac{P}{1+P} |f(t,x,p)|^2 dt dx dp \right)^{\frac{1}{2}}. \end{aligned}$$

To study (3.6), some a priori estimates will be needed for the linear problem

$$\partial_t h + p_x \partial_x h = g\gamma(n_0 Lh + \gamma G), \quad h(0, \cdot, \cdot) = h_0, \quad (3.7)$$

periodic in x with period 2π . Assume $G \in L^\infty([0, T]; L^2_{(1+|p|)^{-3} \frac{P}{1+P}}(\mathbb{R}^3; H^1_{\text{per}}(0, 2\pi)))$ for $T > 0$. The function $\partial_x h$ is at least formally a solution to

$$\partial_t(\partial_x h) + p_x \partial_x(\partial_x h) = g\gamma(n_0 L(\partial_x h) + \gamma \partial_x G), \quad \partial_x h(0, \cdot, \cdot) = \partial_x h_0, \quad (3.8)$$

periodic in x with period 2π . For existence of solutions to problems of the type (3.7), see [19] or alternatively, consider the Fourier transform in x of (3.7) and argue as in the proof of Lemma 3.1. The solutions are unique and continuous as functions of t into $L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H^1_{\text{per}}(0, 2\pi))$. Multiply the equation by $h \frac{P}{1+P}$, integrate on $[0, T] \times [0, 2\pi] \times \mathbb{R}^3$, and use (2.8) to get

Lemma 3.2 For any $\eta > 0$,

$$\|h(T, \cdot)\|_{2,2}^2 + \gamma \|\nu^{\frac{1}{2}} h_\perp\|_{T,2,2}^2 \leq c(\|h_0\|_{2,2}^2 + \gamma^3 \|\nu^{-\frac{1}{2}} G_\perp\|_{T,2,2}^2 + \gamma^2 \eta \|G_\parallel\|_{T,2,2}^2 + \frac{\gamma^2}{\eta} \|h_\parallel\|_{T,2,2}^2).$$

Lemma 3.3 Assume $\gamma \leq 1$,

$$\int_0^{2\pi} \int P(|p|^2 + gn_0) G(t, x, p) dp dx = \int_0^{2\pi} \int P p_x G(t, x, p) dp dx = 0, \quad t \in (0, T), \quad (3.9)$$

and

$$\int_0^{2\pi} \int P(|p|^2 + gn_0) h_0(x, p) dp dx = \int_0^{2\pi} \int P p_x h_0(x, p) dp dx = 0. \quad (3.10)$$

Then

$$\|h_\parallel\|_{T,2,2}^2 \leq c\left(\gamma^{-1} \|h_0\|_{2,2}^2 + \gamma^2 (\|\nu^{-\frac{1}{2}} G_\perp\|_{T,2,2}^2 + \|G_\parallel\|_{T,2,2}^2)\right).$$

Proof of Lemma 3.3

Let $\{\chi_x = c_1 p_x(1+P), \chi_2 = c_2(|p|^2 + gn_0)(1+P)\}$ be an orthonormal basis for the kernel of L . Consider the Fourier series in x of (3.7),

$$\partial_t h_k - ik p_x h_k = c_k + g\gamma^2 G_k, \quad (3.11)$$

where c_k (resp. G_k) is the k -th Fourier coefficient of $g\gamma n_0 Lh$ (resp. G). Set $h_{k2} = (h_k, \chi_2)$ and $h_{kx} = (h_k, \chi_x)$. Multiply (3.11) by $\frac{P}{1+P} \chi_2$ (resp. $\frac{P}{1+P} \chi_x$) and integrate in p ,

$$\partial_t h_{k2} - ik \kappa (h_{kx} + \kappa^{-1} h_{kp_x \chi_2}) = g\gamma^2 G_{k2}, \quad (3.12)$$

$$\partial_t h_{kx} - ik \kappa (h_{k2} + \kappa^{-1} h_{kp_x \chi_x}) = g\gamma^2 G_{kx}. \quad (3.13)$$

Here $h_{p_x \chi_x}$ and $h_{p_x \chi_2}$, denote non-hydrodynamic moments of h , and $\kappa := \int \frac{P}{1+P} p_x \chi_2 \chi_x dp$.

By adding and subtracting (3.12) resp. (3.13) with their conjugates, and dividing by 2 or $2i$, we get equations for the real and imaginary parts,

$$\partial_t \mathcal{R} h_{k2} - ik \kappa (\mathcal{I} h_{kx} + \kappa^{-1} \mathcal{I} h_{kp_x \chi_2}) = g\gamma^2 \mathcal{R} G_{k2},$$

$$\partial_t \mathcal{I} h_{k2} - ik \kappa (\mathcal{R} h_{kx} + \kappa^{-1} \mathcal{R} h_{kp_x \chi_2}) = g\gamma^2 \mathcal{I} G_{k2},$$

$$\partial_t \mathcal{R} h_{kx} - ik \kappa (\mathcal{I} h_{k2} + \kappa^{-1} \mathcal{I} h_{kp_x \chi_x}) = g\gamma^2 \mathcal{R} G_{kx},$$

$$\partial_t \mathcal{I} h_{kx} - ik \kappa (\mathcal{R} h_{k2} + \kappa^{-1} \mathcal{R} h_{kp_x \chi_x}) = g\gamma^2 \mathcal{I} G_{kx}.$$

Multiply the four equations by respectively $\mathcal{I}h_{kx}$, $\mathcal{R}h_{kx}$, $\mathcal{I}h_{k2}$, $\mathcal{R}h_{k2}$, and sum. This leads to the ∂_t -derivative

$$\partial_t(\mathcal{R}h_{k2}\mathcal{I}h_{kx}) + \partial_t(\mathcal{R}h_{kx}\mathcal{I}h_{k2}),$$

which after integrating with respect to t on $[0, T]$, evaluates at T and at zero. Considering the full expressions for $k \neq 0$ and dividing by $k\kappa$, gives

$$\begin{aligned} \int_0^T (|h_{kx}|^2 + |h_{k2}|^2) dt &\leq c \left(|(\mathcal{R}h_{k2}\mathcal{I}h_{kx})(T)| + |(\mathcal{R}h_{kx}\mathcal{I}h_{k2})(T)| + |(\mathcal{R}h_{k2}\mathcal{I}h_{kx})(0)| + |(\mathcal{R}h_{kx}\mathcal{I}h_{k2})(0)| \right. \\ &\quad \left. + \int_0^T (|h_{kx}| \cdot |h_{kp_x\chi_2}| + |h_{k2}| \cdot |h_{kp_x\chi_x}| + g\gamma^2(|h_{kx}| \cdot |G_{k2}| + |h_{k2}| \cdot |G_{kx}|)) dt \right). \end{aligned}$$

Finally for $k = 0$, $\partial_t h_{02} = g\gamma^2 G_{02}$, $\partial_t h_{0x} = g\gamma^2 G_{0x}$, so that by (3.9)-(3.10),

$$|h_{02}|^2 + |h_{0x}|^2 = 0, \quad t \in (0, T).$$

We conclude that

$$\begin{aligned} \int_0^T \|h_{\parallel}(t, \cdot)\|_{2,2}^2 dt &\leq c \left(\|h_{\parallel}(T, \cdot)\|_{2,2}^2 + \|h_{0\parallel}\|_{2,2}^2 + \eta_1 \int_0^T \|h_{\parallel}(\tau, \cdot)\|_{2,2}^2 dt \right. \\ &\quad \left. + \frac{1}{\eta_1} \int_0^T \|h_{\perp}(t, \cdot)\|_{2,2}^2 d\tau + g\gamma^2 \int_0^T \|G_{\parallel}(t, \cdot)\|_{2,2}^2 dt \right). \end{aligned}$$

Using Lemma 3.2 leads to the statement of the lemma. \blacksquare

Moreover, it follows from the expression of h_{\parallel} in the basis $\{(1+P)p_x, (1+P)(|p|^2 + gn_0)\}$ of the kernel of L , that there is a constant c such that for any $h \in L^2_{\frac{P}{1+P}}$,

$$\|\nu^{\frac{1}{2}}h_{\parallel}\|_{2,2} \leq c \|h_{\parallel}\|_{2,2}. \quad (3.14)$$

Consequently, lemmas 3.2-3.3 give that

Lemma 3.4 *Under the assumptions (3.9) and (3.10), there is a constant $\tilde{\beta}_1 > 0$ such that the x -periodic solution h to the initial value problem (3.7) satisfies*

$$\|h(t, \cdot)\|_{2,2} + \sqrt{\gamma} \|\nu^{\frac{1}{2}}h\|_{T,2,2} \leq \tilde{\beta}_1 \left(\|h_0\|_{2,2} + \gamma^{\frac{3}{2}} (\|\nu^{-\frac{1}{2}}G_{\perp}\|_{T,2,2} + \|G_{\parallel}\|_{T,2,2}) \right), \quad t \in [0, T].$$

Due to the periodic setting, the integrals $\int_0^{2\pi} \partial_x G dx = \int_0^{2\pi} \partial_x h_0 dx = 0$, and so assumptions (3.9)-(3.10) are satisfied by $\partial_x h$, solution to (3.8). Similarly to Lemma 3.4, it holds

Lemma 3.5 *Under the assumptions (3.9) and (3.10) and analogous ones for $\partial_x h_0$ and $\partial_x G$, there is a constant $\tilde{\beta}_2 > 0$ such that the x -periodic solution h to the initial value problem (3.7) satisfies*

$$\|\partial_x h(t, \cdot)\|_{2,2} + \sqrt{\gamma} \|\nu^{\frac{1}{2}}\partial_x h\|_{T,2,2} \leq \tilde{\beta}_2 \left(\|\partial_x h_0\|_{2,2} + \gamma^{\frac{3}{2}} (\|\nu^{-\frac{1}{2}}\partial_x G_{\perp}\|_{T,2,2} + \|\partial_x G_{\parallel}\|_{T,2,2}) \right), \quad t \in [0, T].$$

In the rest of the paper, the notation

$$\beta_1 = \max\{\tilde{\beta}_1, \tilde{\beta}_2\} \quad (3.15)$$

will be used.

4 Proof of the main theorem.

We shall now use the a priori estimates for the linearized equations in Section 3 to construct solutions to the two component model (1.6-9), and begin with local in time solutions $\tilde{\Phi}$ and \tilde{R} to the equations

$$\partial_t \tilde{\Phi} - i\partial_x^2 \tilde{\Phi} = S_1(\tilde{R})\tilde{\Phi} - ign_0\tilde{\Phi} + U(\tilde{\Phi}, \tilde{R}), \quad \tilde{\Phi}(0, \cdot) = \tilde{\Phi}_i, \quad (4.1)$$

$$\partial_t \tilde{R} + p_x \partial_x \tilde{R} = g\gamma \left(n_0 L\tilde{R} + \gamma(L_{1,\tilde{\Phi}}\tilde{R} + Q_{1,\tilde{\Phi}}(\tilde{R}, \tilde{R})) \right), \quad \tilde{R}(0, \cdot, \cdot) = \tilde{R}_i, \quad (4.2)$$

obtained from (1.8), and (1.6). Here,

$$S_1(\tilde{R}) = -ign_0 - \frac{g}{2}\gamma^2 \int PL\tilde{R}dp - 2ig\gamma \int P\tilde{R}dp - \frac{g}{2}\gamma^3 \int Q(\tilde{R}, \tilde{R})dp,$$

$$\begin{aligned} U(\tilde{\Phi}, \tilde{R}) &= -\frac{\sqrt{n_0}}{2}g\gamma \int PL\tilde{R}dp - 2ig\sqrt{n_0} \int P\tilde{R}dp - i\sqrt{n_0}g\gamma(2|\tilde{\Phi}|^2 + \tilde{\Phi}^2) \\ &\quad - \frac{\sqrt{n_0}}{2}g\gamma^2 \int Q(\tilde{R}, \tilde{R})dp - ig\gamma^2 \tilde{\Phi}|\tilde{\Phi}|^2, \end{aligned}$$

$$L_{1,\tilde{\Phi}}\tilde{R} := \left(\sqrt{n_0}(\tilde{\Phi} + \bar{\tilde{\Phi}}) + \gamma|\tilde{\Phi}|^2 \right) Ls = \frac{|\psi|^2 - n_0}{\gamma} L\tilde{R} =: L_{1,\psi}\tilde{R},$$

$$Q_{1,\tilde{\Phi}}(\tilde{R}, \tilde{R}) = (n_0 + \gamma\sqrt{n_0}(\tilde{\Phi} + \bar{\tilde{\Phi}}) + \gamma^2|\tilde{\Phi}|^2) \frac{Q(\tilde{R}, \tilde{R})}{P}.$$

Denote by

$$\begin{aligned} \|\tilde{\Phi}\|_{\infty T} &:= \sup_{t \in [0, T]} \|\tilde{\Phi}(t, \cdot)\|_{H^1(0, 2\pi)}, \\ \|\tilde{R}\|_{q, 2, H^1} &:= \left(\int_0^T \|\tilde{R}(t, \cdot, \cdot)\|_{L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H^1(0, 2\pi))}^q dt \right)^{\frac{1}{q}}, \quad q \in [2, +\infty], \\ \|\tilde{R}(t)\|_{2, H^1} &:= \|\tilde{R}(t, \cdot, \cdot)\|_{L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H^1(0, 2\pi))}. \end{aligned}$$

Some constants to be used later, are introduced next. By rescaling it is enough to consider $n_0 = 1$. The constant β_1 was defined in (3.15). Set

$$\beta_2 = \left(\int P(1+P)dp \right)^{\frac{1}{2}}, \quad \beta'_2 = \left(\int P(1+P)\nu dp \right)^{\frac{1}{2}}. \quad (4.3)$$

Denote by β_3 (resp. β_4) the norm of the injection from $H^1(0, 2\pi)$ into $L^4(0, 2\pi)$ (resp. $L^\infty(0, 2\pi)$). By the Cauchy-Schwartz inequality and Lemmas 2.2 to 2.6, there are constants $(\beta_i)_{5 \leq i \leq 8}$ such that for any function $g \in L^2_{\frac{P}{1+P}}$ (resp. $g \in L^2_{\nu \frac{P}{1+P}}$, $h \in L^2_{\frac{P}{1+P}}$),

$$\left| \int PLgdp \right| \leq \beta_5 \left(\int g^2 \frac{P}{1+P} dp \right)^{\frac{1}{2}}, \quad (4.4)$$

(resp.

$$\left(\int \nu^{-1} \frac{P}{1+P} (Lg)^2 dp \right)^{\frac{1}{2}} \leq \beta_6 \left(\int \nu \frac{P}{1+P} (g_\perp)^2 dp \right)^{\frac{1}{2}}, \quad (4.5)$$

$$|\int Q(g, g)dp| \leq \beta_7 \left(\int g^2 \frac{P}{1+P} dp \right)^{\frac{1}{2}} \left(\int \nu(p) g^2 \frac{P}{1+P} dp \right)^{\frac{1}{2}}, \quad (4.6)$$

$$\left(\int \nu^{-1} \frac{P}{1+P} \left(\frac{Q(g, h)}{P} \right)^2 dp \right)^{\frac{1}{2}} \leq \beta_8 \left(\int \nu \frac{P}{1+P} g^2(p) dp \int \frac{P}{1+P} h^2(p) dp \right)^{\frac{1}{2}}. \quad (4.7)$$

A control of $\| \nu^{-\frac{1}{2}} \frac{Q(g, h)}{P} \|_{2,2,H^1}$ will also be needed and is given in the following lemma.

Lemma 4.1

For cylindrically symmetric functions g and h in $L^2(0, T; L^2_{\nu \frac{P}{1+P}}(\mathbb{R}^3; H^1(0, 2\pi)))$,

$$\| \nu^{-\frac{1}{2}} \frac{Q(g, h)}{P} \|_{2,2,H^1} \leq 2\beta_4 \beta_8 (\| g \|_{\infty,2,H^1} \| \nu^{\frac{1}{2}} h \|_{2,2,H^1} + \| h \|_{\infty,2,H^1} \| \nu^{\frac{1}{2}} g \|_{2,2,H^1}). \quad (4.8)$$

Proof of Lemma 4.1

By definition of $\| \cdot \|_{2,2,H^1}$,

$$\begin{aligned} & \| \nu^{-\frac{1}{2}} \frac{Q(g, h)}{P} \|_{2,2,H^1}^2 \\ & \leq \int \nu^{-1} \left(\frac{Q(g, h)}{P} \right)^2 \frac{P}{1+P} dp dx dt + 2 \int \nu^{-1} \left(\frac{Q(g, \partial_x h)}{P} \right)^2 \frac{P}{1+P} dp dx dt + 2 \int \nu^{-1} \left(\frac{Q(\partial_x g, h)}{P} \right)^2 \frac{P}{1+P} dp dx dt \\ & \leq 2\beta_8^2 \left(\int \left(\int g^2 \frac{P}{1+P} dp \right) \left(\int \nu(h^2 + (\partial_x h)^2) \frac{P}{1+P} dp \right) dx dt \right. \\ & \quad \left. + \int \left(\int h^2 \frac{P}{1+P} dp \right) \left(\int \nu(g^2 + (\partial_x g)^2) \frac{P}{1+P} dp \right) dx dt \right). \end{aligned}$$

Moreover,

$$\begin{aligned} & \int \left(\int g^2 \frac{P}{1+P} dp \right) \left(\int \nu(h^2 + (\partial_x h)^2) \frac{P}{1+P} dp \right) dx dt \\ & \leq \left(\sup_{(t,x) \in [0,T] \times [0,2\pi]} \int g^2(t, x, p) \frac{P}{1+P} dp \right) \| \nu^{\frac{1}{2}} h \|_{2,2,H^1}^2 \\ & \leq \left(\sup_{t \in [0,T]} \int \left(\sup_{x \in [0,2\pi]} |g(t, x, p)| \right)^2 \frac{P}{1+P} dp \right) \| \nu^{\frac{1}{2}} h \|_{2,2,H^1}^2 \\ & \leq \beta_4^2 \left(\sup_{t \in [0,T]} \int (g^2 + (\partial_x g)^2)(t, x, p) \frac{P}{1+P} dx dp \right) \| \nu^{\frac{1}{2}} h \|_{2,2,H^1}^2 \\ & = \beta_4^2 \| g \|_{\infty,2,H^1}^2 \| \nu^{\frac{1}{2}} h \|_{2,2,H^1}^2. \end{aligned}$$

Applying an analogous inequality to $\int \left(\int h^2 \frac{P}{1+P} dp \right) \left(\int \nu(g^2 + (\partial_x g)^2) \frac{P}{1+P} dp \right)$ leads to (4.8).

Proposition 4.1

Let $(\tilde{\Phi}_i, \tilde{R}_i) \in H^1_{\text{per}}(0, 2\pi) \times L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H^1_{\text{per}}(0, 2\pi))$, satisfy

$$\int \tilde{R}_i(x, p) p_x P dx dp = \int \tilde{R}_i(x, p) (|p|^2 + gn_0) P dx dp = 0, \quad (4.9)$$

and

$$\int (|\psi_i|^2 - 1 + \gamma \int_{\mathbb{R}^3} P \tilde{R}_i dp) dx = 0. \quad (4.10)$$

There are $T_0, \gamma_0 \in]0, 1]$ and $c_1 \geq 1$, such that for $\gamma \in]0, \gamma_0]$, there is a unique solution

$$(\tilde{\Phi}, \tilde{R}) \in \mathcal{C}_b([0, T_0]; H_{\text{per}}^1(0, 2\pi)) \times \mathcal{C}_b([0, T_0]; L_{\frac{P}{1+P}}^2(\mathbb{R}^3; H_{\text{per}}^1(0, 2\pi)))$$

to (4.1-2), for which

$$\tilde{R} \in L_{\frac{(1+|p|)^3}{P(1+P)}}^2([0, T_0] \times \mathbb{R}^3; H_{\text{per}}^1(0, 2\pi)),$$

and

$$\|\tilde{\Phi}\|_{\infty T} + \|\tilde{R}\|_{\infty, 2, H^1} + \sqrt{\gamma} \|\nu^{\frac{1}{2}} \tilde{R}\|_{2, 2, H^1} \leq c_1 (\|\tilde{\Phi}_i\|_{H^1} + \|\tilde{R}_i\|_{2, H^1}),$$

where

$$c_1 = \max\{1, 2(\beta_1 + 2e^{12\pi g^2})\}. \quad (4.11)$$

Given $\eta_0 > 0$, when $\|\tilde{\Phi}_i\|_{H^1} + \|\tilde{R}_i\|_{2, H^1}$ is bounded by η_0 , then T_0 and γ_0 can be taken to depend only on η_0 .

Proof of Proposition 4.1.

Denote by $\eta_0 > 0$ a bound of $\|\tilde{\Phi}_i\|_{H^1} + \|\tilde{R}_i\|_{2, H^1}$. Let \bar{c}_1 be the maximum of

$$30g(1 + 2\eta_0 e^{12\pi g^2})(\beta_2 + \beta_5 + \beta_1\beta_4\beta_7\eta_0 + 2\eta_0 e^{12\pi g^2})e^{60g^2(1 + \beta_1^2\eta_0^2(\beta_2^2 + \beta_5^2 + (\beta_1\beta_4\beta_7\eta_0)^2))}, \quad (4.12)$$

$$1 + 4\beta_1^2\beta_4\beta_8(1 + \beta_4)\eta_0(1 + 4\eta_0^2 e^{24\pi g^2}), \quad (4.13)$$

and

$$\begin{aligned} & 12g(c_1\eta_0(3 + c_1\eta_0 + 4\beta_2 + \beta_5 + 4\beta_2'\beta_4\beta_8(1 + c_1\eta_0)) + 2\beta_2 + \beta_5)e^{12g^2(3 + c_1^2\eta_0^2((\beta_2 + \beta_5)^2 + (\beta_4\beta_7c_1\eta_0)^2))} \\ & + 6\beta_1\beta_4c_1\eta_0((\beta_6 + 4\beta_4\beta_8c_1\eta_0)(1 + c_1\eta_0) + 2\beta_4\beta_8(1 + 2c_1\eta_0 + c_1^2\eta_0^2) + 1). \end{aligned} \quad (4.14)$$

Let γ_0 (resp. T_0) be chosen so that

$$\gamma_0 \leq \min\left\{1, \frac{e^{-12\pi g^2}}{8\beta_1\beta_6\eta_0(1 + \eta_0 e^{12\pi g^2})}, \frac{e^{-24\pi g^2}}{(16\beta_1(1 + \beta_4)\beta_6\eta_0(1 + 2\eta_0 e^{12\pi g^2}))^2}, \frac{1}{4\beta_1\beta_6(1 + 2\beta_4)c_1\eta_0(2 + c_1\eta_0)}, \frac{1}{16\bar{c}_1^2}\right\}, \quad (4.15)$$

$$T_0 \leq 1, \quad T_0 \leq \frac{1}{16\bar{c}_1^2}. \quad (4.16)$$

Let us prove by induction the existence and uniqueness of sequences $(\tilde{\Phi}^n)$ in $C([0, T_0], H_{\text{per}}^1(0, 2\pi))$ and (\tilde{R}^n) in $C([0, T_0], L_{\frac{P}{1+P}}^2(\mathbb{R}^3, H_{\text{per}}^1(0, 2\pi)))$ solutions to

$$\tilde{\Phi}^0 = 0, \quad \tilde{R}^0 = 0, \quad (4.17)$$

$$\partial_t \tilde{\Phi}^{n+1} - i\partial_x^2 \tilde{\Phi}^{n+1} = S_1(\tilde{R}^n)\tilde{\Phi}^{n+1} - ig\overline{\tilde{\Phi}^{n+1}} + U(\tilde{\Phi}^n, \tilde{R}^n), \quad \tilde{\Phi}^{n+1}(0, x) = \tilde{\Phi}_i(x), \quad (4.18)$$

$$\partial_t \tilde{R}^{n+1} + p_x \partial_x \tilde{R}^{n+1} = g\gamma \left(L\tilde{R}^{n+1} + \gamma(L_{1, \tilde{\Phi}^n} \tilde{R}^{n+1} + Q_{1, \tilde{\Phi}^n}(\tilde{R}^n, \tilde{R}^n)) \right), \quad \tilde{R}^{n+1}(0, \cdot, \cdot) = \tilde{R}_i, \quad (4.19)$$

and such that for any $T \leq T_0$ and $\gamma \leq \gamma_0$,

$$\begin{aligned} & \| \delta_n \tilde{\Phi} \|_{\infty T} + \| \delta_n \tilde{R} \|_{\infty, 2, H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}} \delta_n \tilde{R} \|_{2, 2, H^1} \\ & \leq \bar{c}_1 (\sqrt{T} + \sqrt{\gamma}) (\| \delta_{n-1} \tilde{\Phi} \|_{\infty T} + \| \delta_{n-1} \tilde{R} \|_{\infty, 2, H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}} \delta_{n-1} \tilde{R} \|_{2, 2, H^1}), \end{aligned} \quad (4.20)$$

and

$$\| \tilde{\Phi}^n \|_{\infty T} + \| \tilde{R}^n \|_{\infty, 2, H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}} \tilde{R}^n \|_{2, 2, H^1} \leq c_1 (\| \tilde{\Phi}_i \|_{H^1} + \| \tilde{R}_i \|_{2, H^1}), \quad n \in \mathbb{N}. \quad (4.21)$$

Here $\delta_n \tilde{\Phi} = \tilde{\Phi}^{n+1} - \tilde{\Phi}^n$ and $\delta_n \tilde{R} = \tilde{R}^{n+1} - \tilde{R}^n$.

The existence and uniqueness of $\tilde{\Phi}^1$ in $C([0, T_0], H_{per}^1(0, 2\pi))$ follow from Lemma 3.1. By (3.2) and the first bound on T_0 in (4.16),

$$\| \tilde{\Phi}^1 \|_{\infty T} \leq 2 \| \tilde{\Phi}_i \|_{H^1} e^{12\pi g^2}, \quad T \in]0, T_0]. \quad (4.22)$$

The existence and uniqueness of \tilde{R}^1 in $C([0, T_0], L^2_{\frac{P}{1+P}}(\mathbb{R}^3, H_{per}^1(0, 2\pi)))$ follow from analogous arguments for the solution of the linearized Boltzmann equation. Lemmas 3.4-5 apply to \tilde{R}^1 and $\partial_x \tilde{R}^1$, by the assumption (4.3) on \tilde{R}_i , and because by periodicity

$$\int \partial_x \tilde{R}^1(t, x, p) (|p|^2 + g) P dx dp = \int \partial_x \tilde{R}^1(t, x, p) p_x P dx dp = 0.$$

Hence,

$$\| \tilde{R}^1 \|_{\infty, 2, H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}} \tilde{R}^1 \|_{2, 2, H^1} \leq \beta_1 \| \tilde{R}_i \|_{2, H^1}, \quad (4.23)$$

so that

$$\| \tilde{\Phi}^1 \|_{\infty, T} + \| \tilde{R}^1 \|_{\infty, 2, H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}} \tilde{R}^1 \|_{2, 2, H^1} \leq \frac{C_1}{2} (\| \tilde{\Phi}_i \|_{H^1} + \| \tilde{R}_i \|_{2, H^1}). \quad (4.24)$$

And so, (4.21) holds for $n = 1$.

The existence and uniqueness of $\tilde{\Phi}^2$ in $C([0, T_0], H_{per}^1(0, 2\pi))$ follow from Lemma 3.1, since $S_1(\tilde{R}^1)$ and $U(\tilde{\Phi}^1, \tilde{R}^1)$ belong to $L^\infty(0, T_0; H_{per}^1(0, 2\pi))$, $H^1(0, 2\pi)$ being an algebra. By the proof of Lemma 3.1,

$$\begin{aligned} \| \delta_1 \tilde{\Phi} \|_{\infty T} & \leq 3\sqrt{T} \left(\int_0^T \| (U(\tilde{\Phi}^1, \tilde{R}^1) + (S_1(\tilde{R}^1) + ig)\tilde{\Phi}^1)(r) \|_{H^1}^2 dr \right)^{\frac{1}{2}} e^{3(\int_0^T \| S^1(\tilde{R}^1)(r) \|_{H^1}^2 dr + 2\pi g^2)}, \\ & \quad T \in]0, T_0]. \end{aligned}$$

Moreover, by (4.4) and (4.6),

$$\begin{aligned} & \| U(\tilde{\Phi}^1, \tilde{R}^1)(r) + (S^1(\tilde{R}^1)(r) + ig)\tilde{\Phi}^1 \|_{H^1} \\ & \leq 2g(1 + \| \tilde{\Phi}^1 \|_{\infty T}) (\beta_5 + \beta_2 + \beta_4 \beta_7 \gamma^2 \| \nu^{\frac{1}{2}} \tilde{R}^1 \|_{2, 2, H^1}) \| \tilde{R}^1(r) \|_{2, H^1} \\ & + 10g \| \tilde{\Phi}^1 \|_{\infty T}^2 (1 + \| \tilde{\Phi}^1 \|_{\infty T}), \quad r \in [0, 1]. \end{aligned}$$

And so,

$$\begin{aligned} & \| \delta_1 \tilde{\Phi} \|_{\infty T} \\ & \leq 30g\sqrt{T} (1 + \| \tilde{\Phi}^1 \|_{\infty T}) \left((\beta_5 + \beta_2 + \beta_4 \beta_7 \gamma^2 \| \nu^{\frac{1}{2}} \tilde{R}^1 \|_{2, 2, H^1}) \| \tilde{R}^1 \|_{\infty, 2, H^1} + \| \tilde{\Phi}^1 \|_{\infty T}^2 \right) \\ & \quad \times e^{60g^2(1 + (\beta_2^2 + \beta_5^2)) \| \tilde{R}^1 \|_{2, 2, H^1}^2 + \beta_4^2 \beta_7^2 \gamma \| \tilde{R}^1 \|_{\infty, 2, H^1}^2 \| \nu^{\frac{1}{2}} \tilde{R}^1 \|_{T, 2, 2}^2} \\ & \leq 30g\sqrt{T} (1 + 2\eta_0 e^{12\pi g^2}) \left((\beta_2 + \beta_5 + \beta_1 \beta_4 \beta_7 \eta_0) \| \tilde{R}^1 \|_{\infty, 2, H^1} + 2\eta_0 e^{12\pi g^2} \| \tilde{\Phi}^1 \|_{\infty T} \right) \\ & \quad \times e^{60g^2(1 + \beta_1^2 \eta_0^2 (\beta_2^2 + \beta_5^2 + (\beta_1 \beta_4 \beta_7 \eta_0)^2))}. \end{aligned} \quad (4.25)$$

A solution $\tilde{R}^2 \in C([0, T_0], L^2_{\frac{P}{1+P}}(\mathbb{R}^3, H^1_{per}(0, 2\pi)))$ to (4.19) can be obtained as the limit of the sequence $(r^k)_{k \in \mathbb{N}}$ defined by

$$\begin{aligned} r^0 &= 0, \quad \partial_t r^{k+1} + p_x \partial_x r^{k+1} = g\gamma \left(Lr^{k+1} + \gamma(L_{1\Phi^1} r^k + Q_{1\Phi^1}(\tilde{R}^1, \tilde{R}^1)) \right), \\ r^k(0, \cdot, \cdot) &= \tilde{R}_i. \end{aligned}$$

By Lemma 3.4, (4.5) and the second condition in (4.15),

$$\begin{aligned} \|\nu^{\frac{1}{2}} \delta_k r\|_{T,2,2} &\leq \gamma \beta_1 \beta_6 (2 + \|\Phi^1\|_{\infty T}) \|\Phi^1\|_{\infty T} \|\nu^{\frac{1}{2}} \delta_{k-1} r\|_{T,2,2} \\ &\leq \frac{1}{2} \|\nu^{\frac{1}{2}} \delta_{k-1} r\|_{T,2,2}, \quad k \in \mathbb{N}^*. \end{aligned}$$

And so, by a contraction argument, the local existence and uniqueness of \tilde{R}^2 follow. Applying Lemmas 3.4-5 to $\delta_1 \tilde{R}$ leads to

$$\begin{aligned} &\|\delta_1 \tilde{R}\|_{\infty, 2, H^1} + \sqrt{\gamma} \|\nu^{\frac{1}{2}} \delta_1 \tilde{R}\|_{2, 2, H^1} \\ &\leq \beta_1 \gamma^{\frac{3}{2}} \left(\|(\Phi^1 + \bar{\Phi}^1 + \gamma|\Phi^1|^2) \nu^{-\frac{1}{2}} L \delta_1 \tilde{R}\|_{2, 2, H^1} + \|(\Phi^1 + \bar{\Phi}^1 + \gamma|\Phi^1|^2) \nu^{-\frac{1}{2}} L \tilde{R}^1\|_{2, 2, H^1} \right. \\ &\quad \left. + \|\nu^{-\frac{1}{2}} Q_{1, \Phi^1}(\tilde{R}^1, \tilde{R}^1)\|_{2, 2, H^1} \right) \\ &\leq \beta_1 (1 + \beta_4) \gamma^{\frac{3}{2}} \left(2\beta_6 \|\Phi^1\|_{\infty T} (1 + \|\Phi^1\|_{\infty T}) (\|\nu^{\frac{1}{2}} \delta_1 \tilde{R}\|_{2, 2, H^1} + \|\nu^{\frac{1}{2}} \tilde{R}^1\|_{2, 2, H^1}) \right. \\ &\quad \left. + 4\beta_4 \beta_8 (1 + \|\Phi^1\|_{\infty T}^2) \|\tilde{R}^1\|_{\infty, 2, H^1} \|\nu^{\frac{1}{2}} \tilde{R}^1\|_{2, 2, H^1} \right). \end{aligned}$$

And so, by (4.22), (4.23) and the third condition in (4.15),

$$\|\delta_1 \tilde{R}\|_{\infty, 2, H^1} + \frac{3}{4} \sqrt{\gamma} \|\nu^{\frac{1}{2}} \delta_1 \tilde{R}\|_{2, 2, H^1} \leq \left(\frac{1}{4} + 4\beta_1^2 \beta_4 \beta_8 (1 + \beta_4) \eta_0 (1 + 4\eta_0^2 e^{24\pi g^2}) \right) \gamma \|\nu^{\frac{1}{2}} \tilde{R}^1\|_{2, 2, H^1}. \quad (4.26)$$

It results from (4.25), (4.26) and the bounds from below (4.12)-(4.13) on \bar{c}_1 that

$$\begin{aligned} &\|\delta_1 \tilde{\Phi}\|_{\infty T} + \|\delta_1 \tilde{R}\|_{\infty, 2, H^1} + \sqrt{\gamma} \|\nu^{\frac{1}{2}} \delta_1 \tilde{R}\|_{2, 2, H^1} \\ &\leq \bar{c}_1 (\sqrt{T} + \sqrt{\gamma}) (\|\tilde{\Phi}^1\|_{\infty, T} + \|\tilde{R}^1\|_{\infty, 2, H^1} + \sqrt{\gamma} \|\nu^{\frac{1}{2}} \tilde{R}^1\|_{2, 2, H^1}). \end{aligned} \quad (4.27)$$

And so, (4.20) holds for $n = 1$.

The existence and uniqueness of $(\tilde{\Phi}^n)_{n \geq 3}$ in $C([0, T_0], H^1_{per}(0, 2\pi))$ follow from Lemma 3.1, since by induction on n and $H^1(0, 2\pi)$ being an algebra, $S_1(\tilde{R}^n)$ and $U(\tilde{\Phi}^n, \tilde{R}^n)$ belong to $L^\infty(0, T_0; H^1_{per}(0, 2\pi))$.

The existence and uniqueness of $(\tilde{R}^n)_{n \geq 3}$ in $C([0, T_0], L^2_{\frac{P}{1+P}}(\mathbb{R}^3, H^1_{per}(0, 2\pi)))$ follow from similar arguments to those previously used for the existence and uniqueness of \tilde{R}^2 .

Assuming (4.20)-(4.21) up to $n - 1$, and using the second (resp. fourth) condition in (4.16) (resp. (4.15)) implies that (4.21) holds for n . Then $\delta_n \tilde{\Phi}$ and $\delta_n \tilde{R}$ satisfy

$$\begin{aligned} \partial_t \delta_n \tilde{\Phi} - i \partial_x^2 \delta_n \tilde{\Phi} &= S_1(\tilde{R}^n) \delta_n \tilde{\Phi} - i g \overline{\delta_n \tilde{\Phi}} + \Gamma_{n-1}, \quad \delta_n \tilde{\Phi}(0, x) = 0, \\ \partial_t \delta_n \tilde{R} + p_x \partial_x \delta_n \tilde{R} &= g\gamma \left(L \delta_n \tilde{R} + \gamma(L_{1, \tilde{\Phi}^n} \delta_n \tilde{R} + G_{n-1}) \right), \quad \delta_n \tilde{R}(0, x, p) = 0. \end{aligned}$$

Here

$$\begin{aligned} \Gamma_{n-1} &= \tilde{\Phi}^n \delta_{n-1}(S_1(\tilde{R})) + \delta_{n-1} U(\tilde{\Phi}, \tilde{R}), \\ G_{n-1} &= \left(\delta_{n-1} \tilde{\Phi} + \overline{\delta_{n-1} \tilde{\Phi}} + \gamma(\tilde{\Phi}^n \overline{\delta_{n-1} \tilde{\Phi}} + \overline{\tilde{\Phi}^{n-1} \delta_{n-1} \tilde{\Phi}}) \right) L \tilde{R}^n + \delta_{n-1} Q_{1, \tilde{\Phi}}(\tilde{R}, \tilde{R}), \end{aligned}$$

and $(G_{n-1})_{\parallel} = 0$. Applying Lemma 3.1, it holds that

$$\|\delta_n \tilde{\Phi}\|_{\infty T}^2 \leq 6T \left(\int_0^T \|\Gamma_{n-1}(t)\|_{H^1}^2 dt \right) e^{6T \int_0^T (\|S_1(\tilde{R}^n)(t, \cdot)\|_{H^1}^2 + 2\pi g^2) dt}.$$

By definition of $S_1(\tilde{R})$ and $U(\tilde{\Phi}, \tilde{R})$,

$$\begin{aligned} \Gamma_{n-1} = & -\frac{g}{2} \gamma \tilde{\Phi}^n \left(\gamma \int PL \delta_{n-1} \tilde{R} dp + 4i \int P \delta_{n-1} \tilde{R} dp + \gamma^2 \int (Q(\tilde{R}^n, \delta_{n-1} \tilde{R}) + Q(\delta_{n-1} \tilde{R}, \tilde{R}^{n-1})) dp \right) \\ & -\frac{g}{2} \gamma \int PL \delta_{n-1} \tilde{R} dp - 2ig \int P \delta_{n-1} \tilde{R} dp \\ & -ig\gamma ((2\tilde{\Phi}^n + \tilde{\Phi}^{n-1} + \tilde{\Phi}^n) \delta_{n-1} \tilde{\Phi} + 2\tilde{\Phi}^{n-1} \delta_{n-1} \tilde{\Phi}) \\ & -\frac{g}{2} \gamma^2 \int (Q(\tilde{R}^n, \delta_{n-1} \tilde{R}) + Q(\delta_{n-1} \tilde{R}, \tilde{R}^{n-1})) dp \\ & -ig\gamma^2 ((\tilde{\Phi}^n + \tilde{\Phi}^{n-1}) \tilde{\Phi}^n \delta_{n-1} \tilde{\Phi} + (\tilde{\Phi}^{n-1})^2 \delta_{n-1} \tilde{\Phi}). \end{aligned}$$

It follows from the Cauchy-Schwartz inequality w.r.t. the p variable and (4.7) that for every $t \in [0, T]$,

$$\begin{aligned} \left(\int \left(\int Q(\tilde{R}^n, \delta_{n-1} \tilde{R})(t, x, p) dp \right)^2 dx \right)^{\frac{1}{2}} & \leq \beta'_2 \left(\int \frac{P}{1+P} \nu^{-1} \left(\frac{Q(\tilde{R}^n, \delta_{n-1} \tilde{R})}{P} \right)^2 dp dx \right)^{\frac{1}{2}} \\ & \leq \beta'_2 \beta_8 \left(\sup_{x \in [0, 2\pi]} \|\tilde{R}^n(t, x, \cdot)\|_{L^2 \frac{P}{1+P}} \right) \|\nu^{\frac{1}{2}} \delta_{n-1} \tilde{R}(t)\|_{2,2} \\ & \leq \beta'_2 \beta_4 \beta_8 \|\tilde{R}^n(t)\|_{2, H^1} \|\nu^{\frac{1}{2}} \delta_{n-1} \tilde{R}(t)\|_{2,2}. \end{aligned}$$

Analogously, $\partial_x Q(\tilde{R}^n, \delta_{n-1} \tilde{R})$ being equal to $Q(\partial_x \tilde{R}^n, \delta_{n-1} \tilde{R}) + Q(\tilde{R}^n, \partial_x \delta_{n-1} \tilde{R})$,

$$\begin{aligned} & \left(\int \left(\partial_x \int Q(\tilde{R}^n, \delta_{n-1} \tilde{R})(t, x, p) dp \right)^2 dx \right)^{\frac{1}{2}} \\ & \leq \beta'_2 \beta_4 \beta_8 \left(\|\delta_{n-1} \tilde{R}(t)\|_{2, H^1} \|\nu^{\frac{1}{2}} \partial_x \tilde{R}^n(t)\|_{2,2} + \|\tilde{R}^n(t)\|_{2, H^1} \|\nu^{\frac{1}{2}} \partial_x \delta_{n-1} \tilde{R}(t)\|_{2,2} \right). \end{aligned}$$

And so,

$$\begin{aligned} & \left\| \int Q(\tilde{R}^n, \delta_{n-1} \tilde{R})(t, \cdot, p) dp \right\|_{H^1} \\ & \leq 2\beta'_2 \beta_4 \beta_8 \left(\|\tilde{R}^n(t)\|_{2, H^1} \|\nu^{\frac{1}{2}} \delta_{n-1} \tilde{R}(t)\|_{2, H^1} + \|\nu^{\frac{1}{2}} \tilde{R}^n(t)\|_{2, H^1} \|\delta_{n-1} \tilde{R}(t)\|_{2, H^1} \right). \end{aligned}$$

Consequently,

$$\begin{aligned} \|\Gamma_{n-1}(t)\|_{H^1} \leq & g\gamma \|\tilde{\Phi}^n\|_{\infty T} \left((\beta_5 \gamma + 4\beta_2) \|\delta_{n-1} \tilde{R}(t)\|_{2, H^1} \right. \\ & + \beta'_2 \beta_4 \beta_8 \gamma^2 (\|\tilde{R}^{n-1}(t)\|_{2, H^1} + \|\tilde{R}^n(t)\|_{2, H^1}) \|\nu^{\frac{1}{2}} \delta_{n-1} \tilde{R}(t)\|_{2, H^1} \\ & \left. + (\|\nu^{\frac{1}{2}} \tilde{R}^{n-1}(t)\|_{2, H^1} + \|\nu^{\frac{1}{2}} \tilde{R}^n(t)\|_{2, H^1}) \|\delta_{n-1} \tilde{R}(t)\|_{2, H^1} \right) \\ & + g(\beta_5 \gamma + 2\beta_2) \|\delta_{n-1} \tilde{R}(t)\|_{2, H^1} \\ & + 3g\gamma (\|\tilde{\Phi}^{n-1}\|_{\infty T} + \|\tilde{\Phi}^n\|_{\infty T}) \|\delta_{n-1} \tilde{\Phi}(t)\|_{H^1} \\ & + g\beta'_2 \beta_4 \beta_8 \gamma^2 (\|\tilde{R}^{n-1}(t)\|_{2, H^1} + \|\tilde{R}^n(t)\|_{2, H^1}) \|\nu^{\frac{1}{2}} \delta_{n-1} \tilde{R}(t)\|_{2, H^1} \\ & + (\|\nu^{\frac{1}{2}} \tilde{R}^{n-1}(t)\|_{2, H^1} + \|\nu^{\frac{1}{2}} \tilde{R}^n(t)\|_{2, H^1}) \|\delta_{n-1} \tilde{R}(t)\|_{2, H^1} \\ & + 2g\gamma^2 (\|\tilde{\Phi}^{n-1}\|_{\infty T}^2 + \|\tilde{\Phi}^n\|_{\infty T}^2) \|\delta_{n-1} \tilde{\Phi}(t)\|_{H^1}. \end{aligned}$$

Moreover, for any $t \in [0, T]$,

$$\| S_1(\tilde{R}^n)(t) \|_{H^1} \leq 2g \left(1 + \gamma(\beta_5\gamma + \beta_2) \| \tilde{R}^n(t) \|_{2,H^1} + \beta_4\beta_7\gamma^3 \| \tilde{R}^n(t) \|_{2,H^1} \| \nu^{\frac{1}{2}}\tilde{R}^n(t) \|_{2,H^1} \right).$$

And so, using (4.21) at steps $n-1$ and n , and the first condition in (4.16)

$$\begin{aligned} \| \delta_n \tilde{\Phi} \|_{\infty T} &\leq 6g\sqrt{T}e^{12g^2(3+c_1^2\eta_0^2((\beta_2+\beta_5)^2+(\beta_4\beta_7c_1\eta_0)^2))} \\ &\quad \left(2c_1\eta_0(3+2c_1\eta_0) \| \delta_{n-1}\tilde{\Phi} \|_{\infty T} + (c_1\eta_0(4\beta_2+\beta_5+2\beta_2'\beta_4\beta_8(1+c_1\eta_0))+2\beta_2+\beta_5) \| \delta_{n-1}\tilde{R} \|_{\infty,2,H^1} \right. \\ &\quad \left. + 2c_1\eta_0(1+c_1\eta_0)\beta_2'\beta_4\beta_8\sqrt{\gamma} \| \nu^{\frac{1}{2}}\delta_{n-1}\tilde{R} \|_{2,2,H^1} \right). \end{aligned} \quad (4.28)$$

Multiplying the equation for $\delta_n\tilde{R}$ by $(|p|^2+g)P$ (resp. p_xP) and integrating on $(0, 2\pi) \times \mathbb{R}^3$, gives

$$\frac{d}{dt} \int \delta_n \tilde{R}(t, x, p)(|p|^2+g)P dx dp = 0, \quad (\text{resp. } \frac{d}{dt} \int \delta_n \tilde{R}(t, x, p)p_x P dx dp = 0).$$

Indeed, it follows from Lemma 2.1 that

$$\int (L\delta_n\tilde{R})(|p|^2+g)P dp = \int (\delta_n\tilde{R})L((|p|^2+g)(1+P))\frac{P}{1+P} dp = 0.$$

Similarly, the $\delta_{n-1}Q_{1\tilde{\Phi}}(\tilde{R}, \tilde{R})(|p|^2+g)P$ term vanishes after integration, by the δ_0 factor in the definition of $Q(\tilde{R}, \tilde{R})$.

Being zero initially, $\int \delta_n\tilde{R}(t, x, p)(|p|^2+g)P dx dp$ and $\int \delta_n\tilde{R}(t, x, p)p_x P dx dp$ remain identically zero, so that Lemmas 3.4-5 apply to the equation for $\delta_n\tilde{R}$, and $\delta_n\partial_x\tilde{R}$. Hence,

$$\| \delta_n \tilde{R} \|_{\infty,2,H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}}\delta_n\tilde{R} \|_{2,2,H^1} \leq \beta_1\gamma^{\frac{3}{2}} \| \nu^{-\frac{1}{2}}(L_{1,\Phi_n}\delta_n s + G_{n-1}) \|_{2,2,H^1}. \quad (4.29)$$

Moreover,

$$\begin{aligned} \| \nu^{-\frac{1}{2}}L_{1,\tilde{\Phi}^n}\delta_n\tilde{R} \|_{2,2,H^1} &\leq \beta_6(1+2\beta_4) \| \tilde{\Phi}^n \|_{\infty T} (2+ \| \tilde{\Phi}^n \|_{\infty T}) \| \nu^{\frac{1}{2}}\delta_n\tilde{R}_\perp \|_{2,2,H^1} \\ &\leq \beta_6(1+2\beta_4)c_1\eta_0(2+c_1\eta_0) \| \nu^{\frac{1}{2}}\delta_n\tilde{R}_\perp \|_{2,2,H^1}, \end{aligned}$$

and, using (4.8),

$$\begin{aligned} &\| \nu^{-\frac{1}{2}}G_{n-1} \|_{2,2,H^1} \\ &\leq 2\beta_4 \left((2+ \| \tilde{\Phi}^{n-1} \|_{\infty T} + \| \tilde{\Phi}^n \|_{\infty T}) \| \delta_{n-1}\tilde{\Phi} \|_{\infty T} (\beta_6+4\gamma\beta_4\beta_8 \| \tilde{R}^n \|_{\infty,2,H^1}) \| \nu^{\frac{1}{2}}\tilde{R}^n \|_{2,2,H^1} \right. \\ &\quad \left. + 2\beta_4\beta_8(1+2 \| \tilde{\Phi}^{n-1} \|_{\infty T} + \| \tilde{\Phi}^{n-1} \|_{\infty T}^2) (\| \nu^{\frac{1}{2}}\tilde{R}^{n-1} \|_{2,2,H^1} + \| \nu^{\frac{1}{2}}\tilde{R}^n \|_{2,2,H^1}) \| \delta_{n-1}\tilde{R} \|_{\infty,2,H^1} \right. \\ &\quad \left. + (\| \tilde{R}^{n-1} \|_{\infty,2,H^1} + \| \tilde{R}^n \|_{\infty,2,H^1}) \| \nu^{\frac{1}{2}}\delta_{n-1}\tilde{R} \|_{2,2,H^1} \right) \\ &\leq 4\beta_4c_1\eta_0\gamma^{-\frac{1}{2}} \left((\beta_6+4\beta_4\beta_8c_1\eta_0)(1+c_1\eta_0) \| \delta_{n-1}\tilde{\Phi} \|_{\infty T} \right. \\ &\quad \left. + 2\beta_4\beta_8(1+2c_1\eta_0+c_1^2\eta_0^2) \| \delta_{n-1}\tilde{R} \|_{\infty,2,H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}}\delta_{n-1}\tilde{R} \|_{2,2,H^1} \right). \end{aligned}$$

And so, using (4.28), the fourth condition on γ_0 in (4.15), in order to move the

$$\beta_1\beta_6(1+2\beta_4)c_1\eta_0(2+c_1\eta_0)\gamma^{\frac{3}{2}} \| \nu^{\frac{1}{2}}\delta_n\tilde{R}_\perp \|_{2,2,H^1}$$

term from the r.h.s. of (4.29) to its l.h.s., and using the bound from below (4.14) of \bar{c}_1 ,

$$\begin{aligned} &\| \delta_n \tilde{\Phi} \|_{\infty T} + \| \delta_n \tilde{R} \|_{\infty,2,H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}}\delta_n\tilde{R} \|_{2,2,H^1} \\ &\leq \bar{c}_1(\sqrt{T} + \sqrt{\gamma})(\| \delta_{n-1}\tilde{\Phi} \|_{\infty T} + \| \delta_{n-1}\tilde{R} \|_{\infty,2,H^1} + \sqrt{\gamma} \| \nu^{\frac{1}{2}}\delta_{n-1}\tilde{R} \|_{2,2,H^1}). \end{aligned} \quad (4.30)$$

This proves the induction (4.20) for n .

It follows from (4.20)-(4.21), and the fifth (resp. second) condition in (4.15) (resp. (4.16)), that the sequence $(\tilde{\Phi}^n, \tilde{R}^n)$ converges in

$L^\infty(0, T_0; H_{per}^1(0, 2\pi)) \times L^\infty(0, T_0; L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H_{per}^1(0, 2\pi)))$ when $n \rightarrow +\infty$ to a solution

$$(\tilde{\Phi}, \tilde{R}) \in L^\infty(0, T_0; H_{per}^1(0, 2\pi)) \times L^\infty(0, T_0; L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H_{per}^1(0, 2\pi)))$$

of (4.1-2), satisfying

$$\|\tilde{\Phi}\|_{\infty T_0} + \|\tilde{R}\|_{\infty, 2, H^1} + \sqrt{\gamma} \|\nu^{\frac{1}{2}} \tilde{R}\|_{2, 2, H^1} \leq c_1 (\|\tilde{\Phi}_i\|_{H^1} + \|\tilde{R}_i\|_{2, H^1}). \quad (4.31)$$

The solution belongs to $W^{1,1}(0, T_0; H_{per}^1(0, 2\pi)) \times W^{1,1}(0, T_0; L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H_{per}^1(0, 2\pi)))$, hence

$$(\tilde{\Phi}, \tilde{R}) \in C_b([0, T_0]; H_{per}^1(0, 2\pi)) \times C_b([0, T_0]; L^2_{\frac{P}{1+P}}(\mathbb{R}^3; H_{per}^1(0, 2\pi))).$$

The uniqueness of the solution to (4.1-2) follows similarly, considering the difference of two solutions. \blacksquare

The following lemma on the kinetic and internal energies of ψ , will also be needed to prove the global in time existence result of Theorem 1.1.

Lemma 4.2 *The solution $(f, \psi) = (P(1 + \gamma\tilde{R}), \psi)$ of (1.6-9) satisfies*

$$\begin{aligned} \frac{d}{dt} \int \left(|\partial_x \psi|^2 + \frac{g}{2} (|\psi|^2 - 1)^2 \right) (t, x) dx &= 2ig\gamma \int (\bar{\psi} \partial_x \psi - \psi \partial_x \bar{\psi}) \int P \partial_x \tilde{R} dp dx \\ &- \frac{g}{2} \gamma^2 \int (\bar{\psi} \partial_x \psi + \psi \partial_x \bar{\psi}) \int (PL \partial_x \tilde{R} + \gamma \partial_x Q(\tilde{R}, \tilde{R})) dp dx - g\gamma^2 \int |\partial_x \psi|^2 \int (PL \tilde{R} + \gamma Q(\tilde{R}, \tilde{R})) dp dx \\ &- g^2 \gamma^2 \int |\psi|^2 (|\psi|^2 - 1) \int (PL \tilde{R} + \gamma Q(\tilde{R}, \tilde{R})) dp dx. \end{aligned} \quad (4.32)$$

Proof of Lemma 4.2.

Given (1.10), equation (1.8) satisfied by ψ is

$$\partial_t \psi - i \partial_x^2 \psi = -\psi \left(\frac{D}{2} + iA \right), \quad (4.33)$$

where

$$D = g\gamma^2 \int (PL \tilde{R} + \gamma Q(\tilde{R}, \tilde{R})) dp, \quad A = g(|\psi|^2 - 1 + 2\gamma \int P \tilde{R} dp).$$

Multiply (4.33) (resp. the conjugate of (4.33)) by $\partial_t \bar{\psi}$ (resp. $-\partial_t \psi$), integrate on $[0, 2\pi]$ so that

$$\begin{aligned} \frac{d}{dt} \int \left(|\partial_x \psi|^2 + \frac{g}{2} (|\psi|^2 - 1)^2 \right) (t, x) dx &= \frac{i}{2} \int D (\psi \partial_t \bar{\psi} - \bar{\psi} \partial_t \psi) dx - 2g\gamma \int (\psi \partial_t \bar{\psi} + \bar{\psi} \partial_t \psi) \int P \tilde{R} dp dx \\ &= 2ig\gamma \int (\psi \partial_x^2 \bar{\psi} - \bar{\psi} \partial_x^2 \psi) \int P \tilde{R} dp dx + 2g\gamma \int |\psi|^2 D \int P \tilde{R} dp dx \\ &\quad + \frac{1}{2} \int (\psi \partial_x^2 \bar{\psi} + \bar{\psi} \partial_x^2 \psi) D dx - \int |\psi|^2 A D dx \\ &= 2ig\gamma \int (\bar{\psi} \partial_x \psi - \psi \partial_x \bar{\psi}) \int P \partial_x \tilde{R} dp dx + 2g\gamma \int |\psi|^2 D \int P \tilde{R} dp dx \\ &\quad - \frac{1}{2} \int (\psi \partial_x \bar{\psi} + \bar{\psi} \partial_x \psi) \partial_x D dx - \int |\partial_x \psi|^2 D dx - \int |\psi|^2 A D dx. \end{aligned}$$

This proves the lemma. ■

Proof of Theorem 1.1.

Defining $s = e^{\zeta t} \tilde{R}$, we look for a solution $(\tilde{\Phi}, s)$ to the equations

$$\partial_t \tilde{\Phi} - i \partial_x^2 \tilde{\Phi} = S_1(s) \tilde{\Phi} - i g \tilde{\Phi} + U(\tilde{\Phi}, s), \quad \tilde{\Phi}(0, \cdot) = \tilde{\Phi}_i, \quad (4.34)$$

$$\partial_t s + p_x \partial_x s = g \gamma \left(Ls + \gamma (L_{1, \tilde{\Phi}} s + Q_{1, \tilde{\Phi}}(s, s)) \right) + \zeta s, \quad s(0, \cdot, \cdot) = \tilde{R}_i, \quad (4.35)$$

obtained from (4.1) and (4.2).

Here ζ will be the positive rate of an exponential in time decay of \tilde{R} . Set $c_\zeta = \frac{\zeta}{\gamma}$.

Let γ_0 be given as in Proposition 4.1 when the norm of the initial conditions is bounded by $\eta_0 = 1$.

Let c_ζ and $(c_i)_{2 \leq i \leq 7}$ be the constants defined by

$$c_\zeta = \min \left\{ 1, \frac{10}{\sqrt{1 + g(1 + \beta_3^4)}}, \frac{g\nu_0}{4\beta_1(1 + \sqrt{\nu_0})} \right\}, \quad (4.36)$$

$$c_2 = 4\beta_1 \left(4\beta_1 \beta_4 c_4 \left(\beta_6 + \frac{\beta_8}{\sqrt{\nu_0}} \right) + 1 \right), \quad (4.37)$$

$$c_3 = \sqrt{2} g c_4 \left(2\sqrt{\beta_2} + \beta_4 + \beta_5 \right), \quad (4.38)$$

$$c_4 = \beta_4 (\mathcal{M}_0 + 1)^{\frac{1}{2}}, \quad (4.39)$$

$$c_5 = 4\beta_4 \left(\frac{1}{g} + 2c_4^2 \right) + g + \frac{g c_4^2}{\sqrt{c_\zeta}} (\beta_5 + \beta_4 \beta_7), \quad (4.40)$$

$$c_6 = g \beta_5 (3\beta_4 + \sqrt{g}), \quad (4.41)$$

$$c_7 = \frac{1}{10} + g \beta_4 \beta_7 (3\beta_4 + \sqrt{g}). \quad (4.42)$$

Additionally, it is required that

$$\eta_0 = \min \left\{ 1, \frac{1}{4\beta_1(\beta_6 c_5 + 2\beta_4 \beta_8 c_4^2)}, \frac{1}{4\beta_1(\beta_6 c_5 + 4\beta_4 c_4(\beta_6 + \beta_8 c_4 + \frac{\beta_8}{\sqrt{\nu_0}}))} \right\}, \quad (4.43)$$

and

$$\gamma_0 \leq \min \left\{ 1, \frac{1}{\eta_0 \sqrt{2}}, \frac{1}{5c_6}, \frac{1}{25c_7^2} \right\}. \quad (4.44)$$

Assume that

$$\| \tilde{\Phi}_i \|_{H^1} \leq \frac{\eta \zeta}{10c_1}, \quad \| \tilde{R}_i \|_{2, H^1} \leq \frac{\eta \zeta}{10c_1(1 + c_2)(1 + c_3)}, \quad \text{with } \eta \leq \eta_0. \quad (4.45)$$

Set

$$\alpha(t) = \int (|\partial_x \psi|^2 + \frac{g}{2} (|\psi|^2 - 1)^2)(t, x) dx.$$

To start, there is by Proposition 4.1 a unique solution to problem (4.1-2) on a time interval $[0, T_0]$. From the computation,

$$\int (|\psi_i(x)|^2 - 1)^2 dx = \gamma^2 \int ((\tilde{\Phi}_i + \bar{\tilde{\Phi}}_i)(x) + \gamma|\tilde{\Phi}_i|^2)^2 dx \leq 2\gamma^2 \int (|\tilde{\Phi}_i(x)|^2 + |\tilde{\Phi}_i(x)|^4) dx,$$

it holds that

$$\alpha(0) \leq (1 + g(1 + \beta_3^4))\gamma^2 \|\tilde{\Phi}_i\|_{H^1}^2,$$

hence

$$\alpha(0) \leq \gamma^2 \eta^2, \quad (4.46)$$

by (4.45) and the second bound in (4.36). Consider the set of times $t_1 \leq T_0$ such that on $[0, t_1)$ the solution of (4.1-2) exists and satisfies

$$\alpha(t) \leq 2\gamma^2 \eta^2 \quad \text{and} \quad \|s(t)\|_{2,2}^2 + \|\partial_x s(t)\|_{2,2}^2 + \gamma(\|\nu^{\frac{1}{2}}s\|_{T_1,2,2}^2 + \|\nu^{\frac{1}{2}}\partial_x s\|_{T_1,2,2}^2) \leq 2\eta^2. \quad (4.47)$$

This set is nonempty by continuity. Denote by T_1 its upper bound. We shall next prove that $T_1 = T_0$ and improve the bounds (4.47), which will allow the solution to be continued beyond T_0 . That result is a main step in the proof of global existence, since it will imply that a solution will, as long as it can be continued, stay within the bounds of (4.47).

On $[0, T_1] \times [0, 2\pi]$,

$$|\psi(t, x)| \leq \beta_4 \left(\int (|\psi(t, x)|^2 + |\partial_x \psi(t, x)|^2) dx \right)^{\frac{1}{2}} \leq \beta_4 (\mathcal{M}_0 + \alpha(t))^{\frac{1}{2}} \leq \beta_4 (\mathcal{M}_0 + 1)^{\frac{1}{2}} = c_4, \quad (4.48)$$

by the bound on $\alpha(t)$ in (4.47), the second bound in (4.44) and the definition of c_4 . Moreover,

$$\begin{aligned} \left| |\psi(t, x)|^2 - 1 - \frac{1}{2\pi} \int (|\psi(t, y)|^2 - 1) dy \right| &\leq \beta_4 \left(\int (|\psi(t, u)|^2 - 1 - \frac{1}{2\pi} \int (|\psi(t, y)|^2 - 1) dy)^2 du \right. \\ &\quad \left. + \int (\psi \partial_x \bar{\psi} + \bar{\psi} \partial_x \psi)^2(t, u) du \right) \\ &\leq \beta_4 \left(\int (|\psi(t, u)|^2 - 1)^2 + 4c_4^2 |\partial_x \psi(t, u)|^2 du \right) \\ &\leq \beta_4 \left(\frac{2}{g} + 4c_4^2 \right) \alpha(t) \\ &\leq 4\beta_4 \left(\frac{1}{g} + 2c_4^2 \right) \gamma \eta, \quad t \in [0, T_1]. \end{aligned}$$

Multiplying (1.8) (resp. the conjugate of (1.8)) by $\bar{\psi}$ (resp. ψ), integrating w.r.t. x and adding the resulting equations leads to

$$\begin{aligned} \left| \frac{d}{dt} \int (|\psi(t, x)|^2 - 1) dx \right| &= g\gamma^2 \left| \int |\psi(t, x)|^2 \int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R})) dp dx \right| \\ &\leq gc_4^2 \sqrt{2\pi} \gamma^2 e^{-\zeta t} \|s(t)\|_{2,2} (\beta_5 + \beta_4 \beta_7 \gamma e^{-\zeta t} \|\nu^{\frac{1}{2}}s(t)\|_{2,2}). \end{aligned}$$

And so, using the Cauchy-Schwartz inequality when integrating the previous inequality on $[0, t]$,

$$\begin{aligned} \left| |\psi(t, x)|^2 - 1 \right| &\leq 4\beta_4 \left(\frac{1}{g} + 2c_4^2 \right) \gamma \eta + \frac{1}{2\pi} \left| \int (|\psi_i(x)|^2 - 1) dx \right| + \frac{g}{\sqrt{2\pi}} c_4^2 \beta_5 \gamma^2 (2\zeta)^{-\frac{1}{2}} \|s\|_{T_1,2,2} \\ &\quad + \frac{g}{\sqrt{2\pi}} c_4^2 \beta_4 \beta_7 \gamma^3 (4\zeta)^{-\frac{1}{2}} \sup_{t \in [0, T_1]} \|s(t)\|_{2,2} \|\nu^{\frac{1}{2}}s\|_{2,2,H^1}. \end{aligned}$$

Using (4.46) and the s -part of (4.47), leads to

$$| |\psi(t, x)|^2 - 1 | \leq c_5 \gamma \eta, \quad (4.49)$$

with c_5 defined in (4.40).

On $[0, T_1]$, consider the function s which is a solution to

$$\begin{aligned} \partial_t s + p_x \partial_x s &= g\gamma \left(Ls + \gamma (L_1 \psi s + e^{-\zeta t} |\psi|^2 \frac{Q(s, s)}{P} + \frac{\zeta}{g\gamma^2} s) \right), \\ s(0, x, p) &= \tilde{R}_i(x, p). \end{aligned}$$

By Lemma 3.4, (4.5), (4.7), (4.48), (4.49), the third bound on c_ζ in (4.36), and the s -part of (4.47),

$$\begin{aligned} & \sup_{t \in [0, T_1]} \| s(t) \|_{2,2} + \frac{3}{4} \sqrt{\gamma} \| \nu^{\frac{1}{2}} s \|_{T_1, 2, 2} \\ & < \beta_1 \left(\| \tilde{R}_i \|_{2,2} + \beta_6 c_5 \gamma^{\frac{3}{2}} \eta \| \nu^{\frac{1}{2}} s_\perp \|_{T_1, 2, 2} + \beta_4 \beta_8 c_4^2 \gamma^{\frac{3}{2}} \| s \|_{\infty, 2, H^1} \| \nu^{\frac{1}{2}} s \|_{T_1, 2, 2} \right) \\ & \leq \beta_1 \left(\| \tilde{R}_i \|_{2,2} + (\beta_6 c_5 + 2\beta_4 \beta_8 c_4^2) \gamma^{\frac{3}{2}} \eta_0 \| \nu^{\frac{1}{2}} s \|_{T_1, 2, 2} \right). \end{aligned}$$

It follows from the second bound on η_0 in (4.43) that

$$\sup_{t \in [0, T_1]} \| s(t) \|_{2,2} + \frac{\sqrt{\gamma}}{2} \| \nu^{\frac{1}{2}} s \|_{T_1, 2, 2} < \beta_1 \| \tilde{R}_i \|_{2,2}. \quad (4.50)$$

The function $\partial_x s$ satisfies

$$\begin{aligned} \partial_t \partial_x s + p_x \partial_x \partial_x s &= g\gamma \left(L \partial_x s + (|\psi(t, x)|^2 - 1) L \partial_x s + (\psi \partial_x \bar{\psi} + \bar{\psi} \partial_x \psi) L s \right. \\ & \quad \left. + e^{-\zeta t} \partial_x (|\psi|^2 \frac{Q(s, s)}{P}) + \frac{\zeta}{g\gamma^2} \partial_x s \right), \\ \partial_x s(0, x, p) &= \partial_x \tilde{R}_i(x, p). \end{aligned}$$

Analogously, using Lemma 3.5,

$$\begin{aligned} & \sup_{t \in [0, T_1]} \| \partial_x s(t) \|_{2,2} + \frac{3}{4} \sqrt{\gamma} \| \nu^{\frac{1}{2}} \partial_x s \|_{T_1, 2, 2} < \beta_1 \left(\| \partial_x \tilde{R}_i \|_{2,2} + \beta_6 c_5 \gamma^{\frac{3}{2}} \eta \| \nu^{\frac{1}{2}} \partial_x s \|_{T_1, 2, 2} \right. \\ & \quad \left. + 2c_4 \sqrt{\gamma} \| (\partial_x \psi) \nu^{-\frac{1}{2}} L s \|_{T_1, 2, 2} + \sqrt{\gamma} \| \nu^{-\frac{1}{2}} \partial_x (|\psi|^2 \frac{Q(s, s)}{P}) \|_{T_1, 2, 2} \right). \end{aligned}$$

Here,

$$\begin{aligned} \| (\partial_x \psi) \nu^{-\frac{1}{2}} L s \|_{T_1, 2, 2}^2 &\leq \beta_6^2 \int_0^{T_1} \int |\partial_x \psi|^2(t, x) \frac{P}{1+P} \nu(p) s^2(t, x, p) dp dx dt \\ &\leq \beta_6^2 \left(\sup_{t \in [0, T_1]} \int |\partial_x \psi|^2(t, x) dx \right) \left(\int_0^{T_1} \int \frac{P}{1+P} \nu(p) \sup_{x \in [0, 2\pi]} s^2(t, x, p) dp dt \right) \\ &\leq 2\beta_6^2 \gamma^2 \eta^2 \left(\int_0^{T_1} \int \frac{P}{1+P} \nu(p) \sup_{x \in [0, 2\pi]} s^2(t, x, p) dp dt \right) \quad \text{by (4.47)} \\ &\leq 2\beta_4^2 \beta_6^2 \gamma^2 \eta^2 \left(\| \nu^{\frac{1}{2}} s \|_{T_1, 2, 2}^2 + \| \nu^{\frac{1}{2}} \partial_x s \|_{T_1, 2, 2}^2 \right). \end{aligned}$$

Moreover,

$$\begin{aligned}
& \left\| \nu^{-\frac{1}{2}} \partial_x (|\psi|^2 \frac{Q(s, s)}{P}) \right\|_{T_1, 2, 2}^2 \\
& \leq 4c_4^2 \left(\int_0^{T_1} \int |\partial_x \psi|^2(t, x) \nu^{-1} \left(\frac{Q(s, s)}{P} \right)^2(t, x, p) \frac{P}{1+P} dp dx dt \right. \\
& \quad \left. + c_4^2 \int_0^{T_1} \int \nu^{-1} \left(\frac{Q(\partial_x s, s)}{P} \right)^2(t, x, p) \frac{P}{1+P} dp dx dt \right) \\
& \leq 4\beta_8^2 c_4^2 \left(\frac{1}{\nu_0} \left(\sup_{t \in [0, T_1]} \int |\partial_x \psi|^2(t, x) dx \right) \left(\int_0^{T_1} \int \nu(p) \sup_{x \in [0, 2\pi]} s^2(t, x, p) \frac{P}{1+P} dp dt \right)^2 \right. \\
& \quad \left. + \beta_4^2 c_4^2 \|s\|_{\infty, 2, H^1}^2 \|\nu^{\frac{1}{2}} \partial_x s\|_{T_1, 2, 2}^2 \right) \\
& \leq 4\beta_4^2 \beta_8^2 c_4^2 \left(\frac{2}{\nu_0} \gamma^2 \eta^2 \|\nu^{\frac{1}{2}} s\|_{2, 2, H^1}^4 + c_4^2 \|s\|_{\infty, 2, H^1}^2 \|\nu^{\frac{1}{2}} \partial_x s\|_{T_1, 2, 2}^2 \right) \\
& \leq 8\beta_4^2 \beta_8^2 c_4^2 \left(\frac{2}{\nu_0} \gamma \eta^4 \|\nu^{\frac{1}{2}} s\|_{2, 2, H^1}^2 + c_4^2 \eta^2 \|\nu^{\frac{1}{2}} \partial_x s\|_{T_1, 2, 2}^2 \right)
\end{aligned}$$

by (4.47). It follows from the third bound on η_0 in (4.43) that

$$\left\| \partial_x s(t) \right\|_{2, 2} + \frac{\sqrt{\gamma}}{2} \left\| \nu^{\frac{1}{2}} \partial_x s \right\|_{T_1, 2, 2} \leq \beta_1 \left(8\beta_1 \beta_4 c_4 \left(\beta_6 + \frac{\beta_8}{\sqrt{\nu_0}} \right) \left\| \tilde{R}_i \right\|_{2, 2} + \left\| \partial_x \tilde{R}_i \right\|_{2, 2} \right).$$

Consequently,

$$\begin{aligned}
& \left\| s(t) \right\|_{2, 2} + \left\| \partial_x s(t) \right\|_{2, 2} + \sqrt{\gamma} \left(\left\| \nu^{\frac{1}{2}} s \right\|_{T_1, 2, 2} + \left\| \nu^{\frac{1}{2}} \partial_x s \right\|_{T_1, 2, 2} \right) \\
& \leq c_2 \left(\left\| \tilde{R}_i \right\|_{2, 2} + \left\| \partial_x \tilde{R}_i \right\|_{2, 2} \right), \tag{4.51}
\end{aligned}$$

with c_2 defined in (4.37). And so, using (4.45),

$$\left\| s(t) \right\|_{2, 2} + \left\| \partial_x s(t) \right\|_{2, 2} + \sqrt{\gamma} \left(\left\| \nu^{\frac{1}{2}} s \right\|_{T_1, 2, 2} + \left\| \nu^{\frac{1}{2}} \partial_x s \right\|_{T_1, 2, 2} \right) \leq \frac{\eta \zeta}{10(1+c_3)}, \quad t \in [0, T_1]. \tag{4.52}$$

Recalling that $\zeta \leq 1$, this improves the second inequality in (4.47).

Next consider α . To improve the first inequality in (4.47), each term in the right hand side of (4.32) is first controlled separately. By (4.48) and the Cauchy-Schwartz inequality,

$$\begin{aligned}
& |2ig\gamma \int (\bar{\psi} \partial_x \psi - \psi \partial_x \bar{\psi}) \int P \partial_x \tilde{R} dp dx - \frac{g}{2} \gamma^2 \int (\bar{\psi} \partial_x \psi + \psi \partial_x \bar{\psi}) \int (PL \partial_x \tilde{R} + \gamma \partial_x Q(\tilde{R}, \tilde{R})) dp dx| \\
& \leq gc_4 (2\sqrt{\beta_2} + \beta_5 \gamma) \gamma e^{-\zeta t} \int |\partial_x \psi(t, x)| \left(\int \frac{P}{1+P} |\partial_x s(t, x, p)|^2 dp \right)^{\frac{1}{2}} dx \\
& \quad + 2gc_4 \beta_7 \gamma^3 e^{-\zeta t} \int |\partial_x \psi(t, x)| \left(\int \frac{\nu P}{1+P} |s(t, x, p)|^2 dp \int \frac{P}{1+P} |\partial_x s(t, x, p)|^2 dp \right)^{\frac{1}{2}} dx \\
& \leq gc_4 \gamma e^{-\zeta t} \sqrt{\alpha(t)} \left\| \partial_x s(t) \right\|_{2, 2} \left(2\sqrt{\beta_2} + \beta_5 + \gamma^2 e^{-\zeta t} \sup_{x \in [0, 2\pi]} \left(\int \frac{\nu P}{1+P} s^2(t, x, p) dp \right)^{\frac{1}{2}} \right) \\
& \leq gc_4 \left(2\sqrt{\beta_2} + \beta_5 + \beta_4 \right) \gamma e^{-\zeta t} \sqrt{\alpha(t)} \left\| \partial_x s(t) \right\|_{2, 2} \left(1 + \gamma \left(\left\| \nu^{\frac{1}{2}} s(t) \right\|_{2, 2} + \left\| \nu^{\frac{1}{2}} \partial_x s(t) \right\|_{2, 2} \right) \right) \\
& = \frac{c_3}{\sqrt{2}} \gamma e^{-\zeta t} \sqrt{\alpha(t)} \left\| \partial_x s(t) \right\|_{2, 2} \left(1 + \gamma \left\| \nu^{\frac{1}{2}} s(t) \right\|_{2, H^1} \right), \tag{4.53}
\end{aligned}$$

by the definition (4.38) of c_3 . And so, by the α -part of (4.47) and (4.52),

$$\begin{aligned} & |2ig\gamma \int (\bar{\psi}\partial_x\psi - \psi\partial_x\bar{\psi}) \int P\partial_x\tilde{R}dpdx - \frac{g}{2}\gamma^2 \int (\bar{\psi}\partial_x\psi + \psi\partial_x\bar{\psi}) \int (PL\partial_x\tilde{R} + \gamma\partial_xQ(\tilde{R}, \tilde{R}))dpdx| \\ & \leq \zeta e^{-\zeta t} \frac{\gamma^2\eta^2}{10} (1 + \gamma \| \nu^{\frac{1}{2}}s(t) \|_{2,H^1}). \end{aligned} \quad (4.54)$$

By (4.4), the α -part of (4.47) and (4.52),

$$\begin{aligned} | \int |\partial_x\psi|^2(t, x) \int PL\tilde{R}dpdx | & \leq \beta_5 e^{-\zeta t} \alpha(t) \sup_{x \in [0, 2\pi]} \left(\int s^2(t, x, p) \frac{P}{1+P} dp \right)^{\frac{1}{2}} \\ & \leq 2\beta_4\beta_5\gamma^2\eta^2 e^{-\zeta t} \| s(t) \|_{2,H^1} \\ & \leq \beta_4\beta_5\gamma^2\eta^3\zeta e^{-\zeta t}, \end{aligned} \quad (4.55)$$

and, by (4.6), (4.52) and (4.47),

$$\begin{aligned} & | \int |\partial_x\psi|^2(t, x) \int Q(\tilde{R}, \tilde{R})dpdx | \\ & \leq \beta_7 e^{-2\zeta t} \alpha(t) \sup_{x \in [0, 2\pi]} \left(\int s^2(t, x, p) \frac{P}{1+P} dp \right)^{\frac{1}{2}} \sup_{x \in [0, 2\pi]} \left(\int \nu(p) s^2(t, x, p) \frac{P}{1+P} dp \right)^{\frac{1}{2}} \\ & \leq \beta_4^2\beta_7 e^{-2\zeta t} \alpha(t) \| s \|_{\infty, 2, H^1} \| \nu^{\frac{1}{2}}s(t) \|_{2, H^1} \\ & \leq \beta_4^2\beta_7\gamma^2\eta^3\zeta e^{-2\zeta t} \| \nu^{\frac{1}{2}}s(t) \|_{2, H^1}. \end{aligned} \quad (4.56)$$

The $|\psi|^2(|\psi|^2 - 1)$ factor of the integrand in the last term of the r.h.s. in (4.32) is split into $|\psi|^2(|\psi|^2 - 1) = (|\psi|^2 - 1)^2 + (|\psi|^2 - 1)$. It gives rise to the terms

$$\int (|\psi|^2 - 1)^2 \int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R}))dpdx \quad \text{and} \quad \int (|\psi|^2 - 1) \int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R}))dpdx.$$

Analogously to the previous control of $\int |\partial_x\psi|^2 \int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R}))dpdx$, it holds that

$$| \int (|\psi|^2 - 1)^2 \int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R}))dpdx | \leq \frac{2\beta_4}{g} \gamma^2 \eta^3 \zeta e^{-\zeta t} (\beta_5 + \beta_4\beta_7\gamma e^{-\zeta t} \| \nu^{\frac{1}{2}}s(t) \|_{2, H^1}).$$

Moreover,

$$\begin{aligned} & | \int (|\psi|^2 - 1) \int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R}))dpdx | \\ & \leq \sqrt{\alpha(t)} \left(\int \left(\int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R}))dp \right)^2 dx \right)^{\frac{1}{2}} \\ & \leq 2\sqrt{\frac{\alpha(t)}{g}} e^{-\zeta t} (\beta_5 \| s(t) \|_{2,2} + \beta_4\beta_7\gamma e^{-\zeta t} \| s \|_{\infty, 2, H^1} \| \nu^{\frac{1}{2}}s(t) \|_{2,2}) \\ & \leq \frac{1}{\sqrt{g}} \gamma \eta^2 \zeta e^{-\zeta t} (\beta_5 + \beta_4\beta_7\gamma e^{-\zeta t} \| \nu^{\frac{1}{2}}s(t) \|_{2,2}). \end{aligned}$$

Consequently,

$$\begin{aligned} & g^2\gamma^2 | \int |\psi|^2 (|\psi|^2 - 1) \int (PL\tilde{R} + \gamma Q(\tilde{R}, \tilde{R}))dpdx | \\ & \leq g(2\beta_4 + \sqrt{g})\gamma^3\eta^2\zeta e^{-\zeta t} (\beta_5 + \beta_4\beta_7\gamma e^{-\zeta t} \| \nu^{\frac{1}{2}}s(t) \|_{2, H^1}). \end{aligned} \quad (4.57)$$

And so, it follows from (4.32), (4.54), (4.55), (4.56) and (4.57) that

$$|\alpha'(t)| \leq \left(\frac{1}{10} + c_6\gamma \right) \gamma^2 \eta^2 \zeta e^{-\zeta t} + c_7 \gamma^3 \eta^2 \zeta e^{-\zeta t} \|\nu^{\frac{1}{2}} s(t)\|_{2,H^1}, \quad t \in [0, T_1), \quad (4.58)$$

with c_6 (resp. c_7) defined in (4.41) (resp. (4.42)). Integrating (4.58) on $[0, t]$ for $t \in [0, T_1)$, using the Cauchy-Schwartz inequality in the last term, (4.52) and the two last inequalities in (4.44) gives

$$\begin{aligned} \alpha(t) &\leq \alpha(0) + \left(\frac{1}{10} + c_6\gamma \right) \gamma^2 \eta^2 + c_7 \gamma^3 \eta^2 \|\nu^{\frac{1}{2}} s\|_{2,2,H^1} \\ &\leq \alpha(0) + \left(\frac{1}{10} + c_6\gamma + c_7 \sqrt{\gamma} \eta \zeta \right) \gamma^2 \eta^2 \\ &\leq \alpha(0) + \frac{\gamma^2 \eta^2}{2}, \quad t \in [0, T_1), \end{aligned}$$

so that, by (4.46),

$$\alpha(t) < \frac{3}{2} \gamma^2 \eta^2, \quad t \in [0, T_1). \quad (4.59)$$

Since the bounds obtained in the r.h.s. of (4.52) and (4.59) are better than the ones defining T_1 as the maximal time so that (4.47) holds, it implies that $T_1 = T_0$.

Let T_2 be the maximal time such that the solution exists on $[0, T_2[$ and (4.47) holds.

The family $(\tilde{\Phi}(t))_{t \in [0, T_2[}$ is bounded in $H^1([0, 2\pi])$. Indeed, the α -part of (4.47) implies that $\int_0^{2\pi} |\partial_x \tilde{\Phi}(t, x)|^2 dx \leq 2$. Moreover, (4.49) implies that

$$\gamma(\mathcal{R}\tilde{\Phi})^2 + 2\mathcal{R}\tilde{\Phi} + \gamma(\mathcal{I}\tilde{\Phi})^2 - c_5\eta \leq 0,$$

so that

$$\gamma|\tilde{\Phi}(t, x)| \leq 3 + c_5, \quad (t, x) \in [0, T_2[\times [0, 2\pi].$$

Moreover, it follows from (4.47) that $(\tilde{R}(t))_{t \in [0, T_2[}$ is bounded by 2 in $H^1_{\frac{P}{1+P}}([0, 2\pi] \times \mathbb{R}^3)$.

Consequently, Proposition 4.1 applies with any time $t < T_2$ as initial time and provides a unique solution to (4.34)-(4.35) on an interval of time of length \tilde{T}_0 from Proposition 4.1 when the initial data are bounded by $\eta_0 = 3\left(\frac{3+c_5}{\gamma} + 1\right)$. If T_2 is finite, using Proposition 4.1 with initial time $T_2 - \frac{\tilde{T}_0}{2}$ and arguing as for (4.52), (4.59), it follows that the solution can be continued beyond T_2 up to $T_2 + \frac{\tilde{T}_0}{2}$, so that (4.47) holds. This contradicts T_2 being the maximal time. It results that $T_2 = +\infty$.

Set $\lambda = \gamma^2$ and write $\tilde{R}_i = \gamma R_i$, $\tilde{R} = \gamma R$, $\tilde{\Phi}_i = \gamma \Phi_i$, and $\tilde{\Phi} = \gamma \Phi$.

The existence part of Theorem 1.1 is thus proved for λ_1 in the statement of Theorem 1.1 given by γ_0^2 with γ_0 defined in (4.15)-(4.44), c_ζ given by (4.36), and $\eta_1 = \frac{\eta}{10c_1(1+c_2)(1+c_3)}$ from (4.45) with η smaller than η_0 given by (4.43).

It follows from (4.51) that R and $\partial_x R$ converge exponentially to zero of order ζ . As a consequence using the total mass conservation, $\int |\psi|^2 dx$ converges exponentially to $2\pi n_0$. Using (4.58), it follows that $\alpha_\infty := \lim_{t \rightarrow \infty} \alpha(t)$ exists, and is finite with the convergence to the limit being exponential of order ζ . The solution f is positive. Namely, by (4.52) the magnitude of R is bounded by 1 in L^∞ , and so $|\lambda R| < 1$. \blacksquare

Bibliography

1. Allemand, T.: Derivation of a two-fluids model for a Bose gas from a quantum kinetic system, *Kin. Rel. Models* 2, 379-402 (2009)
2. Arkeryd, L., Nouri, A.: Bose condensates in interaction with excitations - a kinetic model, *Comm. Math. Phys.* 310, 765-788 (2012)
3. Arkeryd, L., Nouri, A.: A Milne problem from a Bose condensate with excitations, *Kin. Rel. Models* 6, 671-686 (2013)
4. Bose, S.N.: Planck's Law and Light Quantum Hypothesis, *Z. Phys.* 26, 178 (1924)
5. Eckern, U.: Relaxation processes in a condensate Bose gas, *J. Low Temp. Phys.* 54, 333-359 (1984)
6. Einstein, A.: *Sitzber. Kgl. Preuss. Akad. Wiss.* 261 (1924)
7. Escobedo, M., Velázquez, J.: Finite time blow-up and condensation for the bosonic Nordheim equation, *Invent. Math.* (2013) doi:10.1007/s00222-014-0539-7
8. Escobedo, M., Pezzotti, F., Valle, M.: Analytical approach to relaxation dynamics of condensed Bose gases, *Ann. Phys.* 326, 808-827 (2011)
9. Golse, F., Levermore, D.: Stokes-Fourier and acoustic limits for the Boltzmann equation; convergence proofs, *Comm. Pure Appl. Math.* 55, 336-393 (2002)
10. Griffin, A., Nikuni, T., Zaremba, E.: *Bose-condensed gases at finite temperatures*, Cambridge University Press, Cambridge 2009
11. Hohenberg, P., Martin, P.: Microscopic theory of superfluid helium, *Ann. Phys.* 34, 291-359 (1965)
12. Imamovic-Tomasovic, M., Kadanoff-Baym, L.: *Kinetic theory for a trapped Bose condensate gas*, Thesis, Univ. Toronto 2001
13. Imamovic-Tomasovic, M., Griffin, A.: Quasiparticle kinetic equation in a trapped Bose gas at low temperature, *J. Low Temp. Phys.* 122, 617-655 (2001)
14. Khalatnikov, I.M.: *Theory of Superfluidity* (in Russian), Nauka, Moskva (1971)
15. Kirkpatrick, T.R., Dorfman, J.R.: Transport in a dilute but condensed nonideal Bose gas: kinetic equations, *J. Low Temp. Phys.* 58, 301-331 (1985)
16. Kirkpatrick, T.R., Dorfman, J.R.: Transport coefficients in a dilute but condensed Bose gas, *J. Low Temp. Phys.* 58, 399-415 (1985)
17. Kane, J., Kadanoff, L.: Green's functions and superfluid hydrodynamics, *Jour. math. Phys.* 6, 1902-1912 (1965)

18. Lu, X.: The Boltzmann equation for Bose-Einstein particles: condensation in finite time, *J. Stat. Phys.* 150, 1138-1176 (2013)
19. Maslova, N.: Nonlinear evolution equations: kinetic approach, World Scientific, Singapore (1993)
20. Nordheim, L.W.: On the kinetic methods in the new statistics and its applications in the electron theory of conductivity, *Proc. Roy. Soc. London Ser. A* 119, 689-698 (1928)
21. Pitaevski, L., Stringari, S.: Bose-Einstein Condensation, Clarendon Press, Oxford (2003)
22. Pomeau, Y., Brachet, M-É., Métens, S., Rica, S.: Théorie cinétique d'un gaz de Bose dilué avec condensat, *CRAS 327 Série II b*, 791-798 (1999)
23. Spohn, H.: Kinetics of the Bose-Einstein condensation, *Physica D* 239, 627-634 (2010)
24. Stoof, H.: Coherent versus incoherent dynamics during Bose-Einstein condensation in atomic gases, *J. Low Temp. Phys.* 114, 11-108 (1999)
25. Uehling, E.A., Uhlenbeck, G.E.: Transport phenomena in Einstein-Bose and Fermi-Dirac gases, *Phys. Rev.* 43, 552-561 (1933)
26. Zaremba, E., Nikuni, T., Griffin, A.: Dynamics of trapped Bose gases at finite temperatures, *J. Low Temp. Phys.* 116, 277-345 (1999)