

# Split group codes

Cunsheng Ding, David R. Kohel, and San Ling

*Abstract*— We construct a class of codes of length  $n$  such that the minimum distance  $d$  outside of a certain subcode is, up to a constant factor, bounded below by the square root of  $n$ , a well-known property of quadratic residue codes. The construction, using the group algebra of an abelian group and a special partition or *splitting* of the group, yields quadratic residue codes, duadic codes, and their generalizations as special cases. We show that most of the special properties of these codes have analogues for split group codes, and present examples of new classes of codes obtained by this construction.

**Index terms:** Split group codes, quadratic residue codes, duadic codes.

## I. INTRODUCTION

The codes of study in this article unify elements common to quadratic residue codes, duadic codes, and their generalizations with a common construction. Binary duadic codes were introduced in Leon, Masley, and Pless [7] as a generalization of quadratic residue codes, and further studies in Pless, Masley, and Leon. [12]. The authors exploited features particular to  $\mathbb{F}_2$  in order to write down an idempotent generator defining the codes. Smid [17] (for summary see [18]) removed the base field restriction and brought the definition in line with the constructive definition for quadratic residue codes. Under this definition, the Q-codes of Pless [11] are duadic codes over  $\mathbb{F}_4$ . Further aspects of these codes can be found in Rushanan [15] and Pless [13], [14].

Previously quadratic residue codes had been generalized in another direction in Camion [3] and in Ward [20]. In the approach of Camion, developed further in van Lint and MacWilliams [9], the generalized quadratic residue codes are defined as ideals in abelian group algebras, a generalization of cyclic codes. Since most of the features of duadic codes also carry over to the abelian group algebras, Rushanan [16] defines duadic codes in this setting, but reverts to a nonconstructive idempotent definition.

In the present work we unify the various abelian group algebra constructions. By working with the dual group  $G$  of and abelian group  $A$ , we can view the group algebra as a ring of functions on  $A$ . As a generalization of the duadic construction we broaden the definition of partitions, or splittings, and show that the main theorems for duadic codes hold in this larger setting. In particular, Theorem IV.5, Theorem IV.10, and Theorem IV.11 are analogues of the main theorems for duadic codes, which hold in this general context as well. By example we show that new subclasses introduced here hold good codes.

The article is structured as follows. In Section II we introduce the abelian group rings and the family of split group codes which form the objects of study in this work.

The main body of this section is devoted to properties of the ideals and ideal codes in these rings. We follow in Section III with an investigation of the problem of determining the minimal subfield  $F$  over which a code can be defined and describe algorithmic aspects of computing with group algebra codes. Section IV treats duality, code extensions, and minimum distance bounds, and Section V gives explicit examples and computational results for select subclasses of split group codes.

## II. SPLIT GROUP CODES

Let  $R$  be a finite commutative ring and let  $A$  be its underlying abelian group. Then  $A$  is a finite abelian group, written additively, whose exponent and order we denote by  $m$  and  $n$ , respectively. Let  $K$  be a field containing all the  $m$ -th roots of unity and in which  $n$  is invertible. We write the group operation additively. Define  $G = \text{Hom}(A, K^*)$  to be the group of homomorphisms from  $A$  to  $K^*$ , or characters, and let  $K[G]$  be the group ring over  $K$ . Since  $G$  and  $A$  are isomorphic, albeit noncanonically, the ring  $K[G]$  is a commutative algebra of dimension  $n$  over  $K$ . For any character  $\psi$  in  $G$  we also denote by  $\psi$  its image in  $K[G]$ . By extending characters linearly, we interpret the elements of  $K[G]$  as functions from  $A$  to  $K$ .

### A. Abelian group rings and decompositions

The following form of the discrete Fourier transform provides the basis for the later study of ideals and ideal codes in the ring  $K[G]$ .

*Theorem II.1:* Evaluation at  $x$  defines a homomorphism  $K[G] \rightarrow K$  with kernel  $\mathfrak{m}_x = \{f \in K[G] \mid f(x) = 0\}$ , such that the map

$$\epsilon : K[G] \longrightarrow K^A = \prod_{x \in A} K$$

$$f \longmapsto (f(x))_{x \in A}$$

is an isomorphism of rings with inverse defined by

$$\lambda : K^A \longrightarrow K[G].$$

$$(c_x)_{x \in A} \longmapsto \frac{1}{n} \sum_{\psi \in G} \left( \sum_{x \in A} c_x \psi(x)^{-1} \right) \psi$$

The theorem is an immediate consequence of the orthogonality relations for characters (see Chapter VIII §4-§5 of Lang [6]). In the special case  $A = \mathbb{Z}/n\mathbb{Z}$ , we have an isomorphism  $K[G] \cong K[X]/(X^n - 1)$  where  $X$  is the image of a generator for  $G$ . For a primitive  $n$ -th root of unity  $\zeta$ , we suppose, under the isomorphism with  $K[G]$ , that  $X$  acts on  $A$  by  $X(r) = \zeta^r$ . Then the evaluation map  $\epsilon$  takes the usual form

$$f(x) \longmapsto (f(1), f(\zeta), \dots, f(\zeta^{n-1})),$$

of the discrete Fourier transform (see for instance §4.3.3 C of Knuth [5]).

An *idempotent* of a ring  $S$  is a nonzero element  $e$  such that  $e^2 = e$ , and is called *primitive* if for every other idempotent  $f$ , either  $ef = e$  or  $ef = 0$ . The primitive idempotents in  $K^A$  are clearly the elements  $\delta_x = (0, \dots, 1, 0, \dots, 0)$ , having a 1 in the  $x$ -position. Under the isomorphism of rings, we obtain the following form for the primitive idempotents of  $K[G]$ .

*Corollary II.2:* The primitive idempotents of  $K[G]$  are given by

$$e_x = \frac{1}{n} \sum_{\psi \in G} \psi(x)^{-1} \psi, \quad (1)$$

one for each  $x$  in  $A$ , and the maximal ideal  $\mathfrak{m}_x$  is generated by  $1 - e_x$ .

We return later to the role of idempotents in decompositions of rings and their ideal codes.

### B. Splittings and codes

For an element  $s$  of  $R$  we denote the corresponding endomorphism  $x \mapsto sx$  of  $A$  by  $\tau_s$ . We denote the induced *pull-back* map on  $G$  and on  $K[G]$  by  $\mu_s$ , defined as  $\mu_s(f) = f \circ \tau_s$ , so that

$$\mu_s(f)(x) = f(sx)$$

for all  $x \in A$  and all  $f \in K[G]$ . We define a *splitting* of  $A$  over  $Z$  to be a triple  $(Z, X_0, X_1)$  giving a partition  $A = Z \cup X_0 \cup X_1$  for which there exists an element  $s$  in the unit group  $R^*$  of  $R$  with  $\tau_s(X_0) = X_1$  and  $\tau_s(X_1) = X_0$ . We say that such an  $s$  *splits*  $(Z, X_0, X_1)$  and we say that an element  $s$  in  $R^*$  such that  $\tau_s(X_0) = X_0$  and  $\tau_s(X_1) = X_1$  *stabilizes* the splitting. This definition generalizes the splittings considered in Leon, Masley, and Pless [7].

For any subset  $X$  of  $A$  we define an ideal

$$I_X = \{f \in K[G] \mid f(x) = 0 \text{ for all } x \in X\}. \quad (2)$$

The *split group code*  $C_0(K)$  over  $K$  associated to a splitting  $(Z, X_0, X_1)$  is defined to be the ideal  $C_0(K) = I_{X_0}$  and the *conjugate split group code* to be  $C_1(K) = I_{X_1}$ . In a like manner, we define the subcodes  $C_0^Z(K) = I_{Z \cup X_0}$ ,  $C_1^Z(K) = I_{Z \cup X_1}$  and  $C_Z(K) = I_{X_0 \cup X_1}$ .

The code  $C_0$  is said to be *split* by a unit  $s$  of  $R$  if  $\mu_s(C_0) = C_1$  and  $\mu_s(C_1) = C_0$  and *stabilized* by  $s$  if  $\mu_s$  stabilizes  $C_0$  and  $C_1$ . One verifies that  $\mu_s$  acts on the set of maximal ideals by sending  $\mathfrak{m}_x$  to  $\mathfrak{m}_{s^{-1}x}$ , from which we obtain the following property of splittings.

*Proposition II.3:* A split group code  $C_0$  is split or stabilized by  $s$  if and only if  $s$  splits or stabilizes  $(Z, X_0, X_1)$ , respectively.

### C. Idempotent decompositions

In this section we indicate how Theorem II.1 gives the idempotent decomposition of  $K[G]$ . The following formulation gives this decomposition in terms of the primitive idempotents.

*Proposition II.4:* The ring  $K[G]$  decomposes as a direct sum  $\bigoplus_{x \in A} Ke_x$  such that  $fe_x = f(x)e_x$ , so that  $f$  has the form

$$f = \sum_{x \in A} f(x)e_x.$$

Every idempotent  $e$  in  $K[G]$  can be uniquely written in the form

$$e = \sum_{x \in X} e_x,$$

for a nonempty subset  $X$  of  $A$ .

*Proof:* By construction of  $e_x$ , under the ring isomorphism  $K[G] \cong K^A$ , the ideal  $Ke_x$  in  $K[G]$  corresponds to the  $x$ -component in  $K^A$  with the image of  $e_x$  equal to the unity in that component. By definition of the isomorphism, the coefficient of  $e_x$  in an element  $f$  of  $K[G]$  is  $f(x)$ . By definition an idempotent  $e$  satisfies  $e^2 = e$ , so is either zero or unity in each component, hence can be written as a sum of the primitive idempotents as indicated.  $\square$

*Corollary II.5:* All nonzero ideals of  $K[G]$  are of the form

$$I_X = \bigoplus_{x \in X^c} Ke_x,$$

generated by the idempotent  $e = \sum_{x \in X^c} e_x$ , for a unique proper subset  $X$  of  $A$ .

*Proof:* Let  $I$  be a nonzero ideal of  $K[G]$  and set

$$X = \{x \in A \mid f(x) = 0 \text{ for all } f \text{ in } I\}.$$

From Proposition II.4 it is clear that if  $x$  is in  $X^c$  then  $I$  contains  $Ie_x = Ke_x$  and that conversely  $I \cap Ke_x = Ie_x = (0)$  for all  $x$  in  $X$ . Thus  $I$  is the ideal  $I_X$ , having the indicated form. Representing  $I$  as the product  $\prod_{x \in X} \mathfrak{m}_x$  (see for instance Proposition 1.10 of Atiyah and MacDonald [1]), Corollary II.2 implies that  $I$  is generated by the idempotent  $e = \prod_{x \in X} (1 - e_x)$ . Expanding  $e$  as a product, we find

$$e = \prod_{x \in X} (1 - e_x) = 1 - \sum_{x \in X} e_x = \sum_{x \in X^c} e_x,$$

proving the form of the generator.  $\square$

*Theorem II.6:* Let  $(Z, X_0, X_1)$  be a splitting and let  $C_0(K)$  be the associated split group code. Then the following results hold.

1. The codes  $C_0(K)$  and  $C_1(K)$  are generated by the idempotents

$$e = \sum_{x \in X_0^c} e_x \text{ and } f = \sum_{x \in X_1^c} e_x.$$

Likewise the codes  $C_0^Z(K)$ ,  $C_1^Z(K)$ , and  $C_Z(K)$  are generated by

$$\sum_{x \in X_1} e_x, \sum_{x \in X_0} e_x, \text{ and } \sum_{z \in Z} e_z.$$

2. If the splitting is given by  $s$ , then  $\mu_s$  induces an equivalence of  $C_0(K)$  with its conjugate  $C_1(K)$ , and of the subcode  $C_0^Z(K)$  with  $C_1^Z(K)$ .

3.  $K[G]$  decomposes as a direct sum  $C_Z(K) \oplus C_0^Z(K) \oplus C_1^Z(K)$ .

*Proof:* The first statement follows from Corollary II.5 and the respective definitions of the codes. The second statement then follows by noting that  $\mu_s(e_x) = e_{s^{-1}x}$ ,

which implies that  $\mu_s$  exchanges the idempotents  $e$  and  $f$ . Since  $\mu_s$  also permutes the code basis  $G$ , this is an equivalence of codes. The decomposition of  $K[G]$  follows by Proposition II.4 and the grouping

$$K[G] = \left( \bigoplus_{x \in Z} Ke_x \right) \oplus \left( \bigoplus_{x \in X_1} Ke_x \right) \oplus \left( \bigoplus_{x \in X_0} Ke_x \right),$$

according to the partition  $A = Z \cup X_0 \cup X_1$ .  $\square$

The next corollary is an elementary consequence of the theorem.

*Corollary II.7:* The codes  $C_0(K)$  and  $C_1(K)$  have dimension  $(n+|Z|)/2$ ; the subcodes  $C_0^Z(K)$  and  $C_1^Z(K)$  have dimension  $(n-|Z|)/2$ ; and the dimension of  $C_Z(K)$  is  $|Z|$ .

### III. DESCENDING THE BASE FIELD

The field  $K$  was defined to contain the  $m$ -th roots of unity. However, we generally want to define codes over fields on which we place no such requirement. Indeed the principal field of interest is  $\mathbb{F}_2$ , which contains no nontrivial roots of unity at all!

In this section we let  $F$  be a subfield of  $K$  of  $q$  elements, and extend the definition of split group codes to  $F$ . The main goal of the section is to give necessary and sufficient conditions for split group codes to be defined over  $F$ , and to describe constructive methods for producing such codes.

For any vector subspace  $V = V(K)$  in  $K^n$  we define the descent problem as follows. Define  $V(F) = V \cap F^n$ , and note that

$$\dim_F(V(F)) \leq \dim_K(V(K)).$$

If equality holds we say that  $V$  is *defined over the field  $F$* . For fixed field  $F$ , over which the vector space  $C_0(K)$  of  $K^n = K[G]$  is defined, we write  $C_0 = C_0(F)$ , and refer to  $C_0$  as the *split group code* over  $F$ . Similarly we write  $C_1$ ,  $C_0^Z$ ,  $C_1^Z$ , and  $C_Z$  for the subcodes in  $F^n = F[G]$  defined over  $F$ .

#### A. Defining fields of codes

The following split group code provides an example for the descent problem, namely, reducing to a minimal field  $F$  over which a code is defined. Implicit in this example and the following one, is the principle that the vector space of an ideal is defined over  $F$  if and only if it contains a generator in  $F[G]$ . The role of  $\langle \tau_q \rangle$ -orbit decompositions should also be apparent to the reader familiar with the cyclotomic coset decompositions and cyclic codes, but we leave the proofs of these result for the next section.

*Example III.1:* Let  $F = \mathbb{F}_2$  and set  $R = \mathbb{Z}/15\mathbb{Z}$ . The 15th cyclotomic polynomial has a factor  $p(X) = X^4 + X + 1$  over  $\mathbb{F}_2$ . Setting  $K = \mathbb{F}_2[T]/(p(T))$ , the image of  $T$  is a 15th root of unity, which we denote  $\zeta$ .

Set  $Z = 3\mathbb{Z}/15\mathbb{Z} \cup 5\mathbb{Z}/15\mathbb{Z}$ . Its complement splits into  $\langle \tau_2 \rangle$ -orbits  $X_0 = \{1, 2, 4, 8\}$  and  $X_1 = \{7, 13, 14, 11\}$ , giving

a splitting  $(Z, X_0, X_1)$  by  $-1$ . The polynomials

$$g_0(X) = (X - \zeta)(X - \zeta^2)(X - \zeta^4)(X - \zeta^8)$$

$$= X^4 + X + 1,$$

$$g_1(X) = (X - \zeta^7)(X - \zeta^{13})(X - \zeta^{14})(X - \zeta^{11})$$

$$= X^4 + X^3 + 1.$$

are then the generator polynomials of split group codes  $C_0$  and  $C_1$  defined over  $\mathbb{F}_2$ . Raising  $g_i(X)$  in  $\mathbb{F}_2[X]/(X^{15} - 1)$  to the power  $2^4 - 1 = 15$  (see Proposition III.12), we obtain the idempotent generators

$$e_0 = X^{12} + X^9 + X^8 + X^6 + X^4 + X^3 + X^2 + X + 1,$$

$$e_1 = X^{14} + X^{13} + X^{12} + X^{11} + X^9$$

$$+ X^7 + X^6 + X^3 + 1,$$

over  $\mathbb{F}_2$  for the split group code  $C_0$  and its conjugate.  $\square$

The above shows how split group codes generalize the duadic codes of Leon, Masley, and Pless [7] – the latter being binary split group codes for splittings of  $A = \mathbb{Z}/n\mathbb{Z}$  over the set  $Z = \{0\}$  in the present terminology. Smid [17] extended the definition to arbitrary finite fields via generator polynomials, modelled on the standard one for quadratic residue codes. We summarize this correspondence in the present language as the following theorem. The proof is omitted, as it is a direct analogue of Theorem 6.9.3 of van Lint [10] for quadratic residue codes, and both theorem and proof can be extracted from the proof and discussion following Theorem 2 of Pless [13].

*Theorem III.2:* Let  $(\{0\}, S_0, S_1)$  be a splitting of  $\mathbb{Z}/n\mathbb{Z}$ , split by  $s$  in  $\mathbb{Z}/n\mathbb{Z}^*$  and stabilized by  $\tau_2$ . Then the element

$$e_0 = ((n+1)/2) + \sum_{a \in S_0} X^a \in \mathbb{F}_2[X]/(X^n - 1)$$

is an idempotent. Let  $G$  be the dual group of  $\mathbb{Z}/n\mathbb{Z}$  and fix a primitive  $n$ -th root of unity  $\zeta$ . Then the isomorphism of rings

$$\mathbb{F}_2[X]/(X^n - 1) \longrightarrow \mathbb{F}_2[G]$$

given by  $X \mapsto \chi$ , where  $\chi(a) = \zeta^a$ , maps the ideal generated by  $e_0$  to a split group code  $C_0$  defined with respect to a splitting  $(\{0\}, X_0, X_1)$  of  $\mathbb{Z}/n\mathbb{Z}$  by  $s$ .

*Remark III.3:* Note that the set  $X_0$  can be effectively recovered as

$$X_0 = \{a \in \mathbb{Z}/n\mathbb{Z} \mid e_0(\zeta^a) = 0\}.$$

Except for the quadratic residue splitting, the map sending  $S_0$  to  $X_0$  is generally not the identity for any choice of root of unity. Thus the idempotent construction of Theorem III.2, used as the definition of duadic codes in Leon et al. [7], gives an entirely different construction for binary cyclic duadic codes. As seen in Example III.1, this special construction does not generalize to binary cyclic split group codes. Moreover, definitions via idempotent relations as employed in Pless [11] and [12] are nonconstructive so are deduced here only as consequences of split group code constructions.  $\square$

To emphasize that not all codes covered by this work are binary, we conclude this section with a pair of duadic codes over the field  $\mathbb{F}_3$ . It has been noted [17] that there exist splittings of  $\mathbb{Z}/n\mathbb{Z}$  over  $\{0\}$  stabilized by  $\tau_q$  if and only

if  $q$  is a square modulo  $n$ . For binary codes, this implies that every prime divisor  $\ell$  of  $n$  is congruent to  $\pm 1 \pmod 8$  (Theorem 2 of Leon et al. [7]). For ternary codes, each  $\ell$  must be congruent to  $\pm 1 \pmod{12}$ . For  $n = 11$  there are only two orbits of  $\langle \tau_3 \rangle$  in  $\mathbb{Z}/11\mathbb{Z}^*$ , so the only ternary duadic code of this length is the  $[11, 6, 5]$ -quadratic residue code. The first new example occurs for block length 13.

*Example III.4:* Let  $F = \mathbb{F}_3$  let  $R = \mathbb{Z}/13\mathbb{Z}$ . Since 3 generates the biquadratic residues in  $R^*$ , there exist two nonequivalent splittings over  $Z = \{0\}$ . These give the quadratic residue code and a distinct duadic code respectively. First consider the quadratic residue code  $Q_0$  of length 13, a  $[13, 7, 5]$ -code over  $\mathbb{F}_3$ . Its subcode  $Q_0^Z$  of functions vanishing on  $Z$  is a  $[13, 6, 6]$ -code with weight enumerator polynomial

$$1 + 104X^6 + 78X^7 + 156X^8 + 130X^9 \\ + 156X^{10} + 78X^{11} + 26X^{12}.$$

Both codes are best possible for this length and their dimensions. For comparison, now consider the non-quadratic residue duadic code  $C_0$ . It is a  $[13, 7, 4]$ -code, but has subcode  $C_0^Z$  also of minimum distance 6. We find the weight enumerator polynomial to be

$$1 + 156X^6 + 494X^9 + 78X^{12},$$

so that the two subcodes  $Q_0^Z$  and  $C_0^Z$  are clearly nonequivalent codes with the same optimal minimum distance.  $\square$

### B. Descent by Galois action

Let  $\mathcal{G} = \text{Gal}(K/F)$  be the Galois group of the extension  $K/F$ , and let  $\sigma$  be the Frobenius automorphism  $c \mapsto c^q$  which generates  $\mathcal{G}$ . Then  $\mathcal{G}$  acts on  $K[G]$  by the natural action on the coefficients, with  $F[G]$  equal to the set of elements fixed under the action of  $\mathcal{G}$ . In this section we are interested in the action of this group on the collection of ideals in  $K[G]$  in order to determine those ideals which are defined, as vector spaces, over  $F$ .

By assumption the integer  $q$  is relatively prime to  $m$ , so that as an element of the finite ring  $R$ , it is invertible, and  $\tau_q$  is a well-defined automorphism of  $A$ . In this section we relate the action of  $\tau_q$  to the Galois group  $\mathcal{G}$ . This lets us reduce the study of the Galois action on ideals in  $K[G]$  to the action of the group  $\langle \tau_q \rangle$  on subsets in  $A$ . The action of the Frobenius automorphism is described by the following elementary lemma.

*Lemma III.5:* The Frobenius automorphism  $\sigma$  acts on the primitive idempotents of  $K[G]$  by  $e_x^\sigma = e_{qx}$  and similarly on the maximal ideals by  $\mathfrak{m}_x^\sigma = \mathfrak{m}_{qx}$ .

*Proposition III.6:* The idempotents of  $F[G]$  are those  $e$  in  $K[G]$  of the form

$$e = \sum_{x \in Y} e_x,$$

for which  $Y$  can be written as a union of orbits of  $\langle \tau_q \rangle$  in  $A$ , and  $e$  is primitive if and only if  $Y = \langle \tau_q \rangle x$  for some  $x$  in  $A$ . Let  $\{e_1, \dots, e_r\}$  be the set of primitive idempotents of  $F[G]$  with corresponding orbits  $\{Y_1, \dots, Y_r\}$ . Then in the

local decomposition

$$F[G] = \bigoplus_{i=1}^r F[G]e_i,$$

each  $F_i = F[G]e_i$  is a field extension of degree  $|Y_i|$  over  $F$ ,  $e_i \cong F$ .

*Proof:* The form of the idempotents of  $K[G]$  follows from Proposition II.4. The idempotents of  $F[G]$  are then the idempotents of  $K[G]$  which are invariant under the Galois group  $\mathcal{G}$ . By Lemma III.5 these must be the idempotents for which the index set  $Y$  is invariant under  $\tau_q$ , that is,  $Y$  decomposes into a union of orbits of the group  $\langle \tau_q \rangle$ .

To show that  $F[G]$  is isomorphic to a product of fields is to show that each local Artin factor  $F_i$  contains no nilpotents. But this is clear since  $K[G]$  is a product of fields, hence contains no nilpotents.

To prove the statement about the degree of the extension  $F_i/F$ , it suffices to show that  $\dim_F(F_i) = |Y_i|$ . Equivalently, we show that  $K[G]e_i = F_i \otimes_F K$  has dimension  $|Y_i|$  over  $K$ . But  $K[G]$  decomposes over  $K$  into one dimensional factors, so we have:

$$\dim_K(K[G]e_i) = \dim_K\left(\bigoplus_{x \in Y_i} Ke_x\right) = |Y_i|,$$

and the statement follows.  $\square$

*Corollary III.7:* Every nonzero ideal  $I$  of  $F[G]$  is of the form  $\bigoplus_{i \in T} F[G]e_i$  and generated by the idempotent  $e = \sum_{i \in T} e_i$ , where  $\{e_1, \dots, e_r\}$  is the set of primitive idempotents and  $T$  is a nonempty subset of  $\{1, \dots, r\}$ . Moreover there exists a unique ideal  $J$  in  $F[G]$  such that  $I \oplus J = F[G]$ .

*Proof:* Let  $I$  be an ideal of  $F[G]$  and set

$$T = \{i \in \{1, \dots, r\} \mid fe_i \neq 0 \text{ for some } f \text{ in } I\}.$$

Then for all  $i$  in  $\{1, \dots, r\}$  and all  $f$  in  $I$  such that  $fe_i$  is nonzero,  $F_i = F_i f$  is contained in  $I$  and  $Ie_j = \{0\}$  for  $j$  not in  $T$ . Thus  $I = \bigoplus_{i \in T} F[G]e_i$ , which is generated by the idempotent  $e = \sum_{i \in T} e_i$ . By Proposition III.6 it is clear that the ideal  $J = \bigoplus_{i \in T^c} F[G]e_i$ , where  $T^c$  is the complement of  $T$  in  $\{1, \dots, r\}$ , is the unique ideal complementing  $I$  in  $F[G]$ .  $\square$

*Theorem III.8:* Let  $I$  be an ideal in  $K[G]$ . Then the following conditions are equivalent.

1. The ideal  $I$  is defined over  $F$ .
2. The set  $X = \{x \in A \mid f(x) = 0 \text{ for all } f \text{ in } I\}$  is a union of  $\langle \tau_q \rangle$ -orbits.
3. The idempotent of  $I$  lies in  $F[G]$ .

Moreover there exists a unique minimal subfield of  $K$  over which  $I$  is defined.

*Proof:* Suppose that  $I$  is defined over  $F$ . By Corollary II.5 we have  $I = I_X$ , where  $X$  is as defined in the theorem. Since  $I$  has a basis in  $F[G]$ , it must be stabilized by the Galois group. By Lemma III.5 we then have

$$I_X = I_X^\sigma = \bigoplus_{x \in X^c} Ke_{qx} = \bigoplus_{x \in qX^c} Ke_x.$$

Therefore  $X^c = qX^c$  and so also  $X = qX$ . Suppose now that  $X$  is stabilized by  $\tau_q$ , and let  $e$  be the idempotent of

I. Then also by Lemma III.5,

$$e^\sigma = \sum_{x \in X^c} e_{qx} = \sum_{x \in X^c} e_x = e,$$

so  $e$  lies in  $F[G]$ . To complete the cycle, assume that  $I = K[G]e$  for  $e$  in  $F[G]$ . Then since  $F[G]e$  is contained in  $I(F) = F[G] \cap I$ , we have

$$\dim_F(I(F)) \geq \dim(F[G]e).$$

But  $K[G]e = K \otimes_F F[G]e$ , so the dimensions of vector spaces are preserved:

$$\dim_F(F[G]e) = \dim_K(K[G]e) = \dim_K(I).$$

Thus by definition  $I$  is defined over  $F$ .  $\square$

*Corollary III.9:* If  $C_0(K)$  is defined over  $F$  then so is  $C_1(K)$ , and  $F[G]$  has the decomposition

$$F[G] = C_Z(F) \oplus C_0^Z(F) \oplus C_1^Z(F).$$

If  $s$  gives the splitting, then  $\mu_s$  determines an equivalence of  $C_0$  with  $C_1$  and of  $C_0^Z$  with  $C_1^Z$ .

*Proof:* Since  $q$  lies in the center of  $R$ , the automorphisms  $\tau_s$  and  $\tau_q$  commute, hence  $\tau_s$  permutes the  $\langle \tau_q \rangle$ -orbits. In particular, as the image of  $X_0$ , the set  $X_1$  must also be the union of  $\langle \tau_q \rangle$ -orbits, hence  $C_1(K)$  is defined over  $F$ . By Corollary III.7, the ideals  $C_1^Z(F)$  and  $C_0^Z(F)$ , as the complementary ideals to  $C_0(F)$  and  $C_1(F)$ , are defined over  $F$ . Moreover, either as the complement to  $C_0^Z(F) \oplus C_1^Z(F)$ , or as the intersection of  $C_0(F)$  and  $C_1(F)$ , we find that  $C_Z(F)$  is defined over  $F$ . the decomposition of  $F[G]$  and the equivalence of codes follow from the corresponding results of Theorem II.6 over  $K$ .  $\square$

*Theorem III.10:* The block length, dimension, and minimum distance are well-defined invariants of  $C_0$ , independent of the field  $F$  over which  $C_0$  is defined.

*Proof:* The block length and dimension are invariant by definition. Let  $F$  be the minimal field of definition for  $C_0$ . Consider an extension  $L/F$ , and let  $\sigma$  be the Frobenius automorphism. For any element  $g = \sum_{\psi \in G} a_\psi \psi$  in  $C_0(L)$  of minimum nonzero weight, we choose  $\phi$  in  $\text{Supp}(g)$ . Then  $\text{Supp}(g) = \text{Supp}(g^\sigma)$  and  $\text{Supp}(a_\phi^\sigma g - a_\phi g^\sigma)$  is a proper subset, since it does not contain  $\phi$ . Since  $g$  has minimal nonzero weight in  $C_0(L)$ , we must have  $a_\phi^\sigma g - a_\phi g^\sigma = 0$ . Setting  $h = g/a_\phi = g^\sigma/a_\phi^\sigma$ , it follows that  $h$  is defined over  $F$ , so the minimum distance of  $C_0(F)$  equals that of  $C_0(L)$ .  $\square$

We note that the proof makes no use of special properties of the code  $C_0$ ; in fact the theorem holds for any linear code.

### C. Explicit constructions

Every finite abelian group  $A$  of exponent  $m$  is isomorphic to a unique product of the form

$$\mathbb{Z}/m_1\mathbb{Z} \times \cdots \times \mathbb{Z}/m_r\mathbb{Z},$$

where  $m_i$  divides  $m_{i+1}$  and  $m_r = m$ . We set  $M$  equal to  $\text{Hom}(A, \mathbb{Z}/m\mathbb{Z})$ . Then for such a decomposition of  $A$ , we choose generators  $x_1, \dots, x_r$  of the factors, and take  $\pi_1, \dots, \pi_r$  in  $M$  defined by

$$\pi_i(x_j) = \begin{cases} n_i & \text{if } j = i, \\ 0 & \text{otherwise,} \end{cases}$$

where  $n_i m_i = m$ . One readily verifies that  $\{\pi_1, \dots, \pi_r\}$  is a basis for  $M$ .

Suppose  $\chi$  is a fixed primitive character of  $\mathbb{Z}/m\mathbb{Z}$ . For any  $v$  in  $M$  we define  $\chi^v = \chi \circ v$ , and in particular set  $\chi_i = \chi^{\pi_i}$  for  $1 \leq i \leq r$ . Then  $\chi_1, \dots, \chi_r$  generate  $G$  and we fix an isomorphism

$$F[G] = F[\chi_1, \dots, \chi_r] \cong \frac{F[X_1, \dots, X_r]}{(X_1^{m_1} - 1, \dots, X_r^{m_r} - 1)},$$

sending  $\chi_i$  to  $X_i$ . Any  $v$  in  $M$  can be written as  $c_1\pi_1 + \cdots + c_r\pi_r$ , so  $\chi^v = \chi_1^{c_1} \cdots \chi_r^{c_r}$  which is represented by the monomial  $X_1^{c_1} \cdots X_r^{c_r}$  under the above isomorphism. We denote this element  $\mathbf{X}^v$  when we want to think of it as a monomial in the quotient polynomial ring and as  $\chi^v$  when we view it as a function.

For any subset  $Y$  of  $A$  stabilized by  $\langle \tau_q \rangle$  we construct the ideal  $I_Y$  as follows. For  $Y = Y_1 \cup Y_2$ , we have  $I_Y = I_{Y_1} I_{Y_2}$ , so it suffices to determine the maximal ideals  $\mathfrak{m}_Y = I_Y$ . For  $Y = \langle \tau_q \rangle z$ , we define  $M_Y = \{v \in M \mid v(z) = 0\}$ , and let  $B_Y$  be a basis.

Let  $\delta$  be the largest divisor of  $m$  such that  $Y \subseteq \delta A$ . We find  $a_1, \dots, a_r \in \mathbb{Z}/m\mathbb{Z}$  such that  $\sum_{i=1}^r a_i \pi_i(z) = \delta$ , and define  $\pi = \sum_{i=1}^r a_i \pi_i$ . With this notation we can write down a collection of generators for  $\mathfrak{m}_Y$ .

*Proposition III.11:* The maximal ideal  $\mathfrak{m}_Y$  is generated by the set

$$\mathcal{S} = \{g^\pi = \prod_{y \in Y} (\mathbf{X}^\pi - \chi^\pi(y))\} \cup \{\mathbf{X}^v - 1 \mid v \in B_Y\}.$$

*Proof:* By definition, for any  $y$  in  $Y$  and  $v$  in  $B_Y$  we have

$$\mathbf{X}^v(y) = \chi(v(y)) = \chi(0) = 1.$$

Likewise it is clear that  $g^\pi(y) = 0$ , so  $\mathcal{S}$  is contained in  $\mathfrak{m}_Y$ . It suffices to show the converse: if  $y$  is a root of all functions in  $\mathcal{S}$ , then  $y$  lies in  $Y$ . Let  $\phi : \mathbb{Z}/m\mathbb{Z} \rightarrow A$  be a splitting of  $\pi$ , i.e.  $\phi : \mathbb{Z}/m\mathbb{Z} \rightarrow A$  so that  $\pi\phi$  is the identity on  $\mathbb{Z}/m\mathbb{Z}$ . Suppose that  $y$  is in  $A$  such that  $\chi^v(y) = 1$  for all  $v$  in  $B_Y$ . Then  $y$  lies in the subgroup  $\mathbb{Z}/m\mathbb{Z}z$  of  $\mathbb{Z}/m\mathbb{Z}\phi(1)$  generated by  $Y$ . Since  $Y$  is in bijection with  $\pi(Y)$ , in order to show that a root of  $g^\pi$  in  $\mathbb{Z}/m\mathbb{Z}\phi(1)$  actually lies in  $Y$ , we show that  $g = (g^\pi)^\phi$  in  $F[\chi]$  has roots only in  $\pi(Y)$ . But by definition  $g$  equals

$$\prod_{y \in Y} (\mathbf{X}^{\pi\phi} - \chi^{\pi\phi}(y)) = \prod_{y \in Y} (\chi - \chi^\pi(y)).$$

So the roots of  $g$  in  $\mathbb{Z}/m\mathbb{Z}$  are precisely those in  $\pi(Y)$ .  $\square$

A set of generators for an ideal  $I$  may be reduced to the single idempotent generator by the following proposition.

*Proposition III.12:* Let  $g$  be a nonzero element in  $F[G]$ , let  $d$  be the order of the group  $\langle \tau_q \rangle$ , and set  $\ell = q^d - 1$ . Then  $e = g^\ell$  is an idempotent. For any collection of generators  $g_1, \dots, g_t$  of an ideal  $I$  with corresponding idempotents  $e_1 = g_1^\ell, \dots, e_t = g_t^\ell$ , the element

$$e = \sum_{i=1}^t e_i \prod_{j < i} (1 - e_j),$$

is the idempotent generator for  $I$ .

*Proof:* By Proposition III.6, the group algebra  $F[G]$  is isomorphic to a product of field extensions of  $F$ , each of

degree dividing  $d$  over  $F$ . In particular  $e = g^\ell$  is congruent to one or zero in every quotient, hence is an idempotent. To prove the main statement, we note that a collection of elements  $f_1, \dots, f_t$  generates the ideal  $I = I_Y$  if and only if the intersection of the zero sets  $Y_i$  of  $f_i$  equals  $Y$ . It is clear that each  $e_i$  has the same zeros as the corresponding  $g_i$ . Suppose that  $e_1$  and  $e_2$  are idempotents whose zero sets are subsets  $Y_1$  and  $Y_2$  of  $A$ . Then it is immediately verified that  $e = e_1 + e_2 - e_1 e_2$  is an idempotent, and by evaluating at each  $x$  in  $A$ , we check that  $e$  has zero set  $Y_1 \cap Y_2$ . By induction on  $t$ , it follows that the idempotent of  $I$  has the indicated form.  $\square$

In the above construction we have omitted the issue of constructing the character  $\chi$ . The definition of  $\chi$  requires only that we construct an extension field of  $F$  with a prescribed  $n$ th root of unity. This reduces to a standard polynomial factorization problem over finite fields, as already seen in Example III.1. The following provides a complete worked example of the idempotent construction.

*Example III.13:* Let  $F = \mathbb{F}_2$  and set  $R = \mathbb{Z}/7\mathbb{Z} \times \mathbb{Z}/7\mathbb{Z}$ . Define  $K = F_2[\zeta]$  and  $\chi(1) = \zeta$  where  $\zeta$  satisfies the irreducible factor  $X^3 + X + 1$  of the 7th cyclotomic polynomial. If  $Y$  is the orbit generated by  $(1, 2)$ , then the group  $M_Y$  has basis  $\{v\}$  where  $v((x_1, x_2)) = x_1 + 3x_2$ . We can take  $\pi$  to be the homomorphism defined by  $\pi((x_1, x_2)) = x_1$ . By Proposition III.11 this gives generators  $X_1 X_2^3 + 1$  and  $X_1^3 + X_1 + 1$ , so we find a generating set of idempotents:

$$\begin{aligned} e_0 &= (X_1^3 + X_1 + 1)^7 = X_1^4 + X_1^2 + X_1, \\ e_1 &= (X_1 X_2^3 + 1)^7 = X_1^2 X_2^6 + X_1^4 X_2^5 + X_1^6 X_2^4 \\ &\quad + X_1 X_2^3 + X_1^3 X_2^2 + X_1^5 X_2. \end{aligned}$$

The idempotent generator  $e = e_0 + e_1 - e_0 e_1$  of  $\mathfrak{m}_Y$  is then:

$$\begin{aligned} &(X_1^6 + X_1^4 + X_1^3 + X_1^2)X_2^6 + (X_1^6 + X_1^5 + X_1^4 + X_1)X_2^5 \\ &+ (X_1^6 + X_1^3 + X_1 + 1)X_2^4 + (X_1^5 + X_1^3 + X_1^2 + X_1)X_2^3 \\ &+ (X_1^5 + X_1^4 + X_1^3 + 1)X_2^2 + (X_1^6 + X_1^5 + X_1^2 + 1)X_2 \\ &+ X_1^4 + X_1^2 + X_1. \end{aligned}$$

In terms of the basis  $\{1, X_1, \dots, X_1^6, X_2, \dots, X_1^6 X_2^6\}$ , the idempotent is the weight 27 vector  $(0110100101001110011100111010110100101001110011101)$ , and a subset of  $\{X_1^i X_2^j e \mid 0 \leq i < 7, 0 \leq j < 7\}$  gives a vector space basis for the ideal generated by  $e$ .  $\square$

#### IV. DUALITY, EXTENSIONS, AND MINIMUM DISTANCE

In this section we analyze the decomposition of split group codes into orthogonal ideals under certain conditions on the splittings. In IV-B we introduce extensions of split group codes, define an inner product, and find conditions for the extended codes to be self-dual. In general the extension need not exist; we require that the base field  $F$  contain sufficiently many roots of unity. In a typical situation  $Z$  is a subgroup of  $A$  and the condition is that the image of  $Z$  under all characters in  $G$  is contained in  $F$ . For the duadic codes  $Z = \{0\}$  and this condition is void. In the general case this extends the theory of duality for duadic codes to general split group codes.

#### A. Orthogonal decompositions

For  $f = \sum_{\psi \in G} a_\psi \psi$  in  $K[G]$  we define the *support* of  $f$  to be the set

$$\text{Supp}(f) = \{\psi \in G \mid a_\psi \neq 0\},$$

such that the *weight* of  $f$  is  $\|f\| = |\text{Supp}(f)|$ .

*Proposition IV.1:* The standard Euclidean inner product of  $f$  and  $g$  in  $F[G]$  is

$$\langle f, g \rangle = \frac{1}{n} \sum_{x \in A} f(x)g(-x).$$

*Proof:* Let  $f = \sum_{\psi} a_\psi \psi$  and  $g = \sum_{\psi} b_\psi \psi$ , and set  $f * g = f \mu_{-1}(g)$ . Then the coefficient of the trivial character in  $f * g$  is  $\langle f, g \rangle = \sum_{\psi} a_\psi b_\psi$ . Expanding  $f * g$  in terms of its idempotent decomposition, we find

$$f * g = \frac{1}{n} \sum_{\psi \in G} \sum_{x \in A} f(x)g(-x)\psi^{-1}(x)\psi.$$

But by Theorem II.1 the coefficient of the trivial character is also  $n^{-1} \sum_x f(x)g(-x)$ .  $\square$

*Remark IV.2:* Since  $f$  and  $g$  lie in  $F[G]$ , the inner product lies in  $F$  even though the summation occurs in the extension  $K$ .

*Corollary IV.3:* Suppose that  $Z$  is stabilized by  $-1$ . Then the dual of  $C_Z$  is  $C_0^Z \oplus C_1^Z$ . If  $-1$  splits  $C_0$  then  $C_0^\perp = C_0^Z$ , and if  $-1$  stabilizes  $C_0$  then  $C_0^\perp = C_1^Z$ . In the latter case the ideal decomposition

$$F[G] = C_Z \oplus C_0^Z \oplus C_1^Z$$

is an orthogonal decomposition of  $F[G]$ .

*Proof:* Suppose that  $-1$  stabilizes  $Z$ , and let  $f$  lie in  $C_Z$  and  $g$  lie in  $C_0^Z \oplus C_1^Z$ . Then

$$\langle f, g \rangle = \frac{1}{n} \sum_{x \in A} f(x)g(-x) = 0,$$

since  $f(x) = 0$  for all  $x$  in  $X_0 \cup X_1$  and  $g(-x) = 0$  for all  $x$  in  $Z$ . The remaining orthogonality results are similar.  $\square$

We can now state the following theorem, which generalizes a well-known property of binary quadratic residue codes.

*Theorem IV.4:* Suppose that  $F = \mathbb{F}_2$  and  $C_0^Z$  is contained in its dual. Then the weight of every codeword in  $C_0^Z$  is congruent to 0 mod 4.

*Proof:* Since 0 lies in  $Z$ , every element  $f$  of  $C_0^Z$  satisfies  $f(0) = 0$ , hence has even weight. Since  $C_0^Z$  is contained in its dual,

$$0 = \langle f, g \rangle = |\text{Supp}(f) \cap \text{Supp}(g)| \pmod{2}$$

for all  $f$  and  $g$  in  $C_0^Z$ . From the equality

$$\|f + g\| = \|f\| + \|g\| - 2|\text{Supp}(f) \cap \text{Supp}(g)|,$$

we conclude that  $\|\cdot\| : C_0^Z \rightarrow \mathbb{Z}/4\mathbb{Z}$  is a well-defined group homomorphism with image in  $2\mathbb{Z}/4\mathbb{Z}$ . Let  $D_0^Z$  be the kernel of this map. Then it is clear that it is an ideal for  $F[G]$ , and is of codimension zero or one in  $C_0^Z$ . By Corollary III.7 there exists a complementary ideal  $J$  such that  $C_0^Z = D_0^Z \oplus J$ . By Proposition III.6 the only ideal of dimension one is that which corresponds to the orbit  $\{0\}$ . The idempotent generator of this ideal, the odd weight element  $\sum_{\psi \in G} \psi$ , does not lie in  $C_0^Z$ , so  $J = 0$  and  $C_0^Z = D_0^Z$ .  $\square$

### B. Code extensions

Let  $(Z, X_0, X_1)$  be a splitting of  $A$  giving a code  $C_0$  over  $F$ . Suppose that  $-1$  either splits or stabilizes the code and that  $F$  contains the image of  $Z$  under all characters in  $G$ .

Let  $F^Z = \prod_{z \in Z} F$  and define a map  $F[G]$  to  $F^Z$  by evaluation at elements of  $Z$ . For each  $f \in F[G]$  we define  $f_Z = (f(z))_{z \in Z}$ . The extended code  $\bar{C}_0$  is defined to be the subspace

$\bar{C}_0 = \{\tilde{f} = (f, f_Z) \mid f \in C_0\} \subseteq F[G] \times F^Z$ ,  
 $\bar{C}_1$  is defined similarly. We define an inner product on  $F^Z$  by

$$\langle u, v \rangle = -\frac{1}{n} \sum_{z \in Z} u_z v_{-z},$$

for all  $u = (u_z)_{z \in Z}$  and  $v = (v_z)_{z \in Z}$  in  $F^Z$ . Using this inner product on  $F^Z$ , we extend the usual Euclidean inner product on  $F[G]$  to the extended word space  $F[G] \times F^Z$  by defining  $F[G]$  and  $F^Z$  to be orthogonal subspaces.

*Theorem IV.5:* The extended codes  $\bar{C}_0$  and  $\bar{C}_1$  are equivalent. If  $-1$  splits  $C_0$ , then  $\bar{C}_0$  and  $\bar{C}_1$  are self dual, and if  $-1$  stabilizes  $C_0$ , then  $\bar{C}_0$  is dual to  $\bar{C}_1$ .

*Proof:* The equivalence of  $\bar{C}_0$  and  $\bar{C}_1$  follows immediately from the equivalence of  $C_0$  and  $C_1$  in Corollary III.9. Suppose that  $\mu_{-1}(C_0) = C_1$ , and let  $f$  and  $g$  be in  $C_0$ . Then  $f * g = f \mu_{-1}(g)$  lies in  $C_0 \cdot C_1 = C_0 \cap C_1$ . Thus  $f * g$  has the form

$$\sum_{z \in Z} f(z)g(-z)e_z = \frac{1}{n} \sum_{\psi \in G} \sum_{z \in Z} f(z)g(-z)\psi^{-1}(z)\psi.$$

On the other hand if we write  $f = \sum_{\psi} a_{\psi} \psi$  and  $g = \sum_{\phi} b_{\phi} \phi$ , then expanding the product, we find  $f * g$  equals

$$\left( \sum_{\psi \in G} a_{\psi} \psi \right) \left( \sum_{\phi \in G} b_{\phi} \phi^{-1} \right) = \sum_{\psi \in G} \sum_{\phi \in G} a_{\psi} b_{\phi} \psi \phi^{-1}.$$

Equating coefficients of the trivial character, we find that

$$\langle f, g \rangle = \sum_{\psi \in G} a_{\psi} b_{\psi} = \frac{1}{n} \sum_{z \in Z} f(z)g(-z) = -\langle f_Z, g_Z \rangle,$$

which implies the triviality of the inner product  $\langle \tilde{f}, \tilde{g} \rangle$ . The duality of  $\bar{C}_0$  and  $\bar{C}_1$  when  $\mu_{-1}(C_0) = C_0$  follows similarly.  $\square$

### C. Minimum distance bounds

Let  $C_0$  be the split group code over  $F$  relative to a splitting  $(Z, X_0, X_1)$ . We assume fixed an element  $s$  which splits  $C_0$  and for  $f \in F[G]$  define  $f^* = f \mu_s(f)$ .

Let  $N$  be the additive subgroup of  $A$  generated by  $Z$ . Define  $H$  to be the subgroup of  $G$  vanishing on  $Z$  (equivalently on  $N$ ), and let  $\mathcal{C}$  be a set of coset representatives for  $G/H$ . Then  $G/H$  is identified with the dual of  $N$  and  $H$  with the dual of  $A/N$ , whose order we denote by  $h$ . We define an element  $e_H$  by

$$e_H = \frac{1}{h} \sum_{\psi \in H} \psi.$$

*Lemma IV.6:* Let  $C_0$  be a split group code over a field of  $q$  elements. If  $\tau_s$  agrees with a power of  $\tau_q$  on  $Z$ , then for each  $z$  in  $Z$ ,  $f(z) = 0$  if and only if  $f^*(z) = 0$ .

*Proof:* If  $\tau_s = \tau_q^r$  on  $Z$ , then  $f(sz) = f(q^r z) = f(z)^{q^r}$ , for all  $z$  in  $Z$ .  $\square$

*Lemma IV.7:* If  $C_0$  is split by  $s$  then the subgroup  $H$  is stabilized by  $\mu_s$ .

*Proof:* Elements  $\phi$  in  $H$  are characterized by the condition that  $\phi(z) = 0$  for all  $z$  in  $Z$ . Since  $\tau_s$  stabilizes  $Z$ , for any  $\phi$  in  $H$  and  $z$  in  $Z$ ,  $\mu_s(\phi)(z) = \phi(sz) = 0$ , so  $\mu_s(\phi)$  lies in  $H$ .  $\square$

*Lemma IV.8:* The element  $e_H$  is the idempotent generator of  $I_{N^c}$ , and every  $g$  in  $C_Z$  is of the form

$$g = \sum_{\phi \in \mathcal{C}} c_{\phi} e_H \phi.$$

In particular, the support of  $g$  is a union of cosets of  $H$ .

*Proof:* Viewing  $H$  as the group of characters on  $A/N$ , the value of  $e_H$  equals 1 on  $N$  and zero elsewhere by the orthogonality relations for characters. Therefore  $e_H$  is the idempotent generator for  $I_{N^c}$ . Since  $Z^c$  contains  $N^c$ , the ideal  $I_{N^c}$  contains  $C_Z = I_{Z^c}$ , so every element  $g$  of  $C_Z$  can be written

$$g = f e_H = \sum_{\phi \in \mathcal{C}} c_{\phi} e_H \phi,$$

as indicated.  $\square$

*Proposition IV.9:* Let  $C_0$  be split by  $s$  and let  $f$  be in  $C_0$ . Then  $f^*$  is of the form  $\sum_{\phi \in \mathcal{C}} c_{\phi} e_H \phi$ . Moreover, if the support of  $f$  is contained in a coset of  $H$ , then  $f^*$  is of the form

$$f^* = c_{\xi} e_H \xi$$

for some  $\xi$  in  $G$ . If  $s = -1$  then the coefficient  $c_{\xi}$  of the trivial character  $\varepsilon$  is  $\langle f, f \rangle h$ .

*Proof:* Since  $f^*$  lies in  $C_0 C_1 = C_Z$ , by Lemma IV.8 we have  $f^* = \sum_{\phi \in \mathcal{C}} c_{\phi} e_H \phi$ . Since  $Q = \text{Supp}(f)$  is contained in a coset  $H\rho$  of  $H$ , by Lemma IV.7 the set  $\text{Supp}(\mu_s(f)) = \mu_s(Q)$  is contained in  $H\mu_s(\rho)$ . Then

$$\text{Supp}(f^*) \subseteq Q\mu_s(Q) \subseteq H\xi,$$

where  $\xi = \rho^*$ . Thus  $f^*$  has support on  $H\xi$  and is of the form  $f^* = c_{\xi} e_H \xi$ .

Now suppose that  $s = -1$ , then  $\xi = \rho\rho^{-1}$  is the trivial character. Write  $f = \sum_{\psi \in G} a_{\psi} \psi$ . Then the coefficient of the trivial character in the expansion

$$f^* = \sum_{\phi \in G} \sum_{\psi \in G} a_{\psi} a_{\phi} \psi \phi^{-1},$$

is  $\sum_{\psi \in G} a_{\psi}^2 = \langle f, f \rangle$ . It follows that  $c_{\varepsilon} = \langle f, f \rangle h$ .  $\square$

*Theorem IV.10:* Suppose that  $C_0$  is the split group code over a field of  $q$  elements, and suppose that  $\tau_s$  agrees with a power of  $\tau_q$  on  $Z$ . Then the minimum weight  $d$  of a codeword in  $C_0 \setminus C_0^Z$  satisfies the bound

$$h \leq \begin{cases} d^2 - d + 1 & \text{if } s = -1, \\ d^2 & \text{otherwise.} \end{cases}$$

*Proof:* If  $f$  lies in  $C_0 \setminus C_0^Z$ , then  $f^*$  is nonzero by Lemma IV.6. Since  $f^*(x) = 0$  for all  $x$  in  $X_0 \cup X_1$ , we have

$$f^* = \sum_{z \in Z} f^*(z) e_z = \frac{1}{n} \sum_{\psi \in G} \sum_{z \in Z} f^*(z) \psi(z)^{-1} \psi.$$

Since  $\phi(z) = \psi(z)$  for all  $\phi$  and  $\psi$  in the same coset of  $H$  and all  $z$  in  $Z$ , the sum for  $f^*$  can be expressed as

$$f^* = \frac{h}{n} \sum_{\phi \in C} \left( \sum_{z \in Z} f^*(z) \phi(z)^{-1} \right) e_H \phi,$$

Thus  $f^*$  has a positive multiple of  $h$  nonzero coefficients. On the other hand  $f^*$  has at most  $d^2$  nonzero coefficients, which implies  $d^2 \geq h$ . If  $C_0$  is split by  $s = -1$ , then  $d$  coefficients of expansion for  $f^*$  contribute to the coefficient of the trivial character, so  $d^2 - d + 1 \geq h$ .  $\square$

We define an *incidence relation*  $i : K[G] \times G \rightarrow \{0, 1\}$  by setting  $i(f, \psi) = 1$  if  $\psi$  is in the support of  $f$  and  $i(f, \psi) = 0$  otherwise. In reference to the incidence structure, we refer to functions in  $K[G]$  as *lines* and characters in  $G$  as *points*. Moreover, if  $i(f, \psi) = 1$ , then we say that  $f$  *meets*  $\psi$ .

*Theorem IV.11:* Let  $C_0$  be a split group code over  $F$  which is split by  $-1$ , and suppose that  $\tau_{-1} = \tau_q^r$  on  $Z$  for some integer  $r$ . If  $C_0 \setminus C_0^Z$  has a codeword  $f$  of minimum weight  $d$ , satisfying  $h = d^2 - d + 1$ , then if  $f$  meets the trivial character, the following statements hold.

1. The set  $Q = \text{Supp}(f)$  is a difference set for  $H$  with parameter  $\lambda = 1$ .
2. Each coset  $\mathbb{P}_\phi = H\phi$  comprises the set of points of a combinatorial projective plane of order  $d - 1$  with respect to the set of lines  $\mathcal{L}_\phi = fH\phi$ .
3. The minimum distance of  $C_0$  is  $d$ .

*Proof:* Since  $f^*$  lies in  $C_Z$ , by Proposition IV.9 its support is a union of cosets of  $H$ . On the other hand  $\text{Supp}(f^*)$  lies in  $\{\psi\phi^{-1} \mid \psi, \phi \in Q\}$ , so has at most  $h = d^2 - d + 1$  elements. Since  $f^*$  is nonzero, it follows that  $\text{Supp}(f^*) = H$ , and every nontrivial element of  $H$  can be uniquely represented in the form  $\psi\phi^{-1}$  for  $\psi$  and  $\phi$  in  $Q$ . Since  $f$  meets the trivial character,  $Q$  is contained in  $H$ , and the first statement holds.

The support of a line in  $\mathcal{L}_\phi = fH\phi$  is clearly contained in the set  $\mathbb{P}_\phi = H\phi$ . It suffices to show that two distinct lines in  $\mathcal{L}_\phi$  meet at a unique point. Let  $g_1 = f\nu$  and  $g_2 = f\xi$  be two such lines with support  $Q_1 = Q\nu$  and  $Q_2 = Q\xi$ , respectively. Consider the product map

$$Q_1 \times Q_2^{-1} \rightarrow H\nu\xi^{-1} = H$$

given by the restriction of the group law on  $G$ . The inverse image of  $\nu\xi^{-1}$  has  $d$  elements and elsewhere the map is bijective. In particular, since  $\nu \neq \xi$ , there is a unique pair  $(\psi, \psi^{-1})$  mapping to the trivial character, implying  $g_1$  and  $g_2$  meet uniquely at the point  $\psi$ .

Now let  $g$  be in  $C_0^Z$ . We may assume that  $g$  meets the trivial character.  $C_0$  and  $C_0^Z$  are dual by Corollary IV.3, so  $\langle f\psi^{-1}, g \rangle = 0$ . Thus if  $f\psi^{-1}$  and  $g$  meet, they must do so at more than one point. By construction  $g$  and  $f\psi^{-1}$  meet at the trivial character for all  $\psi$  in  $Q$ . Since the lines  $\{f\psi^{-1} \mid \psi \in Q\}$  are pairwise disjoint away from the trivial character, it follows that  $g$  has weight at least  $d + 1$ .  $\square$

## V. EXAMPLES OF SPLIT GROUP CODES

In this section we demonstrate two constructions by which we remove the restriction that  $q$  be a square modulo all prime divisors of the block length. We focus on the cyclic case; examples of the general construction for non-cyclic abelian groups will be reserved for treatment in a later article.

### A. Dual nonresidue split group codes.

Let  $\ell$  and  $m$  be distinct primes such that  $q$  is not a square mod  $\ell$  or mod  $m$ . Set  $R = \mathbb{Z}/n\mathbb{Z}$  where  $n = \ell m$ , and let  $A$  its abelian group. Let

$$\left( \frac{\cdot}{n} \right) : \mathbb{Z}/n\mathbb{Z} \rightarrow \{0, 1, -1\}$$

be the Kronecker symbol (see §1.4 in Cohen [4]). We set  $Z = m\mathbb{Z}/n\mathbb{Z} \cup \ell\mathbb{Z}/n\mathbb{Z}$  and take  $X_0$  to be the set

$$X_0 = \{a \in R \mid \left( \frac{a}{n} \right) = 1\},$$

which is stable under  $\langle \tau_q \rangle$ , and  $X_1$  its complement outside of  $Z$ . The following theorem shows that the codes  $C_0$  relative to this splitting are bad.

*Theorem V.1:* Suppose  $(Z, X_0, X_1)$  is a splitting of an abelian group  $A$  such that  $Z$  contains a subgroup  $N$  of order  $m$ . Let  $G$  be the dual group of  $A$ , and let  $H$  be the subgroup which is trivial on  $N$ . Then  $f = \sum_{\phi \in H} \phi$  is a codeword in  $C_0$  of weight  $n/m$ . In particular the minimum distance is bounded above by  $n/m$ .

*Proof:* We may view  $H$  as the group of characters on  $A/N$ . Then  $f(x) = \sum_{\phi \in H} \phi(1) = |H|$  for  $x$  in  $N$ , and by the orthogonality relations for characters, is 0 on all other elements of  $A$ . Since  $X_0$  does not meet  $N \subseteq Z$ , it is clear that  $f$  is in  $C_0$ .  $\square$

Applying the theorem to the construction, we find that the code  $C_0$  has minimum distance bounded above by  $\ell$ , the smaller of the two prime divisors of  $n$ . In contrast, the subcode  $C_0^Z$  contains no obvious codeword of small weight, and experimentally, appears to generally have large minimum weight. The simplest example is that in Example III.1. In Table I we present data for the block length, dimension, and minimum weight of these codes over  $\mathbb{F}_2$  up to block length 209.

### B. Twisted lifts of split group codes.

Although the codes  $C_0^Z$  described above perform well, in order to have reasonable minimum distance for  $C_0$ , it is clear from Theorem V.1 that we should avoid splittings for which  $Z$  contains a large subgroup of  $A$ . Also in light of Theorem IV.11 we may want to consider splittings for which  $Z$  is a small subgroup of  $A$ . We thus present another example in which the block length is divisible by a single small prime in which  $q$  is a quadratic nonresidue.

*Example V.2:* Let  $F = \mathbb{F}_2$ , set  $R = \mathbb{Z}/21\mathbb{Z}$  and let  $A$  be its additive group. Since the subset  $7R^*$  of  $A$  consists of a single  $\langle \tau_2 \rangle$ -orbit of two elements, we set  $Z = 7\mathbb{Z}/21\mathbb{Z}$ , and



TABLE I  
DUAL NONRESIDUE CYCLIC SPLIT CODES OVER  $\mathbb{F}_2$

$C_0$					$C_0^Z$		
$\ell$	$m$	$n$	$k$	$d^a$	$n$	$k$	$d^a$
3	5	15	11	3	15	4	8
3	11	33	23	3	33	10	12
3	13	39	27	3	39	12	12
5	11	55	35	5	55	20	16
3	19	57	39	3	57	18	16
5	13	65	41	5	65	24	16
5	13 <sup>c</sup>	65	41	5	65	24	16
3	29	87	59	3	87	28	24
5	19	95	59	5	95	36	16
3	37	111	75	3	111	36	24
3	43	129	87	3	129	42	28 <sup>b</sup>
11	13	143	83	11	143	60	24 <sup>b</sup>
5	29	145	89	5	145	56	24 <sup>b</sup>
5	29 <sup>c</sup>	145	89	5	145	56	26 <sup>b</sup>
3	53	159	107	3	159	52	32 <sup>b</sup>
3	59	177	119	3	177	58	30 <sup>b</sup>
3	61	183	123	3	183	60	36 <sup>b</sup>
5	37	185	113	5	185	72	24 <sup>b</sup>
5	37 <sup>c</sup>	185	113	5	185	72	32 <sup>b</sup>
3	67	201	135	3	201	66	36 <sup>b</sup>
11	19	209	119	11	209	90	30 <sup>b</sup>

<sup>a</sup>Minimum distance computed with Magma [8].

<sup>b</sup>Upper bound found by probabilistic search for minimum weight vectors with Magma; in each case the minimum distance is provably bounded below by 18.

<sup>c</sup>For  $\gcd(\ell-1, m-1)$  greater than 2 there exist additional splittings over  $Z$ ; for the pairs (5, 13), (5, 29), and (5, 37) we find an additional inequivalent splitting, reported here.

take  $X_0$  to be

$$X_0 = \{a \in R \mid \left(\frac{a}{7}\right) = 1\},$$

where  $(a/7)$  is the Legendre symbol. Since 2 is a quadratic residue in  $\mathbb{Z}/7\mathbb{Z}$ , it is clear that  $\tau_2$  stabilizes the splitting. Moreover,  $-1$  is a quadratic nonresidue mod 7, so gives the splitting. The associated split group code, denoted  $Q_0$ , is a  $[21, 12, 3]$ -code, while the subcode  $Q_0^Z$  of functions vanishing on  $Z$  is a  $[21, 9, 4]$ -code, both poor codes. If instead we choose  $X_0$  equal to

$$\{a \in R^* \mid \left(\frac{a}{7}\right) = 1\} \cup \{a \in 3R^* \mid \left(\frac{a}{7}\right) = -1\},$$

then we obtain a split group code  $C_0$  with parameters  $[21, 12, 5]$  and subcode  $C_0^Z$  with parameters  $[21, 9, 8]$ , both of which are best possible for length 21 and their respective dimensions.  $\square$

In the next construction we develop this idea further, showing that the special case above is typical. We describe first a formal construction for provably bad codes, and discuss how to “twist” them to obtain codes which experimentally perform well.

Let  $(W, Y_0, Y_1)$  be a splitting of the additive group of a finite ring  $S$  with associated split group code  $Q_0$  and subcode  $Q_0^W$ . Suppose that there exists a surjective homomorphism  $\pi : R \rightarrow S$  with kernel of order  $\ell$ . We form the

lifted splitting  $(Z, X_0, X_1)$  of  $A$  by setting

$$Z = \pi^{-1}(W), \quad X_0 = \pi^{-1}(Y_0), \quad \text{and} \quad X_1 = \pi^{-1}(Y_1),$$

and let  $C_0$  and  $C_0^Z$  be the associated split group codes. For such codes we have the following theorem, which shows that these codes are bad.

*Theorem V.3:* The weight enumerator polynomial of  $C_0$  is  $w(T)^\ell$ , where  $w(T)$  is the weight enumerator polynomial of  $Q_0$ . The same relation holds between the respective weight enumerator polynomials of  $C_0^Z$  and of  $Q_0^W$ . In particular the minimum distances of  $C_0$  and  $C_0^Z$  are the same as the minimum distances of  $Q_0$  and  $Q_0^W$ , respectively.

*Proof:* Denote the additive groups of  $R$  and  $S$  by  $A$  and  $B$ , respectively, and set  $M = \text{Hom}(B, K^*)$  and  $G = \text{Hom}(A, K^*)$ . Then the pullback  $\pi^* : M \rightarrow G$  is an injective homomorphism with cokernel of order  $\ell$ . Denote also by  $\pi^*$  the induced ring homomorphism  $F[M] \rightarrow F[G]$ . Under this homomorphism  $F[G]$  decomposes as an  $F[M]$ -module into a direct sum  $F[G] = \bigoplus \pi^*(F[M]) \psi$ , where  $\psi$  ranges over coset representatives of  $G/M$ . Consider the idempotent  $e_0$  for  $Q_0$ . Then by definition, for all  $x$  in  $A$ ,  $\pi^*(e_0)$  satisfies

$$\pi^*(e_0)(x) = e_0(\pi(x)) = \begin{cases} 0 & \text{if } \pi(x) \in Y_0, \\ 1 & \text{otherwise,} \end{cases}$$

so is the idempotent for  $C_0$ . In particular  $C_0$  is generated by the image of  $Q_0$ , so we have an  $F[M]$ -module decomposition

$$C_0 = F[G] \pi^*(Q_0) = \bigoplus \pi^*(Q_0) \psi.$$

The form of the weight enumerator polynomial follows from the fact that each  $\pi^*(F[M]) \psi$ , hence  $\pi^*(Q_0) \psi$ , has disjoint support in  $G$ , and from the independence of the equivalent subcodes  $\pi^*(Q_0) \psi$ . The decomposition of  $C_0^Z$  into copies of  $Q_0^W$  follows by the same argument, and equality of the minimum distances is then clear.  $\square$

We apply the above theorem to the following setting. Let  $S = \mathbb{Z}/m\mathbb{Z}$ , for a prime  $m$ , let  $R = \mathbb{Z}/\ell m\mathbb{Z}$ , and let  $\pi$  be the surjective homomorphism  $R \rightarrow S$ . Let  $Q_0$  and  $Q_0^W$  be the quadratic residue codes of length  $m$  defined by the splitting  $(\{0\}, Y_0, Y_1)$ , where

$$Y_0 = \{a \in S \mid \left(\frac{a}{m}\right) = 1\},$$

and  $Y_1$  is the complement in  $S \setminus \{0\}$ .

First we consider the splitting over  $Z = m\mathbb{Z}/\ell m\mathbb{Z}$  with  $X_0 = \pi^{-1}(Y_0)$  and  $X_1 = \pi^{-1}(Y_1)$ ; specifically we note that

$$X_0 = \{a \in R \mid \left(\frac{a}{m}\right) = 1\}.$$

The associated codes  $C_0$  and  $C_0^Z$  are liftings of  $Q_0$  and  $Q_0^W$ , so by Theorem V.3 the minimum distances of  $C_0$  and  $C_0^Z$  are exactly the same as for the quadratic residue codes themselves.

Instead, we note that the set  $R \setminus Z$  splits into the orbits of  $R^*$  and those of  $\ell R^*$ . We thus choose  $X_0$  to be the subset of  $A \setminus Z$  twisted by a quadratic nonresidue mod  $m$  on one of these sets:

$$\{a \in R^* \mid \left(\frac{a}{m}\right) = 1\} \cup \{a \in \ell R^* \mid \left(\frac{a}{m}\right) = -1\}.$$

Then any  $s$  in  $R^*$  which is not a square mod  $m$  splits  $(Z, X_0, X_1)$ , but this twisted lift avoids the conditions of

Theorem V.3 which produced poor codes. Indeed in the tables below we find that the minimum distances of the resulting twisted lifts well exceed those of the corresponding quadratic residue codes, and give some new minimum distance records.

From Theorem V.3 it is clear that for each  $[m, k, d]$ -code above there exists a lifted  $[\ell m, \ell k, d]$ -code. Thus we provide Table II as reference for the parameters of quadratic residue codes of block length  $m$ , and in Table III give only the parameters of the twisted lifts, taking  $\ell = 3$ . In view of Theorem IV.10 we also indicate the smallest value  $d_0$  for which  $d_0^2 \geq m$  when  $m \equiv 1 \pmod{8}$ , or for which  $d_0^2 - d_0 + 1 \geq m$  when  $m \equiv 7 \pmod{8}$ .

We note that the minimum distances of the examples in Table III are consistently close to the best known of their length and dimension. In particular, the  $[21, 12, 5]$  and  $[93, 48, 14]$  codes  $C_0$  and the  $[21, 9, 8]$  code  $C_0^Z$  match the best known in Brouwer [2]. The minimum distance of the  $[141, 69, 24]$  code  $C_0^Z$  gives a new minimum distance record for codes of length 141 and dimension 69. By the Spoiling Lemma [19, Lemma 1.1.34] this improves the lower bound on the best possible minimum distance of a code for the ten values of  $[n, k]$  listed in Table IV.

TABLE II  
QUADRATIC RESIDUE CODES OVER  $\mathbb{F}_2$

$C_0$			$C_0^Z$			$d_0$
$m$	$k$	$d^a$	$m$	$k$	$d^a$	
7	4	3	7	3	4	3
17	8	5	17	7	6	5
23	12	7	23	11	8	6
31	16	7	31	15	8	6
41	21	9	41	20	10	7
47	24	11	47	23	12	8
71	36	11	71	35	12	9
73	37	13	73	36	14	9
79	40	15	79	38	16	9
89	45	17	89	44	18	10
97	49	15	97	48	16	10
103	52	19	103	51	20	11
113	57	15	113	56	16	11
127	64	19	127	63	20	12

<sup>a</sup>Minimum distance computed with Magma [8].

## VI. CONCLUSION

We have shown that the main results and methods for various abelian group codes can be studied as a uniform family of codes, and that these generalize duadic codes. Moreover, within this family there exist subclasses not included in the previously described family of duadic codes. As demonstrated by examples constructed within special subclasses, these include codes which have good parameters. It is also of interest that results are obtained regarding classes of codes which have poor parameters. The exam-

TABLE III  
TWISTED LIFTS OF QR CODES OVER  $\mathbb{F}_2$  FOR  $\ell = 3$

$m$	$C_0$			$C_0^Z$			$d_0$
	$n$	$k$	$d^a$	$n$	$k$	$d^a$	
7	21	12	5	21	9	8	3
17	51	27	9	51	24	10	5
23	69	36	11	69	33	12	6
31	93	48	14	93	45	16	6
41	123	63	18	123	60	20	7
47	141	72	21	141	69	24	8
71	213	108	22 <sup>b</sup>	213	105	24 <sup>b</sup>	9
73	219	111	26 <sup>b</sup>	219	108	28 <sup>b</sup>	9
79	237	120	30 <sup>b</sup>	237	117	32 <sup>b</sup>	9
89	267	135	35 <sup>b</sup>	267	132	36 <sup>b</sup>	10
97	291	147	29 <sup>b</sup>	291	144	30 <sup>b</sup>	10
103	309	156	39 <sup>b</sup>	309	153	40 <sup>b</sup>	11
113	339	171	46 <sup>b</sup>	339	168	48 <sup>b</sup>	11
127	391	197	57 <sup>b</sup>	381	194	60 <sup>b</sup>	12

<sup>a</sup>Minimum distance computed with the Magma [8].

<sup>b</sup>Upper bound found by probabilistic search for minimum weight vectors with Magma; in each case the minimum distance is provably bounded below by 16.

TABLE IV  
NEW MINIMUM DISTANCE BOUNDS OVER  $\mathbb{F}_2$

$n$	$k$	$d$	$d_0^a$
141	69	24	23
140	69	23	22
140	68	24	23
139	68	23	22
139	67	24	23
138	67	23	22
138	66	24	23
137	66	23	22
137	65	24	23
136	65	23	22

<sup>a</sup>Minimum distance of previously best known code in Brouwer [2].

ples in this work emphasize the cyclic split group codes; in future work we expect to extend the computational efforts to split group codes in for noncyclic abelian groups, which include the generalized quadratic residue codes and generalized duadic codes.

**Acknowledgement.** D. Kohel was supported by an NSTB grant as a Postdoctoral Fellow at the National University of Singapore when this work was written, and C. Ding and S. Ling were partially supported by the NSTB-MOE grant number RP 960668/M. Part of this work was done while S. Ling was visiting the Graduate School of Mathematical Sciences of the University of Tokyo under the Hitachi Fellowship from the Hitachi Scholarship Foundation. He would like to thank these institutions and Professor S. Nakajima for their hospitality and support. The

authors thank the referees for their valuable comments that improved this paper considerably.

#### REFERENCES

- [1] M. F. Atiyah and I. G. MacDonald. *Introduction to commutative algebra*. Reading, MA: Addison-Wesley, 1969.
- [2] A. E. Brouwer. "Bounds on the size of linear codes," in *Handbook of Coding Theory. Vol. I*. V. Pless and W. Huffman, Eds. Amsterdam: North-Holland, 1998, pp. 295-462.
- [3] P. Camion. "Global quadratic Abelian codes," in *Information Theory*, G. Longo, Ed. *CISM Courses and Lectures*, vol.219, Vienna: Springer-Verlag, 1975, pp. 293-310.
- [4] H. Cohen. *A Course in Computational Algebraic Number Theory. Graduate Texts in Mathematics*, vol.138, Berlin: Springer-Verlag, 1993.
- [5] D. Knuth. *The Art of Computer Programming. Volume 2: Seminumerical algorithms*, 2nd ed. Reading, MA: Addison-Wesley, 1981.
- [6] S. Lang. *Algebra*. 3rd ed. Reading, MA: Addison-Wesley, 1993.
- [7] J. S. Leon, J. M. Masley, and V. Pless. "Duadic codes," *IEEE Trans. Inform. Theory*, vol.30, pp. 709-714, 1984.
- [8] W. Bosma, J. Cannon, and C. Playoust. "The Magma algebra system I: The user language," *J. Symb. Comp.*, vol.24, no. 3-4, pp. 235-265, 1997.
- [9] J. H. van Lint and F. J. MacWilliams. "Generalized quadratic residue codes," *IEEE Trans. Inform. Theory*, vol.24, no. 6, pp. 730-737, 1978.
- [10] J. H. van Lint. *Introduction to Coding Theory. Graduate Texts in Mathematics*, vol.86, Berlin: Springer-Verlag, 1992.
- [11] V. Pless. "Q-Codes," *J. Combin. Theory Ser. A*, vol.43, pp. 258-276, 1986.
- [12] V. Pless, J. M. Masley, and J. S. Leon. "On weights in duadic codes," *J. Combin. Theory Ser. A*, vol.44, pp. 6-21, 1987.
- [13] V. Pless. "Duadic codes revisited," *Congressus Numerantium*, vol.59, pp. 225-233, 1987.
- [14] V. Pless. "Duadic codes and their generalizations," *Eurocode '92 (Udine, Italy), CISM Courses and Lectures*, vol.339, pp. 3-15, Vienna: Springer-Verlag, 1993.
- [15] J. J. Rushanan. "Topics in integral matrices and Abelian group codes," Ph.D. dissertation, Caltech, 1986.
- [16] J. J. Rushanan. "Duadic codes and difference sets," *J. Combin. Theory Ser. A*, vol.57, pp. 254-261, 1991.
- [17] M. Smid. "On Duadic Codes," Dept. Math., Tech. Univ. Eindhoven, The Netherlands, Tech. Report 86-WSK-04, May 1986.
- [18] M. Smid. "Duadic Codes," *IEEE Trans. Inform. Theory*, vol.33, no. 3, pp. 432-433, 1987.
- [19] M. A. Tsfasman and S. G. Vladut. *Algebraic-Geometric Codes. Mathematics and its Applications (Soviet Series)*, vol.58, Dordrecht: Kluwer, 1991.
- [20] H. N. Ward. "Quadratic residue codes and symplectic groups," *J. Algebra*, vol.29, pp. 150-171, 1974.

Cunsheng Ding, Department of Computer Science, National University of Singapore, Singapore 117543  
E-mail: dingcs@comp.nus.edu.sg

David R. Kohel, School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia  
E-mail: kohel@maths.usyd.edu.au

San Ling, Department of Mathematics, National University of Singapore, Singapore 117543  
E-mail: matlings@math.nus.edu.sg