

NONSPECIAL VARIETIES AND GENERALIZED LANG-VOJTA CONJECTURES

ERWAN ROUSSEAU, AMOS TURCHET, AND JULIE TZU-YUEH WANG

ABSTRACT. We construct a family of fibered threefolds $X_m \rightarrow (S, \Delta)$ that are weakly special but not special in the sense of Campana. We prove that if m is big enough, the threefolds X_m present behaviours that contradict the function field and analytic analogues of a conjecture of Abramovich and Colliot-Thélène, thus providing evidences towards Campana’s conjecture. This is obtained using an orbifold analogue of [CZ10, Main Theorem] in the function field and analytic setting, which we prove adapting the recent method of Ru and Vojta in [RV]. We also formulate some generalizations of known conjectures on exceptional loci that fit into Campana’s program and prove some cases over function fields.

1. INTRODUCTION

A fundamental problem in Diophantine Geometry is to describe the distribution of rational points in an algebraic variety X defined over a number field k . The expectation is that the global geometry of X controls the distribution of $X(k)$. In the case in which X is a variety of general type, conjectures of Lang and Vojta (see e.g. [Lan91, Conjecture 3.7], [Voj87, Conjecture 3.4.3]) predict the existence of a proper Zariski closed subset, called the *exceptional set*, such that in its complement rational points $X(L)$ should be finite for every finite extension $L \supset k$; moreover the exceptional set is expected to be independent of the field of definition. On the opposite side there are the varieties whose rational points are potentially dense, i.e. there exists a finite field extension L/k such that the set of L -rational points $X(L)$ is Zariski dense. The conjectures of Lang and Vojta thus predict that rational points in a variety of general type are not potentially dense.

It is natural to look for geometrical properties characterizing varieties where the set of rational points is potentially dense. The conjectures mentioned above imply that for such a property to be satisfied on X , not only X but none of its finite étale covers should dominate a positive-dimensional variety of general type: such varieties are called *weakly special* (see Section 4 for details). Abramovich and Colliot-Thélène [HT00, Conjecture 1.2] conjectured that a variety X has a potentially dense set of rational points if and only if X is weakly special.

In [Cam04], Campana introduced the stronger notion of *specialness*: a variety X is *special* if it does not admit any fibration of general type, in the sense of Campana (see Section 2 for details). Campana conjectured that specialness rather than weak-specialness should characterize potential density. Note that these two characterizations differ from each other (even if they agree in dimension up to two) since there exist projective varieties which are weakly special but not special.

2010 *Mathematics Subject Classification.* 14G40, 11J97, 14G05, 32A22.

Key words and phrases. Campana’s conjectures, function fields, Nevanlinna Theory, orbifolds, hyperbolicity.

Such examples were first constructed in [BT04] as simply-connected threefolds equipped with an elliptic fibration $\pi : X \rightarrow S$ where S has Kodaira dimension 1. Nevertheless, these examples do not explicitly contradict Abramovich and Colliot-Thélène’s conjecture since we still lack a method to control the distribution of rational points on these varieties. Therefore it is natural to consider the analogous problem in the analytic and function fields settings. The goal of this article is to study the two conjectures in these setting giving evidence towards Campana conjecture.

In the analytic setting, the Green–Griffiths–Lang conjecture predicts that entire curves in varieties of general type should be contained in a proper Zariski closed subset, the already mentioned exceptional set. Following this analogy, Campana has conjectured that specialness (and therefore potential density) should correspond to the existence of Zariski dense entire curves (see [Cam04]). The analytic analogue of Abramovich and Colliot-Thélène conjecture would imply that a weakly special variety admits a Zariski dense entire curve. This was already disproved in [CP07], where it was proven that in some of the examples of weakly special but not special varieties constructed in [BT04], all entire curves are algebraically degenerate (i.e. with non Zariski dense image).

Similarly, the conjectures of Lang and Vojta admit analogues for varieties defined over function fields. In our setting, Vojta’s height conjecture predicts that given a general type variety X defined over a characteristic zero function field $\kappa(\mathcal{C})$ of a smooth integral curve \mathcal{C} , and an ample line bundle \mathcal{L} , there exists a positive constant α such that sections $s : \mathcal{C} \rightarrow X$ not contained in the exceptional set, satisfy $\deg s^* \mathcal{L} \leq \alpha(2g(\mathcal{C}) - 2)$. Following Demailly’s notion of algebraic hyperbolicity in [Dem97], we say that varieties X that satisfies this condition are *pseudo algebraically hyperbolic*. A function field analogue of Abramovich and Colliot-Thélène conjecture would imply that a weakly special variety X admits no dominant map $\pi : X \rightarrow Y$ such that there is a degree bound as above for morphisms $\pi \circ s : \mathcal{C} \rightarrow Y$.

One of the goals of this paper is to present a list of new examples of varieties that are weakly special but not special: these are of the form $\pi : X_m \rightarrow (S, \Delta_\pi)$ where X_m is a threefold fibered in elliptic curves, \bar{S} is obtained as a blow-up of \mathbb{P}^2 and $\Delta_\pi = (1 - 1/m)\widetilde{D}_1$ is an orbifold divisor (see Section 6 for the precise definition). The pairs (S, Δ_π) can be seen as orbifold generalizations of surfaces appearing in [CZ10] which provide examples of simply-connected quasi-projective surface with a non Zariski dense set of integral points. For this class of examples we show that the function field and analytic analogues of Abramovich and Colliot-Thélène conjecture fail. In particular, we prove the following:

Theorem 1.1 (see Theorem 6.5). *There exists m_0 such that for all $m \geq m_0$, the following hold:*

- (1) *given an ample line bundle \mathcal{L} on S , there exists a positive constant α and a closed subset $Z \subset \bar{S}$ such that for every section $s : \mathcal{C} \rightarrow \bar{X}_m$, such that $(\pi \circ s)(\mathcal{C}) \not\subset Z$, the following holds:*

$$\deg(\pi \circ s)^* \mathcal{L} \leq \alpha \cdot \left(2g(\mathcal{C}) - 2 + N_{\pi \circ s}^{[1]}(\bar{S} \setminus S) \right);$$

- (2) *every entire curve in X_m is algebraically degenerate.*

To prove Theorem 1.1 we combine Corvaja-Zannier degeneracy statements in [CZ04] with the recent framework of Ru and Vojta [RV] to obtain a generalization of [CZ04, Main Theorem] in the orbifold setting for the function field and analytic case.

Theorem 1.2 (see Corollary 5.8 and Theorem 5.11). *Let $X \subset \mathbb{P}^m$ be a complex nonsingular projective surface and $D = D_1 + \cdots + D_q$ be a divisor with $q \geq 2$, such that*

- (1) *No three of the components D_i meet at a point;*
- (2) *There exists a choice of positive integers p_i such that*
 - *the divisor $D_p := p_1D_1 + p_2D_2 + \cdots + p_qD_q$ is ample;*
 - *The following inequality holds:*

$$2D_p^2\xi_i > (D_p \cdot D_i)\xi_i^2 + 3D_p^2p_i,$$

for every $i = 1, \dots, q$ where ξ_i is the minimal positive solution of the equation $D_i^2x^2 - 2(D_p \cdot D_i)x + D_p^2 = 0$.

Let Δ be the \mathbb{Q} -divisor defined as

$$\Delta = \sum_{i=1}^q \left(1 - \frac{1}{m_i}\right) D_i.$$

Then, there exists a positive integer m such that, if $m_i \geq m$ for every i ,

- (1) *every orbifold entire curve $\psi : \mathbb{C} \rightarrow (X, \Delta)$, is algebraically degenerate;*
- (2) *(X, Δ) is pseudo algebraically hyperbolic.*

Structure of the paper.

In Section 2 we recall basic facts about special varieties and orbifolds, following Campana. Then, in Section 3, we present a general framework where we generalize Lang’s notion of exceptional loci to nonspecial varieties and Campana orbifolds. In Section 4 we discuss a general procedure to construct weakly special but not special varieties, generalizing [BT04]. In Section 5 we prove Theorem 1.2 and we apply it in Section 6 to prove Theorem 1.1. Finally, in Section 7 we prove some generalizations of Ru and Vojta [RV, General Theorem] for function fields and entire curves.

Acknowledgements

We thank Laura Capuano, Pietro Corvaja, Lionel Darondeau, Ariyan Javanpeykar, Stefan Kebekus and Min Ru for useful discussions. We thank Min Ru and Paul Vojta for providing us with an earlier copy of their paper. We are especially grateful to Frédéric Campana for all the discussions on this subject and especially his explanations on the construction of weakly special but not special varieties.

This collaboration was initiated at the conference “Topics On Nevanlinna Theory And Complex Hyperbolicities” at the Shanghai Center for Mathematical Sciences; we thank the organizers for making this possible. Part of this work was finalized during the visit of the first and second named author to the Freiburg Institute for Advanced Studies; they thank the Institute for providing an excellent working environment.

Research of E.R. was supported by the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 754340, and by the ANR project “FOLIAGE”, ANR-16-CE40-0008. Research of A.T. was supported in part by funds from NSF grant DMS-1553459 and in part by Centro di Ricerca Matematica Ennio de Giorgi. Research of J.W. was supported by Taiwan’s MoST 108-2115-M-001-001-MY2.

2. SPECIAL VARIETIES

We collect here basic definitions and constructions related to special varieties, while referring to [Cam04] for more details.

2.1. *Special Manifolds via Bogomolov sheaves*

Let X be a connected complex nonsingular projective variety of complex dimension n . For a rank-one coherent subsheaf $\mathcal{L} \subset \Omega_X^p$, denote by $H^0(X, \mathcal{L}^m)$ the space of sections of $\mathrm{Sym}^m(\Omega_X^p)$ which take values in \mathcal{L}^m (where as usual $\mathcal{L}^m := \mathcal{L}^{\otimes m}$). The *Iitaka dimension* of \mathcal{L} is $\kappa(X, \mathcal{L}) := \max_{m>0} \{\dim(\Phi_{\mathcal{L}^m}(X))\}$, i.e. the maximum dimension of the image of rational maps $\Phi_{\mathcal{L}^m} : X \dashrightarrow \mathbb{P}(H^0(X, \mathcal{L}^m))$ defined at the generic point of X , where by convention $\dim(\Phi_{\mathcal{L}^m}(X)) := -\infty$ if there are no global sections. Thus $\kappa(X, \mathcal{L}) \in \{-\infty, 0, 1, \dots, \dim(X)\}$. In this setting, a theorem of Bogomolov in [Bog79] shows that, if $\mathcal{L} \subset \Omega_X^p$, then $\kappa(X, \mathcal{L}) \leq p$.

Definition 2.1. Let $p > 0$. A rank one saturated coherent sheaf $\mathcal{L} \subset \Omega_X^p$ is called a *Bogomolov sheaf* if $\kappa(X, \mathcal{L}) = p$, i.e. if \mathcal{L} has the largest possible Iitaka dimension.

The following remark shows that the presence of Bogomolov sheaves on X is related to the existence of fibrations $f : X \rightarrow Y$ where Y is of general type.

Remark 2.2. If $f : X \rightarrow Y$ is a fibration (by which we mean a surjective morphism with connected fibers) and Y is a variety of general type of dimension $p > 0$, then the saturation of $f^*(K_Y)$ in Ω_X^p is a Bogomolov sheaf of X ,

Campana introduced the notion of specialness in [Cam04, Definition 2.1] to generalize the absence of fibration as above.

Definition 2.3. A nonsingular variety X is said to be *special* (or *of special type*) if there is no Bogomolov sheaf on X . A projective variety is said to be *special* if some (or any) of its resolutions are special.

By the previous remark if there is a fibration $X \rightarrow Y$ with Y of general type then X is nonspecial. In particular, if X is of general type of positive dimension, X is not of special type.

2.2. *Special Manifolds via orbifold bases*

Campana gave a characterization of special varieties using his theory of orbifolds. We briefly recall the construction.

Let Z be a normal connected compact complex variety. An *orbifold divisor* Δ is a linear combination $\Delta := \sum_{\{D \subset Z\}} c_\Delta(D) \cdot D$, where D ranges over all prime divisors of Z , the *orbifold coefficients* are rational numbers $c_\Delta(D) \in [0, 1] \cap \mathbb{Q}$ such that all but finitely many are zero. Equivalently,

$$\Delta = \sum_{\{D \subset Z\}} \left(1 - \frac{1}{m_\Delta(D)}\right) \cdot D,$$

where only finitely *orbifold multiplicities* $m_\Delta(D) \in \mathbb{Q}_{\geq 1} \cup \{+\infty\}$ are larger than 1.

An orbifold pair is a pair (Z, Δ) where Δ is an orbifold divisor; they interpolate between the compact case where $\Delta = \emptyset$ and the pair $(Z, \emptyset) = Z$ has no orbifold structure, and the *open*, or *purely-logarithmic case* where $c_j = 1$ for all j , and we identify (Z, Δ) with $Z \setminus \text{Supp}(\Delta)$.

When Z is smooth and the support $\text{Supp}(\Delta) := \cup D_j$ of Δ has normal crossings singularities, we say that (Z, Δ) is *smooth*. When all multiplicities m_j are integral or $+\infty$, we say that the orbifold pair (Z, Δ) is *integral*, and when every m_j is finite it may be thought of as a virtual ramified cover of Z ramifying at order m_j over each of the D_j 's.

Consider a fibration $f: X \rightarrow Z$ between normal connected complex projective varieties. In general, the geometric invariants (such as $\pi_1(X), \kappa(X), \dots$) of X do not coincide with the ‘sum’ of those of the base (Z) and of the generic fiber (X_η) of f . Replacing Z by the ‘orbifold base’ (Z, Δ_f) of f , which encodes the multiple fibers of f , leads in some favorable cases to such an additivity (on suitable birational models at least).

Definition 2.4 (Orbifold base of a fibration). Let $f: (X, \Delta) \rightarrow Z$ be a fibration $X \rightarrow Z$ as above and let Δ be an orbifold divisor on X . We shall define the *orbifold base* (Z, Δ_f) of (f, Δ) as follows: to each irreducible Weil divisor $D \subset Z$ we assign the multiplicity $m_{(f, \Delta)}(D) := \inf_k \{t_k \cdot m_\Delta(F_k)\}$, where the scheme theoretic fiber of D is $f^*(D) = \sum_k t_k \cdot F_k + R$, R is an f -exceptional divisor of X with $f(R) \subsetneq D$ and F_k are the irreducible divisors of X which map surjectively to D via f .

Remark 2.5. Note that the integers t_k are well-defined, even if X is only assumed to be normal.

Let (Z, Δ) be an orbifold pair. Assume that $K_Z + \Delta$ is \mathbb{Q} -Cartier (this is the case if (Z, Δ) is smooth, for example): we will call it the *canonical bundle* of (Z, Δ) . Similarly we will denote by the *canonical dimension* of (Z, Δ) the Kodaira dimension of $K_Z + \Delta$ i.e. $\kappa(Z, \Delta) := \kappa(Z, \mathcal{O}_Z(K_Z + \Delta))$. Finally, we say that the orbifold (Z, Δ) is of *general type* if $\kappa(Z, \Delta) = \dim(Z)$.

Definition 2.6. A fibration $f: X \rightarrow Z$ is said to be of *general type* if (Z, Δ_f) of general type.

The idea of Campana was that in order to characterize varieties that have a potentially dense set of rational points it was not sufficient to exclude the presence of étale covers that dominate varieties of general type. One would need to exclude the presence of every fibration of general type in the above sense. This turns out to be equivalent to the specialness condition of Definition 2.3 as proven by Campana.

Theorem 2.7 (see [Cam04, Theorem 2.27]). *A variety X is special if and only if it has no fibrations of general type.*

2.3. Orbifold Morphisms

We recall here the main definition of orbifold morphism, following [Cam11, Definition 2.3], in the special case in which the source is a curve.

Definition 2.8. Let (X, Δ_X) and $(\mathcal{C}, \Delta_{\mathcal{C}})$ two orbifold pairs, with X and \mathcal{C} normal projective varieties defined over a field k , \mathcal{C} a curve, and $\Delta_X, \Delta_{\mathcal{C}}$ the two orbifold divisor of the form

$$\Delta_X = \sum_{i=1}^r \left(1 - \frac{1}{m_i}\right) \Delta_i \quad \Delta_{\mathcal{C}} = \sum_{j=1}^s \left(1 - \frac{1}{n_j}\right) P_j$$

For every prime divisor D of X we denote by $m_X(D)$ its multiplicity, i.e. the number m_i if $D = \Delta_i$ or 0 otherwise. For every point $Q \in \mathcal{C}$ we similarly define $n_{\mathcal{C}}(Q)$. A morphism $f : \mathcal{C} \rightarrow X$ is an *orbifold morphism*, denoted by $f : (\mathcal{C}, \Delta_{\mathcal{C}}) \rightarrow (X, \Delta_X)$ if

- (1) $f(\mathcal{C})$ is *not* contained in Δ_X ;
- (2) for every prime divisor D of X , if $f^*D = \sum_k n_k Q_k$, one has $n_k \cdot n_{\mathcal{C}}(Q_k) \geq m_X(D)$.

We recall the following definition introduced by Demailly in [Dem97] in the compact case, while the logarithmic and orbifold analogue were introduced in [Che04] and [Rou10] respectively.

Definition 2.9. Let (X, D) be a pair of a non-singular projective variety X defined over k and let D be a normal crossing divisor on X . We say that (X, D) is *algebraically hyperbolic* if there exists a ample line bundle \mathcal{L} on X and a positive constant α such that for every non-singular projective curve \mathcal{C} and every morphism $\varphi : \mathcal{C} \rightarrow X$ the following holds:

$$(2.1) \quad \deg \varphi^* \mathcal{L} \leq \alpha \cdot (2g(\mathcal{C}) - 2 + N_{\varphi}^{[1]}(D)).$$

where $N_{\varphi}^{[1]}(D)$ is the cardinality of the *support* of $\varphi^*(D)$. We say that (X, D) is *pseudo algebraically hyperbolic* if there exists a proper closed subvariety Z of X such that (2.1) holds for every morphism $\varphi : \mathcal{C} \rightarrow X$ such that $\varphi(\mathcal{C}) \not\subseteq Z$.

The notion of *pseudo-algebraic hyperbolicity* was defined first in [vBJK19, JX19]. In Lang's terminology [Lan86], the notion is the ‘‘pseudofication’’ of the notion of algebraic hyperbolicity.

Remark 2.10. Note that the degree $\deg \varphi^* \mathcal{L}$, is a Weil Height for φ viewed as a point in the function field $k(\mathcal{C})$, with respect to the ample line bundle \mathcal{L} . In the next sections, we refer to it both as a degree and as the height $h_{\mathcal{L}}(\varphi)$.

Definition 2.9 can be extended to the orbifold setting as follows.

Definition 2.11. Let (X, Δ) be an orbifold; we say that (X, Δ) is *algebraically hyperbolic* if there exists a ample line bundle \mathcal{L} on X and a positive constant α such that for every non-singular projective curve \mathcal{C} and every *orbifold* morphism $\psi : (\mathcal{C}, \Delta_{\mathcal{C}}) \rightarrow (X, \Delta)$ the following holds:

$$(2.2) \quad \deg \psi^* \mathcal{L} \leq \alpha \cdot (2g(\mathcal{C}) - 2 + \deg \Delta_{\mathcal{C}}).$$

We say that (X, Δ) is *pseudo algebraically hyperbolic* if there exists a proper closed subvariety Z of X such that (2.2) holds for every orbifold morphism $\psi : (\mathcal{C}, \Delta_{\mathcal{C}}) \rightarrow (X, \Delta)$ such that $\psi(\mathcal{C}) \not\subseteq Z$.

3. GENERALIZED LANG CONJECTURES

In this section we propose a generalization of the conjectures of Lang and Vojta compatible with Campana's dichotomy between special and nonspecial.

3.1. Arithmetic and Analytic Exceptional sets

In [Lan86], Lang introduced the following exceptional sets.

Definition 3.1. Let X be a projective variety defined over a field k .

- (1) If k is a number field, then the *Diophantine exceptional set* $\text{Exc}_{\text{dio}}(X)$ is the smallest Zariski-closed subset Z of X such that $(X \setminus Z)(L)$ is finite for all finite extensions $L \supset k$.
- (2) If $k = \mathbb{C}$ then the *holomorphic exceptional set* $\text{Exc}_{\text{hol}}(X)$ is the Zariski closure of the union of all entire curves i.e. images of non-constant holomorphic maps $f : \mathbb{C} \rightarrow X$.

Lang has conjectured [Lan86] that X is of general type if and only if both these exceptional sets are proper subsets of X . Moreover he conjectured that $\text{Exc}_{\text{dio}}(X) = \text{Exc}_{\text{hol}}(X)$.

Given Campana’s notion of special variety, it is natural to try to extend the notion of exceptional sets to *nonspecial* varieties in the sense of Campana [Cam04].

The starting point is the following conjecture, formulated by Campana [Cam04, Conjecture 9.2 and Conjecture 9.20].

Conjecture 1. *Let X be a projective variety defined over a field k .*

- (1) *If k is a number field, then $X(L)$ is not Zariski dense for all finite extensions $L \supset k$ if and only if X is not special*
- (2) *If $k = \mathbb{C}$ then $f(\mathbb{C})$ is not Zariski dense for all entire curves $f : \mathbb{C} \rightarrow X$ if and only if X is not special.*

When one consider nonspecial varieties, it is easy to see that there are examples of nonspecial varieties where the two exceptional sets in Definition 3.1 are the entire variety. As an example, consider the nonspecial variety $X = \mathcal{C} \times \mathbb{P}^1$ defined over a number field k , where \mathcal{C} is a smooth projective curve of genus ≥ 2 : in this case for every rational point P in $\mathcal{C}(L)$ the curve $\{P\} \times \mathbb{P}^1$ is in $\text{Exc}_{\text{dio}}(X)$. Therefore since $X(\bar{k})$ is dense in X , Lang’s Diophantine exceptional set coincide with X . Similarly one can show that $\text{Exc}_{\text{hol}}(X) = X$. We suggest that, in order to define meaningful exceptional sets, one should consider the projectivized tangent bundle $X_1 := \mathbb{P}(T_X)$.

Definition 3.2.

- (1) If k is a number field, the Diophantine exceptional set $\text{Exc}_{\text{dio}}^1(X)$ is the smallest Zariski-closed subset Z of X_1 such that for or all finite extensions $L \supset k$, $\mathbb{P}(T_{Y_L}) \subset Z$, where Y_L is the Zariski closure of $X(L)$.
- (2) If $k = \mathbb{C}$, the holomorphic exceptional set $\text{Exc}_{\text{hol}}^1(X)$ is the Zariski closure of the union of all entire curves $g : \mathbb{C} \rightarrow X_1$ obtained as liftings of entire curves $f : \mathbb{C} \rightarrow X$.

The main motivation behind Definition 3.2 is Campana’s Core construction in [Cam04, Section 3]. Given a smooth projective variety X there is a functorial fibration $c_X : X \rightarrow C(X)$, called the *core* of X such that the fibers of c_X are special varieties and the base $C(X)$ is either a point or an orbifold of general type. The idea behind considering X_1 as the natural space where the exceptional set live, is that the core of X identifies the “special direction” in X_1 and therefore, assuming that $C(X)$ has positive dimension, i.e. X is nonspecial, this exceptional set should not be the whole X_1 .

Therefore we propose the following generalization of Lang conjecture for nonspecial varieties.

Conjecture 2. *Let X be a projective variety defined over a field k .*

- (1) *If k is a number field then $\text{Exc}_{\text{dio}}^1(X) \neq X_1$ if and only if X is not special*

(2) If $k = \mathbb{C}$ then $\text{Exc}_{\text{hol}}^1(X) \neq X_1$ if and only if X is not special.

In the example above $X = \mathcal{C} \times \mathbb{P}^1$ with \mathcal{C} an hyperbolic curve, we see that for every number field L the closed subvariety Y_L , the closure of $X(L)$, is the union of finitely many rational curves, corresponding to the fibers of $\text{pr}_1 = c_X$ over the L -rational points of \mathcal{C} . Thus $\text{Exc}_{\text{dio}}^1(X) = \mathbb{P} \text{pr}_2^*(T_{\mathbb{P}^1})$ and in particular $\text{Exc}_{\text{dio}}^1(X) \neq X_1$.

Comparing Conjecture 1 and Conjecture 2 suggest interesting questions. In particular a strong uniform degeneracy statement in X_1 should imply a non density statement in X .

Conjecture 3. *Let X be a projective variety defined over a field k .*

(1) *If k is a number field and $\text{Exc}_{\text{dio}}^1(X) \neq X_1$ then $X(k)$ is not potentially dense.*

(2) *If $k = \mathbb{C}$ and $\text{Exc}_{\text{hol}}^1(X) \neq X_1$ then entire curves $f : \mathbb{C} \rightarrow X$ are algebraically degenerate i.e. the images $f(\mathbb{C})$ are not Zariski dense.*

Interestingly, some examples of this phenomenon have already been proved in the foliated setting. If one considers a complex projective manifold X equipped with a holomorphic (singular) foliation \mathcal{F} such that all entire curves $f : \mathbb{C} \rightarrow X$ are tangent to this foliation, then it is proved in [McQ98] that if X is a surface of general type and \mathcal{F} is a foliation by curves, then entire curves are indeed algebraically degenerate. More recently, the same conclusion is shown in [BRT19] if \mathcal{F} is a transversely hyperbolic foliation of codimension 1 and X an arbitrary complex projective manifold X (not necessarily of general type!).

3.2. Function Fields

Given the well-known analogy between number fields and function fields, we formulate the above conjectures in the function field setting. Let κ be an algebraically closed field of characteristic 0, let \mathcal{C} be a smooth complex projective curve and let $\kappa(\mathcal{C})$ be its function field over κ . Let X be a proper variety defined over $\kappa(\mathcal{C})$ and let $g : \mathcal{X} \rightarrow \mathcal{C}$ be the fibration associated to a proper model of X over \mathcal{C} . In this setting, $\kappa(\mathcal{C})$ -rational points correspond to sections $s : \mathcal{C} \rightarrow \mathcal{X}$ of g and points defined over a finite extension $\kappa(\mathcal{C}') \supset \kappa(\mathcal{C})$ correspond to section of the base change $\mathcal{X}' := \mathcal{X} \times_{\mathcal{C}} \mathcal{C}' \rightarrow \mathcal{C}'$ via a ramified cover $\mathcal{C}' \rightarrow \mathcal{C}$.

In this setting, Lang conjecture can be formulated in the following way (see [Lan86, Historical appendix: algebraic families] for an historical discussion).

Conjecture 4. *Let X be non-isotrivial. Then X is of general type if and only if for all finite extensions $\kappa(\mathcal{C}') \supset \kappa(\mathcal{C})$ there is a proper algebraic subset $Z \subset X$ such that $X \setminus Z$ contains only finitely many $\kappa(\mathcal{C}')$ -points.*

Special cases of Conjecture 5 have been proved in the literature, e.g. [Nog82].

Similarly, the generalization of Conjecture 1 is formulated as follows.

Conjecture 5. *Let X be non-isotrivial. Then X is not special if and only if for all finite extensions $\kappa(\mathcal{C}') \supset \kappa(\mathcal{C})$, $\kappa(\mathcal{C}')$ -points in X are not Zariski dense.*

Lang's stronger conjecture predicts that the closed subset Z , the *exceptional set*, should be independent of the field of definition of the points. Like in the number field case, this cannot be

translated directly to nonspecial varieties. Instead we propose the following analogue of Conjecture 2 in the function field setting.

Conjecture 6. *X is not special if and only if for all finite extensions $\kappa(\mathcal{C}') \supset \kappa(\mathcal{C})$ there is a proper algebraic subset $Z \subset X_1$ such that sections $s : \mathcal{C}' \rightarrow \mathcal{X}'$ whose liftings $s_1 : \mathcal{C}' \rightarrow \mathcal{X}_1$ are not contained in Z are finite.*

In the number field case, conjectures of Vojta (see [Voj87, Conjecture 3.4.3] and [Voj11, Conjecture 24.3]) predict a height bound for rational points in varieties of general type outside of the exceptional locus, which implies a non-density statement.

In the function field setting, an height bound is also expected, which is the content of the Lang-Vojta conjecture.

Conjecture 7. *If X is of general type and \mathcal{L} is an ample line bundle on X , then there exist a proper algebraic subset $Z \subsetneq X$ and a positive constant α such that sections $s : \mathcal{C} \rightarrow \mathcal{X}$ not contained in Z (the model of Z) satisfy the inequality $\deg s^*\mathcal{L} \leq \alpha(2g(\mathcal{C}) - 2)$, with \mathcal{L} a model of L .*

The isotrivial case of Conjecture 7 is known for subvarieties of abelian varieties [Yam15]. For certain cases of Conjecture 7 in the logarithmic setting see [CZ08, CZ13, Tur17, ADT18, CT19].

One should notice that for function fields, the height bound predicted by Conjecture 7 can only imply a non-density result when the variety is not isotrivial. We note that in the isotrivial case, one can expect that a natural generalization of Conjecture 7 predicts the following

Conjecture 8. *If (X, Δ) is an orbifold of general type then (X, Δ) is pseudo algebraically hyperbolic as in Definition 2.11.*

Motivated by Campana’s core construction, as introduced above, we propose the following extension of Conjecture 7 to nonspecial varieties.

Conjecture 9. *If X is not special, then there are a rational dominant map $\pi : X \rightarrow Y$ with $\dim Y > 0$, a proper algebraic subset $Z \subsetneq X_1$, a positive constant α and an ample line bundle L on Y such that sections $s : \mathcal{C} \rightarrow \mathcal{X}$, whose liftings $s_1 : \mathcal{C} \rightarrow \mathcal{X}_1$ are not contained in Z , satisfy the inequality $\deg(\pi \circ s)^*\mathcal{L} \leq \alpha(2g(\mathcal{C}) - 2)$.*

Finally, in the isotrivial case, a natural generalization of Conjecture 9 to the logarithmic case predicts the following

Conjecture 10. *If (X, D) is not special, then there are a rational dominant map $\pi : (X, D) \rightarrow (Y, D')$ with $\dim Y > 0$, a proper algebraic subset $Z \subsetneq X_1$, a positive constant α and an ample line bundle L on Y such that sections $s : \mathcal{C} \rightarrow X$, whose liftings $s_1 : \mathcal{C} \rightarrow X_1$ are not contained in Z , satisfy the inequality $\deg(\pi \circ s)^*\mathcal{L} \leq \alpha(2g(\mathcal{C}) - 2 + N_{\pi \circ s}^{[1]}(D'))$.*

4. SPECIAL VS WEAKLY SPECIAL

In this section we construct a family of examples of varieties that are weakly special in the sense of Abramovich and Colliot-Thélène but are not special in the sense of Definition 2.3.

4.1. *Weakly special varieties*

We first recall the definition of *weakly special* variety (the terminology is due to Campana).

Definition 4.1. A projective variety X defined over a field k is said to be *weakly special* if there are no finite \bar{k} étale covers $u: X' \rightarrow X$ admitting a dominant rational map $f': X' \rightarrow Z'$ to a positive dimensional variety Z' of general type.

Abramovich and Colliot-Thélène conjectured that weak specialness characterizes potential density of rational points for varieties defined over number fields.

Conjecture 11 (see [HT00, Conjecture 1.2]). *Let X be a projective variety defined over a number field. Then, the set of rational points on X is potentially dense if and only if X is weakly special.*

Conjecture 11 is still open: in fact there is no example of a weakly special variety whose rational points are not potentially dense. On the other hand the conjecture should imply the following statements for function fields and entire curves.

Conjecture 12. *Let X be a projective variety defined over \mathbb{C} . Then, X is weakly special if and only if:*

- (1) *there exists an entire curve $\mathbb{C} \rightarrow X$ with Zariski dense image.*
- (2) *there are no dominant map $\pi: X \rightarrow Y$, an ample line bundle \mathcal{L} , a positive constant α and a proper closed subset Y_{exc} such that for every smooth integral curve \mathcal{C} and morphism $s: \mathcal{C} \rightarrow X$ such that $s(\mathcal{C})$ is not contained in Y_{exc} the following holds:*

$$\deg(\pi \circ s)^* \mathcal{L} \leq \alpha (2g(\mathcal{C}) - 2).$$

As seen in Section 2, if a variety X is special then X is weakly special. However, the two notions are equivalent only for curves and surfaces. We construct here examples of 3-dimensional projective varieties which are weakly special, but not special. In Section 6 we will show that these examples contradict Conjecture 12.

4.2. *Examples of weakly special, but not special threefolds.*

The construction below is a slight extension of the construction in [BT04] and was explained to us by F. Campana. We shall construct simply-connected smooth projective threefolds X having no rational fibrations onto varieties of general type, but having equidimensional (elliptic) fibrations of (orbifold) general type $F: X \rightarrow S$ onto smooth surfaces S . These are thus examples of weakly special but not special varieties in the lowest dimension where the two notion do not agree.

Theorem 4.2. *There exist surfaces T, S and fibrations $f: T \rightarrow \mathbb{P}^1$ and $g: S \rightarrow \mathbb{P}^1$ such that:*

- (1) *the first fibration $f: T \rightarrow \mathbb{P}^1$ has a single multiple fiber $f^{-1}(0) =: m \cdot T_0$, with T_0 a smooth elliptic curve and $m \in \mathbb{Z}_{>1}$, and another (singular) simply-connected fiber;*
- (2) *S is a smooth surface and the fibration $g: S \rightarrow \mathbb{P}^1$ has a smooth fiber $S_0 := g^{-1}(0)$ such that:*
 - (a) *the surface S is not of general type but the orbifold surface*

$$(S, \Delta) := (S, (1 - 1/m) \cdot S_0)$$

is of general type;

(b) the complement of S_0 in S is simply connected.

Furthermore, let X be the total spaces of the natural (orbifold) elliptic fibrations with equidimensional fibers defined by f and g , i.e.

$$X := (S \times_{\mathbb{P}^1} T) \xrightarrow{F} (S, \Delta).$$

Then X is a weakly special projective smooth threefold that is not special.

Proof. The existence is proven in the examples below, so we focus on the last part of the statement. We first notice that the orbifold base of the fibration F is indeed Δ : by construction, the only multiple fibers lie above S_0 , and $F^*S_0 = m \cdot (S_0 \times T_0)$. Since by assumption (S, Δ) is a surface of general type, it follows immediately that X is not of special type.

We now prove that X is weakly special. A key feature of X in this direction is simple-connectedness. Let $D := F^{-1}(S_0)$; then, the fibration $F: (X \setminus D) \rightarrow (S \setminus S_0)$ is a fibration without multiple fibers (since $f: (T \setminus T_0) \rightarrow \mathbb{P}^1 \setminus \{0\}$ has no multiple fibers), and with a simply-connected fiber. This implies that $F_*: \pi_1(X \setminus D) \rightarrow \pi_1(S \setminus S_0)$ is an isomorphism. Using our assumption, the group $\pi_1(X \setminus D)$ is hence trivial. This implies that X is simply-connected since the natural map $\pi_1(X \setminus D) \rightarrow \pi_1(X)$ is surjective.

As a consequence, to prove that X is weakly special, it suffices to show that no fibration $h: X \rightarrow Z$ exists, with Z of general type and of dimension d , with $0 < d \leq 3$, since X does not admit any non-trivial étale cover. Assume by contradiction that such an h exists. Then $d < 3$, because X is not of general type, since it is an elliptic fibration over S ; hence $d = 1, 2$. Note that, since X is simply connected, Z has to be simply connected, thus $d > 1$, since the only simply-connected curve is \mathbb{P}^1 , which is not of general type. We are reduced to the case in which $d = 2$, and $h \neq F$, since by assumption S is not of general type. This shows that Z is covered by the image of the fibers of F , i.e. the elliptic curves $h(X_t)$ with $t \in S$. This contradicts our assumption that Z is a surface of general type. \square

Example 4.3 (First fibration). In order to construct $f: T \rightarrow \mathbb{P}^1$, we consider first a fibration $f': T' \rightarrow \mathbb{P}^1$ of an elliptic surface having no multiple fiber, a smooth fiber $T'_0 := (f')^{-1}(0)$ and a (singular) simply-connected fiber, and moreover such that $p_g(T') := h^{0,2}(T') = 0$. Then T' is simply connected. Let $m > 1$ be an integer. The Kodaira logarithmic transform $f: T \rightarrow \mathbb{P}^1$ of order m on the fiber T'_0 of f' is a new elliptic fibration which replaces T'_0 by a smooth multiple fiber $m \cdot T_0$ of multiplicity m over $0 \in \mathbb{P}^1$, leaving the complements $T' \setminus T'_0$ and $T \setminus m \cdot T_0$ isomorphic as elliptic fibrations over $\mathbb{P}^1 \setminus \{0\}$ (see for example [BT04, Section 3] and for more details [FM94, Section 1.6]). Moreover $p_g(T) = p_g(T') = 0$, and T is still simply connected since f still has a simply connected fiber. This implies that $b_1(T) = 0$ is even and T is Kähler; since $p_g(T) = 0$, we conclude that T is projective (we refer to [FM94, Section 1.6] for the details).

In the original construction of [BT04], S was chosen so that $\kappa(S) = 1$. We shall see that a small variation of the construction permits to chose for (S, Δ_F) a suitable blow-up of any smooth projective surface S' with $\kappa(S') \in \{-\infty, 0, 1\}$ and Δ_F the strict transform of a suitable ample orbifold \mathbb{Q} -divisor on S' .

Example 4.4 (Second fibration). In order to construct $g: S \rightarrow \mathbb{P}^1$ (given m), we consider a simply-connected smooth projective surface S' that is not of general type, together with an ample and smooth divisor C' , which is a member of a pencil C'_t of divisors on S' (whose generic member C'' meets C' transversally at C'^2 distinct points), such that the complete linear system $\mathcal{O}_{S'}(C')$ is base-point free, and such that $K_{S'} + (1 - 1/m) \cdot C'$ is a big divisor on S' .

For example, one can chose $S' = \mathbb{P}^2$, together with C' a smooth curve of degree $d \geq 4$ if $m \geq 5$, and $d \geq 7$ if $m \geq 2$.

Now choose a second generic member C'' of the linear system $|C'|$ meeting transversally C' at $(C')^2$ distinct points. Let $\sigma: S \rightarrow S'$ be the blow-up of S' at these $(C')^2$ points, let Δ be the strict transform of $\Delta' := (1 - 1/m) \cdot C'$ on S , and let E the exceptional divisor of σ . Then the divisor

$$K_S + \Delta = (\sigma^* K_{S'} + E) + (\sigma^* \Delta' - (1 - 1/m) \cdot E) = \sigma^*(K_{S'} + \Delta') + (1/m) \cdot E$$

is big since σ is birational and all the varieties are projective; this implies that the orbifold surface (S, Δ) is of general type.

Let $k > 0$ be the order of divisibility of $[C']$ in $\text{Pic}(S')$. By (a version of) Lefschetz theorem, $\pi_1(S' \setminus C')$ is the cyclic group of order k generated by a small loop around C' . But blowing-up a point a on C' makes this loop become trivial in $\pi_1(\mathbb{C} = E_a \setminus \{a\})$, where $E_a \cong \mathbb{P}^1$ is the exceptional divisor of the blow-up over a , and $a \in C' \cap C'' \subset S'$. This loop thus becomes homotopically trivial in $(S \setminus S_0)$, which is thus simply-connected.

Remark 4.5. The smoothness of C' is not necessary. One may just assume that C' is nodal and that C'' meets C' at smooth points of C' .

5. DEGENERACY RESULTS FOR SURFACES

In this section we extend results of Corvaja and Zannier in [CZ04] for number fields to the orbifold setting, both over function fields and in Nevanlinna Theory. We prove hyperbolicity and degeneracy results for a class of surfaces as a combination of ideas of Corvaja and Zannier together with the recent method introduced by Ru-Vojta in [RV].

In particular we recover the function field and analytic analogues of [CZ04, Main Theorem] in the logarithmic setting. We stress that, even if the arithmetic result for orbifold rational points seems at the moment out of reach, our results give evidence for the arithmetic part of Conjecture 1. Finally, all these results will be applied in Section 6 to the nonspecial threefold X_m defined in Section 4.

5.1. Function Fields

In this section we let κ be an algebraically closed field of characteristic zero and we assume all the varieties defined over κ . The main theorem of this section will be a consequence of the following statement, whose proof is contained in Section 7 (see Theorem 7.6). We start by recalling the definition of the β invariant (we refer to [RV] for discussion and properties).

Definition 5.1. Let X be a complete variety, let \mathcal{L} be a big line bundle on X and let D be a nonzero effective Cartier divisor on X . We define

$$\beta(\mathcal{L}, D) = \liminf_{N \rightarrow \infty} \frac{\sum_{m \geq 1} h^0(X, \mathcal{L}^N(-mD))}{N h^0(X, \mathcal{L}^N)}.$$

Using the constant β we can reformulate the main theorem of [RV] in the function field case. However, for our applications we need to obtain an explicit dependence of the constants on the Euler characteristic of the curve. This is obtained in the following theorem, which we state in the constant case, by use of the more explicit [Wan04, Theorem 1] instead of the function field analogue of the Schmidt Subspace Theorem [Wan04, Main Theorem].

Theorem 5.2. *Let $X \subset \mathbb{P}^m$ be a projective variety over κ of dimension n , let D_1, \dots, D_q be effective Cartier divisors intersecting properly on X , and let \mathcal{L} be a big line sheaf. Let \mathcal{C} be a smooth projective curve over κ , let S be a finite set of points on \mathcal{C} and let $K = \kappa(\mathcal{C})$ be the function field of \mathcal{C} . Then for any $\epsilon > 0$, there exist constants c_1 and c_2 , independent of the curve \mathcal{C} and the set S , such that for any map $x = [x_0 : \dots : x_m] : \mathcal{C} \rightarrow X$, where $x_i \in K$, outside the augmented base locus of \mathcal{L} we have either*

$$\sum_{i=1}^q \beta(\mathcal{L}, D_i) m_{D_i, S}(x) \leq (1 + \epsilon) h_{\mathcal{L}}(x) + c_1 \max\{1, 2g(\mathcal{C}) - 2 + |S|\},$$

or the image of x is contained in a hypersurface (over κ) in \mathbb{P}^m of degree at most c_2 .

Remark 5.3.

- (1) In the case where X is nonsingular, the condition on the proper intersection is equivalent to general position. We refer to [RV, Definition 2.1] for precise statements and properties.
- (2) Even if Theorem 5.2 is stated in the split case we note that our proof carries over the non split case almost verbatim. We focus here only on the split case since it is the one relevant for our applications.
- (3) The constants c_1 and c_2 appearing in Theorem 5.2 can be effectively computed given $X, D_i, \mathcal{L}, \epsilon$ so in particular the algebraic hyperbolicity bounds can be made effective in the same way.

The proof of Theorem 5.2 is included in Section 7 as Theorem 7.6, and we refer to Subsection 7.2 for the definitions of all the quantities involved. For our application we will use the following corollary in dimension 2, whose proof is also included in Section 7 (see Corollary 7.7).

Corollary 5.4. *In the previous setting, if X has dimension 2, then for any $\epsilon > 0$, there exist constants c_1 and c_3 independent of the curve \mathcal{C} and the set S such that for any K -point $x = [x_0 : \dots : x_m] : \mathcal{C} \rightarrow X$, with $x_i \in K$, we have that, either $\deg x(\mathcal{C}) \leq c_3$, or*

$$\sum_{i=1}^q \beta(\mathcal{L}, D_i) m_{D_i, S}(x) \leq (1 + \epsilon) h_{\mathcal{L}}(x) + c_1 \max\{1, 2g(\mathcal{C}) - 2 + |S|\}.$$

We now state the main result of this section in the logarithmic case.

Theorem 5.5 (cfr [CZ04, Main Theorem]). *Let $X \subset \mathbb{P}^m$ be a smooth projective surface and $D = D_1 + \dots + D_r$ be a divisor with $r \geq 2$, both defined over κ , such that*

- (1) No three of the components D_i meet at a point;
- (2) There exists a choice of positive integers p_i such that
 - the divisor $D_p := p_1 D_1 + p_2 D_2 + \cdots + p_r D_r$ is ample;
 - The following inequality holds:

$$2D_p^2 \xi_i > (D_p \cdot D_i) \xi_i^2 + 3D_p^2 p_i,$$

for every $i = 1, \dots, r$ where ξ_i is the minimal positive solution of the equation $D_i^2 x^2 - 2(D_p \cdot D_i)x + D_p^2 = 0$.

Then, (X, D) is pseudo algebraically hyperbolic (see Definition 2.9).

We will prove Theorem 5.5 as an application of Theorem 5.2 thus diverging from the original proof that uses the Schmidt's Subspace Theorem. Nevertheless we will use some of the techniques of [CZ04] in computing the constants β .

Lemma 5.6. *In the same setting as Theorem 5.5, for every $i = 1, \dots, r$, $\beta_i = \beta(D_p, D_i) > p_i$.*

Proof. By the Riemann-Roch Theorem, for N large enough we have that $2h^0(ND_p) = D_p^2 N^2 + O(N)$. In order to compute β_i , we consider the divisor $ND_p - mD_i$ which is effective if $m \leq Np_i$.

By the same computation as in [CZ04] (in particular see pages 718–719) we get

$$(5.1) \quad \sum_{m=0}^{\xi_i N} h^0(ND_p - mD_i) \geq N^3 \left(\frac{\xi_i^2 (D_p \cdot D_i)}{2} - \frac{\xi_i^3 D_i^2}{3} \right) + O(N^2).$$

Recall that by definition of ξ_i , we have $D_i^2 \xi_i^3 = 2(D_p \cdot D_i) \xi_i^2 - D_p^2 \xi_i = 0$. Using this in equation 5.1 we obtain

$$\begin{aligned} 2 \sum_{m=0}^{\xi_i N} h^0(NA - mD_i) &\geq 2N^3 \left(\frac{\xi_i^2 (D_p \cdot D_i)}{2} - \frac{2\xi_i^2 (D_p \cdot D_i)}{3} + \frac{D_p^2 \xi_i}{3} \right) + O(N^2) \\ &= \left(\frac{2}{3} \xi_i D_p^2 - \frac{1}{3} (D_p \cdot D_i) \xi_i^2 \right) N^3 + O(N^2). \end{aligned}$$

This implies that

$$\beta_i = \frac{\frac{2}{3} \xi_i D_p^2 - \frac{1}{3} (D_p \cdot D_i) \xi_i^2}{D_p^2} > p_i.$$

□

We also include an easy lemma which provides a useful trick to get a lower bound of the height of a point with respect to D_p .

Lemma 5.7. *In the same settings as Theorem 5.5, let $\varepsilon = \min\{(\beta - p_i)/p_i\}$, then for every $\varphi : \mathcal{C} \rightarrow X$,*

$$\sum_{i=1}^r \beta_i h_{D_i}(\varphi) \geq (1 + \varepsilon) h_{D_p}(\varphi)$$

Proof. We have that for every i

$$\beta_i h_{D_i}(\varphi) = \left(\frac{\beta_i - p_i}{p_i} + 1 \right) p_i h_{D_i}(\varphi)$$

Then, summing over $i = 1, \dots, r$ we get

$$\sum_{i=1}^r \beta_i h_{D_i}(\varphi) = \sum_{i=1}^r \left(\frac{\beta_i - p_i}{p_i} + 1 \right) p_i h_{D_i}(\varphi) \geq (1 + \varepsilon) h_{D_p} \varphi$$

□

Proof of Theorem 5.5. We will prove that there exists a constant $\alpha > 0$ such that for every morphism $\varphi : \mathcal{C} \rightarrow X$ with $\varphi(\mathcal{C}) \not\subseteq D$ one has

$$(5.2) \quad \deg \varphi^* D_p \leq \alpha \cdot \max\{1, 2g(\mathcal{C}) - 2 + N_\varphi^{[1]}(D_p)\}.$$

We note that such bound implies that (X, D) is pseudo algebraically hyperbolic in the sense of Definition 2.9 provided that the union of images of curves for which $2g(\mathcal{C}) - 2 + N_\varphi^{[1]}(D_p) \leq 0$ is a proper closed subset of X . We will first prove the bound (5.2) and then show that the exceptional set is indeed a proper and closed subset. First we note that condition (1) implies that D_1, \dots, D_r are in general position. We fix

$$\varepsilon = \min_i \left\{ \frac{\beta_i - p_i}{p_i} \right\}$$

which is positive by Lemma 5.6. Let $\varphi : \mathcal{C} \rightarrow X$ be a morphism such that $\varphi(\mathcal{C}) \not\subseteq D$. By Corollary 5.4, applied with $\mathcal{L} = \mathcal{O}_X(D_p)$, there exist constant c_1 and c_3 depending only on X and ε (independent of \mathcal{C} and φ) such that either $\varphi(\mathcal{C})$ has degree bounded by c_3 or

$$(5.3) \quad \sum_{i=1}^r \beta_i m_{D_i, S}(\varphi) \leq (1 + \frac{\varepsilon}{2}) h_{D_p}(\varphi) + c_2 \max\{1, 2g(\mathcal{C}) - 2 + |S|\},$$

where $S = \text{supp}(\varphi^* D)$. In the first case we are done. In the latter, since the support of $\varphi^* D$ is contained in S , $m_{D_i, S}(\varphi) = h_{D_i}(\varphi) + O(1)$.

Lemma 5.7 gives the lower bound

$$\sum_{i=1}^r \beta_i m_{D_i, S}(\varphi) > (1 + \varepsilon) h_{D_p}(\varphi).$$

Together with (5.3), this implies that there exists a constant c'_2 such that

$$\frac{\varepsilon}{2} h_{D_p}(\varphi) < c'_2 \max\{1, 2g(\mathcal{C}) - 2 + |S|\}.$$

Hence,

$$\deg \varphi^* D_P = h_{D_p}(\varphi) < 2c'_2 \varepsilon^{-1} \max\{1, 2g(\mathcal{C}) - 2 + N_\varphi^{[1]}(D_p)\}.$$

To finish the proof we need to show that there exists a closed subvariety Z that contains all images $\varphi(\mathcal{C})$ when $2g(\mathcal{C}) - 2 + N_\varphi^{[1]}(D_p)$ is not positive. First of all, we can reduce to the case in which $\kappa = \mathbb{C}$: it is enough to notice that given (X, D) defined over κ there exists a field κ_0 that is an algebraically closed subfield of κ , which is finitely generated over $\overline{\mathbb{Q}}$, and such that (X, D) has a model over κ_0 . For such κ_0 there exists an embedding $\kappa_0 \rightarrow \mathbb{C}$, and therefore we can reduce the problem to the case in which (X, D) is defined over \mathbb{C} .

In this case, we claim that the exceptional set Z can be chosen to be the exceptional set in [Lev09, Theorem 8.3 B]. In fact, the same strategy of Levin applies to the setting of our theorem (in particular the proof of [CZ04, Main Theorem] shows that D_p is *large* in the sense of Levin) and therefore there exists a proper closed subvariety $Z \subset X$ that contains all the images of entire curve $\mathbb{C} \rightarrow X$. To conclude, it is sufficient to notice that for every curve \mathcal{C} such that $2g(\mathcal{C}) - 2 + N_\varphi^{[1]}(D_p) \leq 0$ there exists a non trivial holomorphic map $\mathbb{C} \rightarrow \mathcal{C} \setminus \text{supp } \varphi^* D_p$. In particular this implies that $\varphi(\mathcal{C})$ has to be contained in Z as wanted. \square

Next we show that Theorem 5.5 leads to a statement for orbifold morphisms, in particular we obtain the conclusion of Conjecture 8.

Corollary 5.8. *Let (X, D) as in Theorem 5.5. Let Δ be the \mathbb{Q} -divisor defined as*

$$\Delta = \sum_{i=1}^r \left(1 - \frac{1}{m_i}\right) D_i,$$

for some integers $m_i \geq 1$. Then, there exists a positive integer $m > 0$ such that, if $m_i > m$ for every i , (X, Δ) is pseudo algebraically hyperbolic (see Definition 2.11).

We obtain the previous corollary applying the following general lemma to Theorem 5.5.

Lemma 5.9. *Let (X, Δ) be an orbifold defined over κ such that $(X, \text{supp}(\Delta))$ is pseudo algebraically hyperbolic with a constant $\alpha > 0$ and with exceptional set $Z \subset X$. Then, there exists a positive integer $m > 0$ such that, if all orbifold multiplicities are greater than m , (X, Δ) is pseudo algebraically hyperbolic with the same constant α and the same exceptional set Z .*

The Lemma and its proof generalizes [Rou10, Theorem 4.8] that was proved in the case where $X = \mathbb{P}^n$.

Proof. Let $\psi : (\mathcal{C}, \Delta_{\mathcal{C}}) \rightarrow (X, \Delta)$ be an orbifold morphism. We will use the notation $\delta := N_\psi^{[1]}(\text{supp}(\Delta))$. We first prove that there is a minimal orbifold structure $\tilde{\Delta}$ on \mathcal{C} which makes ψ into an orbifold morphism and for which it is sufficient to prove the result. Let

$$\begin{aligned} \psi^*(\Delta_j) &= \sum_{i=1}^{\delta} t_{i,j} P_i, \\ \psi^*(\text{supp}(\Delta)) &= \sum_{i=1}^{\delta} t_i P_i, \end{aligned}$$

where the P_i are the distinct points of $\psi^{-1}(\text{supp}(\Delta))$. Then if

$$\Delta_{\mathcal{C}} = \sum_{i=1}^{\delta} \left(1 - \frac{1}{m'_i}\right) P_i,$$

ψ is an orbifold morphism if $m'_i t_i \geq m_j$ for all $j \in \varphi(i)$ where $\varphi(i)$ is the set

$$\varphi(i) := \{k : 1 \leq k \leq \delta \text{ and } \psi(P_i) \in \Delta_k\}.$$

Therefore the minimal orbifold structure $\tilde{\Delta}$ on \mathcal{C} is given by the multiplicities

$$\tilde{m}_i = \sup_{j \in \varphi(i)} \left\lceil \frac{m_j}{t_i} \right\rceil,$$

where as usual $\lceil k \rceil$ denotes the round up of k . Therefore, we have to prove that there exists a constant α_Δ and an ample line bundle \mathcal{L}_Δ on X such that, for every ψ such that $\psi(\mathcal{C}) \not\subseteq Z$ we have

$$\deg \psi^* \mathcal{L}_\Delta \leq \alpha_\Delta (2g(\mathcal{C}) - 2 + \sum_{i=1}^{\delta} \left(1 - \frac{1}{\tilde{m}_i}\right)).$$

We can bound the contribution coming from the orbifold divisor as follows:

$$(5.4) \quad \sum_{i=1}^{\delta} \left(1 - \frac{1}{\tilde{m}_i}\right) \geq \sum_{i=1}^{\delta} \left(1 - \frac{t_i}{\sup_{j \in \varphi(i)} m_j}\right) \geq N_\psi^{[1]}(\text{supp}(\Delta)) - \sum_{i=1}^{\delta} \sum_{j=1}^q \frac{t_{i,j}}{m_j}$$

On the other hand, by definition

$$\sum_{i=1}^{\delta} t_{i,j} = \deg(\psi^* \Delta_j).$$

which implies that we can rewrite (5.4) as

$$(5.5) \quad \sum_{i=1}^{\delta} \left(1 - \frac{1}{\tilde{m}_i}\right) \geq N_\psi^{[1]}(\text{supp}(\Delta)) - \sum_{j=1}^q \frac{\deg(\psi^* \Delta_j)}{m_j},$$

In particular, equation (5.5) can be rewritten as

$$(5.6) \quad N_\psi^{[1]}(\text{supp}(\Delta)) \leq \sum_{i=1}^{\delta} \left(1 - \frac{1}{\tilde{m}_i}\right) + \sum_{j=1}^q \frac{\deg(\psi^* \Delta_j)}{m_j}.$$

On the other hand, our assumption that $(X, \text{supp}(\Delta))$ is pseudo algebraically hyperbolic, implies that, for every ample line bundle \mathcal{L} on L , if $\psi(\mathcal{C}) \not\subseteq Z$, we have

$$(5.7) \quad \deg \psi^* \mathcal{L} \leq \alpha \left(2g(\mathcal{C}) - 2 + N_\psi^{[1]}(\text{supp} \Delta)\right)$$

Finally, combining equation (5.7) with equation (5.6), we obtain

$$\deg \psi^* \mathcal{L} \leq \alpha \left(2g(\mathcal{C}) - 2 + \deg \tilde{\Delta}\right) + \deg \psi^* \left(\sum_{j=1}^q \frac{\alpha}{m_j} \Delta_j\right).$$

Therefore we conclude by noticing that, when the multiplicity m_j are big enough, the line bundle

$$\mathcal{L}_\Delta = \mathcal{L} \otimes \mathcal{O}_X \left(-\sum_{j=1}^q \frac{\alpha}{m_j} \Delta_j\right)$$

is ample on X and therefore we obtain

$$\deg \psi^* \mathcal{L}_\Delta \leq \alpha \left(2g(\mathcal{C}) - 2 + \deg \tilde{\Delta}\right).$$

□

5.2. Holomorphic Maps

In this subsection we are interested in degeneracy properties of holomorphic maps. All varieties will be defined over \mathbb{C} . As before we will obtain our result as a consequence of the following generalization of [RV, General Theorem (Analytic Part)] that includes truncation.

Theorem 5.10. *Let X be a complex projective variety of dimension n and let D_1, \dots, D_q be effective Cartier divisors intersecting properly on X . Let \mathcal{L} be a big line bundle. For each $\varepsilon > 0$, there exists a positive integer Q such that for any algebraically non-degenerate holomorphic map $f : \mathbb{C} \rightarrow X$, the inequality*

$$\sum_{j=1}^q \beta(\mathcal{L}, D_j) T_{D_j, f}(r) - (1 + \varepsilon) T_{\mathcal{L}, f}(r) \leq_{\text{exc}} \sum_{j=1}^q \beta(\mathcal{L}, D_j) N_f^{(Q)}(D_j, r)$$

holds, where \leq_{exc} means that the inequality holds for all $r \in \mathbb{R}^+$ except a set of finite Lebesgue measure.

The proof of Theorem 5.10 will be given in Section 7 (see in particular Theorem 7.1). Using the previous theorem we obtain a Nevanlinna analogue of Theorem 5.5 and Corollary 5.8.

Theorem 5.11. *Let $X \subset \mathbb{P}^m$ be a complex nonsingular projective surface and $D = D_1 + \dots + D_q$ be a divisor with $q \geq 2$, such that*

- (1) *No three of the components D_i meet at a point;*
- (2) *There exists a choice of positive integers p_i such that*
 - *the divisor $D_p := p_1 D_1 + p_2 D_2 + \dots + p_q D_q$ is ample;*
 - *The following inequality holds:*

$$2D_p^2 \xi_i > (D_p \cdot D_i) \xi_i^2 + 3D_p^2 p_i,$$

for every $i = 1, \dots, q$ where ξ_i is the minimal positive solution of the equation $D_i^2 x^2 - 2(D_p \cdot D_i)x + D_p^2 = 0$.

Let Δ be the \mathbb{Q} -divisor defined as

$$\Delta = \sum_{i=1}^q \left(1 - \frac{1}{m_i}\right) D_i.$$

Then, there exists a positive integer m such that, if $m_i \geq m$ for every i , every orbifold entire curve $\psi : \mathbb{C} \rightarrow (X, \Delta)$, is algebraically degenerate.

Proof. First we note that condition (1) implies that D_1, \dots, D_q are in general position. Let $f : \mathbb{C} \rightarrow (X, \Delta)$ be a Zariski dense orbifold entire curve. As before, we denote by $\beta_i = \beta(D_p, D_i)$ and we set

$$\varepsilon = \min\{(\beta_i - p_i)/p_i\}$$

which is positive by Lemma 5.6. By Theorem 5.10, applied with $\mathcal{L} = \mathcal{O}_X(D_p)$, and $\varepsilon/2$, there exists an integer Q such that the following inequality holds:

$$(5.8) \quad \sum_{j=1}^q \beta_j T_{D_j, f}(r) - \left(1 + \frac{\varepsilon}{2}\right) T_{D_p, f}(r) \leq_{\text{exc}} \sum_{j=1}^q \beta_j N_f^{(Q)}(D_j, r).$$

The analogue of Lemma 5.7 in the Nevanlinna setting gives the lower bound

$$\sum_{j=1}^q \beta_j T_{D_j, f}(r) > (1 + \varepsilon) T_{D_p, f}(r).$$

Together with (5.8), this implies that

$$(5.9) \quad \frac{\epsilon}{2} T_{D_p, f}(r) \leq \sum_{j=1}^q \beta_j N_f^{(Q)}(D_j, r).$$

Since f is an orbifold entire curve, one has

$$N_f^{(Q)}(D_j, r) \leq \frac{Q}{m_j} N_f(D_j, r) \leq \frac{Q}{m_j} T_{D_p, f}(r).$$

Therefore, we can rewrite equation (5.9), as

$$\frac{\epsilon}{2} T_{D_p, f}(r) \leq Q \left(\sum_{j=1}^q \frac{\beta_j}{m_j} \right) T_{D_p, f}(r) \leq \frac{Q}{m} \left(\sum_{j=1}^q \beta_j \right) T_{D_p, f}(r).$$

To conclude, it is enough to choose m big enough, such that

$$\frac{Q}{m} \left(\sum_{j=1}^q \beta_j \right) < \frac{\epsilon}{2}.$$

□

As a corollary we obtain the analogue of [CZ04, Main Theorem] and Theorem 5.5 in the Nevanlinna setting.

Corollary 5.12. *Let (X, D) be as in Theorem 5.11. Then, every entire curve $\psi : \mathbb{C} \rightarrow X \setminus D$, is algebraically degenerate. Moreover (X, D) is Brody hyperbolic modulo the exceptional set Z of Theorem 5.5, i.e. $\psi(\mathbb{C})$ is contained in Z .*

Proof. The first statement follows directly by Theorem 5.11. For the second statement one can apply directly [Lev09, Theorem 8.3 B] to our setting (since, as noted before, the proof of [CZ04, Main Theorem] shows that the divisor D_p is *large* in the sense of Levin). □

6. DEGENERACY PROPERTIES OF X_m

Given Lang and Vojta’s dictionary between arithmetic and geometry properties of a variety X , it is expected that varieties with a potentially dense set of rational points should correspond to manifolds admitting Zariski dense entire curves. Therefore the analogue of Conjecture 11 should imply that weakly special manifolds should admit such curves (see Conjecture 12).

Campana and Păun [CP07] have shown that some examples constructed in [BT04] (in particular with $\kappa(S) = 1$) give counterexamples to such a statement. In other words, there are some weakly special manifolds in which all entire curves are algebraically degenerate. The goal of this section is to show that one can produce much more “counterexamples” from the examples given in last section. In particular we show that the weakly special varieties X_m (for m big enough) provide counterexamples to Conjecture 12.

6.1. Construction

Following ideas of Corvaja and Zannier in [CZ10] we begin by giving a series of examples in which the results of Section 5 apply. As in Section 5 we denote by κ and algebraically closed field of characteristic 0. We start by the following definition.

Definition 6.1. Let D_1, D_2 and D_3 , be three projective plane curve in \mathbb{P}_κ^2 and let $D_4 = H$ be an hyperplane such that $D_i \cap D_j \cap D_k$ is empty for every distinct i, j, k . Given three distinct projective curves B_1, B_2, B_3 we say that $(D_1, B_1), (D_2, B_2), (D_3, B_3)$ and H are in *general position* if

- D_i and B_i intersect transversally for every $i = 1, 2, 3$;
- $D_i \cap D_j \cap B_h = \emptyset$, for every distinct i, j and $h \in \{i, j\}$.

The main source of examples will be certain blow-ups of \mathbb{P}^2 . In the next proposition we show that these satisfy the assumptions of Theorem 5.5 and Theorem 5.11.

Proposition 6.2. *Let $(D_1, B_1), (D_2, B_2), (D_3, B_3)$ and H be curves in general position as in Definition 6.1 such that $\deg D_i \geq \deg B_i$. Let T be the set of points $D_i \cap B_i$ for $i = 1, 2, 3$; if $\#(T \cap D_i) < \deg D_i^2$ we add to T smooth points of D_i so that $\#T = \deg D_1^2 + \deg D_2^2 + \deg D_3^2$.*

Let X be the blow up \mathbb{P}^2 along T , and denote by \tilde{D}_i the strict transform of D_i , and by \tilde{H} the strict transform of H . Let Δ be the \mathbb{Q} -divisor defined as

$$\Delta = \sum_{i=1}^3 \left(1 - \frac{1}{m_i}\right) \tilde{D}_i + \left(1 - \frac{1}{m_4}\right) \tilde{H}.$$

Then, there exists a positive integer $m > 0$ such that, if $m_i > m$, we have

- (1) (X, Δ) is pseudo algebraically hyperbolic;
- (2) if $\kappa = \mathbb{C}$, every orbifold entire curve $f : \mathbb{C} \rightarrow (X, \Delta)$ is degenerate.

Proof of Proposition 6.2. Let $d_i = \deg D_i$. By definition, we have the following intersection numbers in X : $\tilde{D}_i \cdot \tilde{D}_j = d_i d_j$, $\tilde{D}_i^2 = 0$ and $\tilde{D}_i \cdot \tilde{H} = d_i$. The result will follow from an application of Theorem 5.5. Let $c := 4d_1 d_2 d_3$ and define $p_i := c/d_i$ for $i = 1, 2, 3$, and $p_4 := 3c/4 = 3d_1 d_2 d_3$. Using the integers p_1, p_2, p_3 and p_4 we define

$$D_p = p_1 \tilde{D}_1 + p_2 \tilde{D}_2 + p_3 \tilde{D}_3 + p_4 \tilde{H}.$$

Then, it is immediate to verify that $D_p \cdot \tilde{H} > 0$ and $D_p \cdot E > 0$ where E is any exceptional divisor. Now let C be a non-exceptional curve in X . Then we can compute

$$(6.1) \quad D_p \cdot C = \left(\sum_{i=1}^3 p_i d_i + p_4 \right) H \cdot \pi_* C - \sum_{i=1}^3 p_i \sum_{Q \in D_i \cap T} \text{mult}_Q(\pi_* C).$$

Then, denoting $c = \deg \pi_* C$, Bezout's Theorem implies that $\sum_{Q \in D_i \cap T} \text{mult}_Q(\pi_* C) \leq c d_i$ and therefore we can rewrite equation (6.1) as

$$D_p \cdot C \geq c \left(\sum_{i=1}^3 p_i d_i + p_4 \right) - \sum_{i=1}^3 p_i (c d_i) = c p_4 > 0.$$

Finally, since $D_p^2 > 0$, by the Nakai-Moishezon criterion [Laz04, Theorem 1.2.23], the divisor D_p is ample.

Let $i = 1, 2, 3$, and let ξ_i be the smallest solution of

$$(6.2) \quad \tilde{D}_i^2 x^2 - 2(D_p \cdot \tilde{D}_i)x + D_p^2 = 0.$$

Using the fact that $\tilde{D}_i^2 = 0$ for $i = 1, 2, 3$ this gives the following expression

$$\xi_i = \frac{D_p^2}{2(D_p \cdot D_i)}.$$

Using this expression, in order to apply Theorem 5.5, we have to verify that the following inequality holds true:

$$(6.3) \quad 2D_p^2 \cdot \frac{D_p^2}{2(D_p \cdot \tilde{D}_i)} > (D_p \cdot \tilde{D}_i) \cdot \frac{D_p^4}{4(D_p \cdot \tilde{D}_i)^2} + 3D_p^2 \cdot p_i$$

This simplify to $D_p^2 > 4p_i(D_p \cdot \tilde{D}_i)$. Using the definition of D_p and p_i we can compute $D_p^2 = 177d^2$ where $d = d_1 d_2 d_3$, and $D_p \cdot \tilde{D}_i/d_i = 11d$. In particular $4p_i(D_p \cdot \tilde{D}_i) = 176d^2$ so that (6.3) reads $177d^2 > 176d^2$ and therefore it is verified in the case in which $i = 1, 2, 3$.

To conclude we need to verify that the same condition holds for $i = 4$. In this case, using that $D_p \cdot \tilde{H} = 15d$, equation (6.2) becomes

$$x^2 - 30dx + 177d^2 = 0$$

A direct computation shows that the smallest solution of the equation is $\xi = d(15 - 4\sqrt{3})$. Then, inequality (6.3) becomes

$$2 \cdot 177 \cdot (15 - 4\sqrt{3}) > 15 \cdot (15 - 4\sqrt{3})^2 + 9 \cdot 177$$

that is a true statement, thus concluding the verification of the hypotheses of Theorem 5.5 and Theorem 5.11 which imply the desired conclusion. \square

Remark 6.3. We note that the orbifold (X, Δ) will be of general type as soon as the m_i s are big enough. Moreover, in the case in which all the multiplicities $m_i = \infty$, we recover the analogue of [CZ10, Proposition 2] in the function field and Nevanlinna case.

6.2. Weakly-Special Threefolds

In order to prove degeneracy results for the threefolds X_m constructed in Section 4, we will use Proposition 6.2 in a special case, that guarantees that the blow-up is not of log general type (but of orbifold general type).

Let $D := D_1 + D_2 + D_3 + D_4$ be a simple normal crossing divisor on \mathbb{P}^2 where D_1 is a curve of degree d and D_2, D_3, D_4 are lines. Let $F := F_1 + F_2 + F_3$ be a simple normal crossing divisor on \mathbb{P}^2 where F_1 is a curve of degree d and F_2, F_3 are lines and such that $F + D$ is a simple normal crossing divisor (so that (D_i, F_i) for $i = 1, 2, 3$ and D_4 are in general position in the sense of Definition 6.1). Let $T_i := D_i \cap F_i$, $1 \leq i \leq 3$ and $T = T_1 \cup T_2 \cup T_3$. Let $\pi : \bar{S} \rightarrow \mathbb{P}^2$ be the blow-up of \mathbb{P}^2 in T and $\tilde{D}_i = \pi_*^{-1} D_i$ the strict transform of D_i for $i = 1, \dots, 4$. In order to construct the fibration g of Example 4.4 we consider the quasi-projective surface S given by $\bar{S} \setminus \tilde{D}_2 \cup \tilde{D}_3 \cup \tilde{D}_4$ and the fibration $g : S \rightarrow \mathbb{P}^1$ to be the fibration induced by D_1 and F_1 . Note that for any integer $m \geq 2$, the divisor $K_{\mathbb{P}^2} + (1 - 1/m)D_1 + D_2 + D_3 + D_4$ is big, which implies in particular that the orbifold

surface $(S, (1 - 1/m)\tilde{D}_1)$ is an orbifold of general type. Moreover $S \setminus \tilde{D}_1$ is simply connected (by the same argument as in Example 4.4).

Remark 6.4. The quasi projective surface $\bar{S} \setminus \tilde{D}_1 \cup \tilde{D}_2 \cup \tilde{D}_3 \cup \tilde{D}_4$ appears in [CZ10, Theorem 3] to produce an example of a simply-connected quasi-projective surface with non Zariski-dense set of integral points. The idea of our construction is to consider instead the *orbifold* surface $(\bar{S}, (1 - 1/m)\tilde{D}_1 + \tilde{D}_2 + \tilde{D}_3 + \tilde{D}_4)$, and prove the degeneracy of orbifold entire curves. Unfortunately, it seems that the arithmetic orbifold analogue is out of reach with the present methods.

Given f as in Example 4.3 we can construct the smooth threefold X_m as in Theorem 4.2: it is a weakly special threefold that is not special. We denote by $\pi : X_m \rightarrow S$ the elliptic fibration induced. Note that by construction the orbifold base (see Definition 2.4) of π coincides with $\Delta_\pi = (1 - 1/m)\tilde{D}_1$. Then, the following theorem proves degeneracy results analogues of Conjecture 10 and the analytic part of Conjecture 1 for the *quasi-projective* threefold X_m .

Theorem 6.5. *There exists m_0 such that for all $m \geq m_0$, the following holds:*

- (1) *there exists a positive constant A such that, for every morphism $\varphi : \mathcal{C} \rightarrow \bar{X}_m$ such that $\pi(\varphi(\mathcal{C}))$ is not contained in the exceptional set of Proposition 6.2, the following holds:*

$$\deg \pi(\varphi(\mathcal{C})) \leq A \left(2g(\mathcal{C}) - 2 + N_{\pi \circ \varphi}^{[1]}(\tilde{D}_2 + \tilde{D}_3 + \tilde{D}_4) \right).$$

- (2) *every entire curve $f : \mathbb{C} \rightarrow X_m$ is algebraically degenerate.*

In particular, X_m give counterexamples to (the logarithmic analogue of) Conjecture 12.

Proof. We first prove the statement in the function field case. Let φ be a morphism as above and consider $\pi \circ \varphi : \mathcal{C} \rightarrow \bar{S}$. By construction, this induces an orbifold morphism

$$\left(\mathcal{C}, \varphi^{-1}(\tilde{D}_2 + \tilde{D}_3 + \tilde{D}_4) \right) \rightarrow \left(\bar{S}, \left(1 - \frac{1}{m} \right) \tilde{D}_1 + \tilde{D}_2 + \tilde{D}_3 + \tilde{D}_4 \right).$$

Then the conclusion follows from Proposition 6.2. Similarly, in the Nevanlinna case, one considers an entire curve $f : \mathbb{C} \rightarrow X_m$: the composition $\pi \circ f$ is an orbifold entire curve to $(\bar{S}, (1 - 1/m)\tilde{D}_1 + \tilde{D}_2 + \tilde{D}_3 + \tilde{D}_4)$. Then the conclusion follows again from Proposition 6.2. \square

7. THE RU-VOJTA METHOD

The purpose of this section is to obtain a truncated version of Ru-Vojta's Theorem [RV, General Theorem (Analytic Part)], as well as its analogue in the function field setting.

7.1. Nevanlinna Theory

We first recall some definitions in Nevanlinna theory. Let D be an effective Cartier divisor on a complex variety X . Let $s = 1_D$ be a canonical section of $\mathcal{O}(D)$, i.e. a global section for which $(s) = D$, and fix a smooth metric $|\cdot|$ on $\mathcal{O}(D)$. The associated Weil function $\lambda_D : X(\mathbb{C}) \setminus \text{Supp } D \rightarrow \mathbb{R}$ is given by

$$\lambda_D(x) := -\log |s(x)|.$$

It is linear in D (over a suitable domain), so by linearity and continuity it can be defined for a general Cartier divisor D on X .

Let $f : \mathbb{C} \rightarrow X$ be a holomorphic map whose image is not contained in the support of divisor D on X . The *proximity function* of f with respect to D is defined by

$$m_f(D, r) = \int_0^{2\pi} \lambda_D(f(re^{i\theta})) \frac{d\theta}{2\pi}.$$

Let $n_f(D, t)$ (respectively, $n_f^{(Q)}(D, t)$) be the number of zeros of $\rho \circ f$ inside $\{|z| < t\}$, counting multiplicity, (respectively, ignoring multiplicity larger than $Q \in \mathbb{N}$) with ρ a local equation of D . The *counting function* and the *truncated counting function* of f of order Q at ∞ are defined, respectively, by

$$N_f(D, r) = \int_1^r \frac{n_f(D, t)}{t} dt \quad \text{and} \quad N_f^{(Q)}(D, r) = \int_1^r \frac{n_f^{(Q)}(D, t)}{t} dt.$$

The height function relative to D is defined, up to $O(1)$, as

$$(7.1) \quad T_{D,f}(r) = m_f(D, r) + N_f(D, r).$$

The following is a truncated version of the analytic part [RV, General Theorem].

Theorem 7.1. *Let X be a complex projective variety of dimension n , let D_1, \dots, D_q be effective Cartier divisors intersecting properly on X , and let \mathcal{L} be a big line sheaf. For each $\varepsilon > 0$, there exists a positive integer Q such that for any algebraically non-degenerate holomorphic map $f : \mathbb{C} \rightarrow X$, the inequality*

$$\sum_{j=1}^q \beta(\mathcal{L}, D_j) T_{D_j, f}(r) - (1 + \varepsilon) T_{\mathcal{L}, f}(r) \leq_{\text{exc}} \sum_{j=1}^q \beta(\mathcal{L}, D_j) N_f^{(Q)}(D_j, r)$$

holds, where \leq_{exc} means the inequality holds for all $r \in \mathbb{R}^+$ except a set of finite Lebesgue measure.

Remark 7.2. When X is a surface, the condition that D_1, \dots, D_q intersect properly on X can be relaxed to a general position assumptions, following the same strategy as in [HR18, Main Theorem].

We recall the following theorem from [GW19], which is a modification of [HR18, Theorem 3.2] by applying the general form of the second main theorem with a Wronskian term proved by Vojta in [Voj97, Theorem 1] and Ru in [Ru97, Theorem 2.3].

Theorem 7.3. *Let X be a complex projective variety and let D be a Cartier divisor on X , let V be a nonzero linear subspace of $H^0(X, \mathcal{O}(D))$, let s_1, \dots, s_q be nonzero elements of V and for each $j = 1, \dots, q$, let D_j be the Cartier divisor (s_j) .*

Let $\Phi = (\phi_1, \dots, \phi_d) : X \dashrightarrow \mathbb{P}^{d-1}$ be the rational map associated to the linear system V , let $f : \mathbb{C} \rightarrow X$ be a holomorphic map with Zariski dense image and let $\Psi = (\psi_1, \dots, \psi_d) : \mathbb{C} \rightarrow \mathbb{P}^{d-1}$ be a reduced form of $\Phi \circ f$, i.e. $\Psi = \Phi \circ f$ and ψ_1, \dots, ψ_d are entire functions without common zeros. Denote by $W(\Psi)$ the Wronskian of ψ_1, \dots, ψ_d . Then for any $\varepsilon > 0$,

$$\int_0^{2\pi} \max_J \sum_{j \in J} \lambda_{D_j} \left(f(re^{i\theta}) \right) \frac{d\theta}{2\pi} + N_{W(\Psi)}(0, r) + \dim V \cdot N_h(0, r) \leq_{\text{exc}} (\dim V + \varepsilon) T_{D,f}(r),$$

where J ranges over all subsets of $\{1, \dots, q\}$ such that the sections $(s_j)_{j \in J}$ are linearly independent, and h is an entire function such that $h\psi_i = \phi_i(f)$ for each $1 \leq i \leq d$, i.e. h is a gcd of $\phi_i(f)$, $1 \leq i \leq d$.

Proof. Most of the proof is identical to the proof of [HR18, Theorem 3.2], therefore we will omit some details and indicate the required adjustments.

We may assume that $d > 1$. Let X' be the closure of the graph of Φ , and let $p: X' \rightarrow X$ and $\phi: X' \rightarrow \mathbb{P}^{d-1}$ be the projection morphisms. Then there is an effective Cartier divisor B on X' such that, for each nonzero $s \in V$, there is a hyperplane H in \mathbb{P}^{d-1} such that $p^*(s) - B = \phi^*H$. Let $\tilde{f}: \mathbb{C} \rightarrow X'$ be the lifting of f . Then $\phi \circ \tilde{f} = (\psi_1, \dots, \psi_d): \mathbb{C} \rightarrow \mathbb{P}^{d-1}$ is a reduced presentation of $\Phi \circ f$. Moreover, let h be an entire function such that $h\psi_i = \phi_i(f)$ for each $1 \leq i \leq d$, i.e. h is a gcd of $\phi_i(f)$, $1 \leq i \leq d$. Then

$$(7.2) \quad N_{\tilde{f}}(B, r) \geq N_h(0, r) + O(1).$$

For each $j = 1, \dots, q$ let H_j be the hyperplane in \mathbb{P}^{d-1} for which

$$(7.3) \quad p^*D_j - B = p^*(s_j) - B = \phi^*H_j.$$

Choose a Weil function λ_B for B . Then, we have

$$\lambda_{p^*D_j} = \lambda_{\phi^*H_j} + \lambda_B + O(1).$$

By the functoriality of Weil functions,

$$\lambda_{p^*D_j}(\tilde{f}(z)) = \lambda_{D_j}(f(z)) \quad \text{and} \quad \lambda_{\phi^*H_j}(\tilde{f}(re^{i\theta})) = \lambda_{H_j}(\phi(\tilde{f}(re^{i\theta}))).$$

By the general form of the second main theorem with a Wronskian term in [Voj97, Theorem 1] and [Ru97, Theorem 2.3], we have

$$\int_0^{2\pi} \max_J \sum_{j \in J} \lambda_{H_j}(\phi(\tilde{f})(re^{i\theta})) \frac{d\theta}{2\pi} + N_{W(\phi(\tilde{f}))}(0, r) \leq_{\text{exc}} (d + \varepsilon) T_{\phi(\tilde{f})}(r).$$

From (7.3), we have

$$T_{\phi(\tilde{f})}(r) = T_{D, f} - T_{B, \tilde{f}}.$$

Since each set J has most $\dim V$ elements and B is effective, it follows from (7.2) that

$$(\#J) \left(m_{\tilde{f}}(B, r) + N_h(0, r) \right) \leq (\#J) \left(m_{\tilde{f}}(B, r) + N_{\tilde{f}}(B, r) \right) \leq d T_{B, \tilde{f}} + O(1).$$

Hence,

$$\begin{aligned} & \int_0^{2\pi} \left(\max_J \sum_{j \in J} \lambda_{H_j}(\phi(\tilde{f})(re^{i\theta})) + \lambda_B(\tilde{f}(re^{i\theta})) \right) \frac{d\theta}{2\pi} + d N_h(0, r) + N_{W(\phi(\tilde{f}))}(0, r) \\ & \leq_{\text{exc}} (d + \varepsilon) T_{\phi(\tilde{f})}(r) + d T_{B, \tilde{f}} + O(1) \\ & \leq_{\text{exc}} (d + \varepsilon) T_{D, f} + O(1). \end{aligned}$$

□

Proof of Theorem 7.1. We will follow the proof of the analytic part of [RV, General Theorem] closely, and only indicate the necessary modification. Let $\varepsilon > 0$ be given. We want to show that

$$(7.4) \quad \sum_{j=1}^q \beta(\mathcal{L}, D_j) \left(T_{D_j, f}(r) - N_f^{(Q)}(D_j, r) \right) \leq_{\text{exc}} (1 + \varepsilon) T_{\mathcal{L}, f}(r).$$

Since the quantities $(T_{D_j, f}(r) - N_f^{(Q)}(D_j, r))/T_{\mathcal{L}, f}(r)$ are bounded when their respective denominators are sufficiently large, it suffices to prove (7.4) with a slightly smaller $\varepsilon > 0$ and with $\beta(\mathcal{L}, D_i)$

replaced by slightly smaller $\beta_i \in \mathbb{Q}$ for all i . It's also clear that we may assume that $\beta_i \neq 0$ for each i .

Choose positive integers N and b such that

$$(7.5) \quad \left(1 + \frac{n}{b}\right) \max_{1 \leq i \leq q} \frac{\beta_i N h^0(X, \mathcal{L}^N)}{\sum_{m \geq 1} h^0(X, \mathcal{L}^N(-mD_i))} < 1 + \epsilon.$$

Let

$$\Sigma = \left\{ \sigma \subseteq \{1, \dots, q\} \mid \bigcap_{j \in \sigma} \text{Supp } D_j \neq \emptyset \right\}.$$

For $\sigma \in \Sigma$, let

$$\Delta_\sigma = \left\{ \mathbf{a} = (a_i) \in \prod_{i \in \sigma} \beta_i^{-1} \mathbb{N} \mid \sum_{i \in \sigma} \beta_i a_i = b \right\}.$$

For $\mathbf{a} \in \Delta_\sigma$ as above, we construct a filtration of $H^0(X, \mathcal{L}^N)$ as follows: For $x \in \mathbb{R}^+$, one defines the ideal $\mathcal{I}_\mathbf{a}(x)$ of \mathcal{O}_X by

$$\mathcal{I}_\mathbf{a}(x) = \sum_{\mathbf{b}} \mathcal{O}_X \left(- \sum_{i \in \sigma} b_i D_i \right)$$

where the sum is taken for all $\mathbf{b} \in \mathbb{N}^{\#\sigma}$ with $\sum_{i \in \sigma} a_i b_i \geq bx$. Let

$$\mathcal{F}(\sigma; \mathbf{a})_x = H^0(X, \mathcal{L}^N \otimes \mathcal{I}_\mathbf{a}(x)),$$

which we regard as a subspace of $H^0(X, \mathcal{L}^N)$. We note that there are only finitely many ordered pairs (σ, \mathbf{a}) with $\sigma \in \Sigma$ and $\mathbf{a} \in \Delta_\sigma$. Let $\mathcal{B}_{\sigma; \mathbf{a}}$ be a basis of $H^0(X, \mathcal{L}^N)$ adapted to the above filtration $\{\mathcal{F}(\sigma; \mathbf{a})_x\}_{x \in \mathbb{R}^+}$.

For a basis \mathcal{B} of $H^0(X, \mathcal{L}^N)$, denote by (\mathcal{B}) the sum of the divisors (s) for all $s \in \mathcal{B}$. We now state the following main lemma in [RV].

Lemma 7.4. [RV, Lemma 6.8] *With the above notation, we have*

$$(7.6) \quad \bigvee_{\substack{\sigma \in \Sigma \\ \mathbf{a} \in \Delta_\sigma}} (\mathcal{B}_{\sigma; \mathbf{a}}) \geq \frac{b}{b+n} \left(\min_{1 \leq i \leq q} \frac{\sum_{m=1}^{\infty} h^0(X, \mathcal{L}^N(-mD_i))}{\beta_i} \right) \sum_{i=1}^q \beta_i D_i.$$

Here, the notation “ \bigvee ” is referred to the least upper bound with respect the partial order on the (Cartier) divisor group of X by the relation $D_1 \leq D_2$ if $D_2 - D_1$ is effective. Write

$$\bigcup_{\sigma; \mathbf{a}} \mathcal{B}_{\sigma; \mathbf{a}} = \mathcal{B}_1 \cup \dots \cup \mathcal{B}_{T_1} = \{s_1, \dots, s_{T_2}\}.$$

For each $i = 1, \dots, T_1$, let $J_i \subseteq \{1, \dots, T_2\}$ be the subset such that $\mathcal{B}_i = \{s_j : j \in J_i\}$. Choose Weil functions $\lambda_{\mathcal{B}_i}$ ($i = 1, \dots, T_1$), and λ_{s_j} ($j = 1, \dots, T_2$) for the divisors (\mathcal{B}_i) , and (s_j) , respectively. Then, by (7.6) for $x \in X$,

$$(7.7) \quad \begin{aligned} & \frac{b}{b+n} \left(\min_{1 \leq i \leq q} \sum_{m \geq 1} \frac{h^0(\mathcal{L}^N(-mD_i))}{\beta_i} \right) \sum_{i=1}^q \beta_i \lambda_{D_i}(x) \\ & \leq \max_{1 \leq i \leq T_1} \lambda_{\mathcal{B}_i}(x) + O(1) = \max_{1 \leq i \leq T_1} \sum_{j \in J_i} \lambda_{s_j}(x) + O(1). \end{aligned}$$

Let $M = h^0(X, \mathcal{L}^N)$ and let the set $\{\phi_1, \dots, \phi_M\}$ be a basis of the vector space $H^0(X, \mathcal{L}^N)$, and let $\Phi = [\phi_1, \dots, \phi_M] : X \rightarrow \mathbb{P}^{M-1}(\mathbb{C})$ be the corresponding (rational) map. Let $f : \mathbb{C} \rightarrow X$ be an algebraically non-degenerate holomorphic map. Let h be a gcd of $\phi_1(f), \dots, \phi_M(f)$, i.e. h is an entire function such that $\psi_1 := h^{-1}\phi_1(f), \dots, \psi_M := h^{-1}\phi_M(f)$ have no common zeros. Then $\Psi = (\psi_1, \dots, \psi_M)$ is a reduced form of $\Phi \circ f$.

By Theorem 7.3,

$$(7.8) \quad \int_0^{2\pi} \max_J \sum_{j \in J} \lambda_{s_j}(f(re^{i\theta})) \frac{d\theta}{2\pi} \leq_{\text{exc}} (M + \epsilon) T_{f, \mathcal{L}^N}(r) - N_{W(\Psi)}(0, r) - MN_h(0, r),$$

here the maximum is taken over all subsets J of $\{1, \dots, T_2\}$ for which the sections s_j , $j \in J$, are linearly independent. Combining (7.7) and (7.8), we have

$$\begin{aligned} & \sum_{i=1}^q \beta_i m_f(D_i, r) \\ & \leq_{\text{exc}} \left(1 + \frac{n}{b}\right) \max_{1 \leq i \leq q} \frac{\beta_i}{\sum_{m \geq 1} h^0(\mathcal{L}^N(-mD_i))} ((M + \epsilon) T_{f, \mathcal{L}^N}(x) - N_{W(\Psi)}(0, r) - MN_h(0, r)). \end{aligned}$$

Using (7.5) and the fact that $T_{f, \mathcal{L}^N}(r) = NT_{f, \mathcal{L}}(r)$, we have

$$\sum_{i=1}^q \beta_i m_f(D_i, r) \leq (1 + 2\epsilon) T_{f, \mathcal{L}}(x) - C(N_{W(\Psi)}(0, r) + MN_h(0, r)),$$

where $C = (1 + \epsilon)(MN)^{-1}$. Applying (7.1) to the above inequality, we have

$$\sum_{j=1}^q \beta_j T_{D_j, f}(r) - (1 + 2\epsilon) T_{\mathcal{L}, f}(r) \leq_{\text{exc}} \sum_{j=1}^q \beta_j N_f(D_j, r) - C(N_{W(\Psi)}(0, r) + MN_h(0, r)).$$

To finish the proof, we will show that there exists a large integer Q (to be determined later) such that

$$(7.9) \quad \sum_{j=1}^q \beta_j N_f(D_j, r) - C(N_{W(\Psi)}(0, r) + MN_h(0, r)) \leq \sum_{j=1}^q \beta_j N_f^{(Q)}(D_j, r).$$

For $z_0 \in \mathbb{C}$, let ρ_j be a local defining function of D_j around one of its open neighborhoods U . To show (7.9), it suffices to deduce the following inequality

$$\sum_{j=1}^q \beta_j v_{z_0}^+(\rho_j \circ f) - C \cdot (v_{z_0}^+(W(\Psi)) + Mv_{z_0}(h)) \leq \sum_{j=1}^q \beta_j \min\{Q, v_{z_0}^+(\rho_j \circ f)\},$$

for each $z_0 \in \mathbb{C}$.

The above inequality holds trivially if $v_{z_0}^+(\rho_j \circ f) \leq Q$ for each $1 \leq j \leq q$. Therefore, we assume that $v_{z_0}^+(\rho_j \circ f) \geq Q$ for some $1 \leq j \leq q$. By Lemma 7.4, there exists a set of basis s_1, \dots, s_M of $H^0(X, \mathcal{L}^N)$ with the following property:

$$(7.10) \quad \begin{aligned} \sum_{i=1}^M (s_i) & \geq \frac{b}{b+n} \left(\min_{1 \leq i \leq q} \sum_{m=1}^{\infty} \frac{h^0(X, \mathcal{L}^N(-mD_i))}{\beta_i} \right) \sum_{i=1}^q \beta_i D_i \\ & \geq \frac{MN}{1+\epsilon} \sum_{i=1}^q \beta_i D_i = \frac{1}{C} \sum_{i=1}^q \beta_i D_i, \end{aligned}$$

where the last inequality is due to (7.5). Since

$$v_{z_0}^+(s_i|_U \circ f) \geq \sum_{j=1}^q \text{ord}_{D_j} s_i \cdot v_{z_0}^+(\rho_j \circ f),$$

we can derive from (7.10) that

$$(7.11) \quad \sum_{i=1}^M v_{z_0}^+(s_i|_U \circ f) \geq \frac{1}{C} \sum_{j=1}^q \beta_j v_{z_0}^+(\rho_j \circ f).$$

On the other hand, since $\{\phi_1, \dots, \phi_M\}$ is a basis of the vector space $H^0(X, \mathcal{L}^N)$, each s_j is a \mathbb{C} -linear combination of the ϕ_i 's, and hence each $h^{-1}s_j(f)$ is a linear combination of the ψ_i 's. From the basic properties of Wronskians, we have

$$(7.12) \quad v_{z_0}^+(W(\Psi)) \geq \sum_{j=1}^M v_{z_0}^+(s_j|_U \circ f) - Mv_{z_0}(h) - \frac{1}{2}M(M-1).$$

Combining (7.11) and (7.12) and the assumption that $v_{z_0}^+(\rho_j \circ f) \geq Q$ for some $1 \leq j \leq q$, we obtain

$$\begin{aligned} \sum_{j=1}^q \beta_j v_{z_0}^+(\rho_j \circ f) - C \cdot (v_{z_0}^+(W(\Phi \circ f)) + Mv_{z_0}(h)) &\leq \frac{1}{2}CM(M-1) \\ &\leq \sum_{j=1}^q \beta_j \min\{Q, v_{z_0}^+(\rho_j \circ f)\}, \end{aligned}$$

where Q is chosen to be $\frac{CM(M-1)}{2 \min_{1 \leq j \leq q} \{\beta_j\}}$. (Note that we have assumed that $\beta_j \neq 0$ for each j .) This completes our proof. \square

7.2. Function Fields

In this section we let κ be an algebraically closed field of characteristic zero, let \mathcal{C} be a smooth projective curve over κ of genus $g(\mathcal{C})$ and let $K = \kappa(\mathcal{C})$ be the function field of \mathcal{C} . For each point $\mathfrak{p} \in \mathcal{C}$, we may choose a uniformizer $t_{\mathfrak{p}}$ to define a normalized order function $v_{\mathfrak{p}} := \text{ord}_{\mathfrak{p}} : K \rightarrow \mathbb{Z} \cup \{\infty\}$ at \mathfrak{p} . For a non-zero element $f \in K$, the *height* $h(f)$ counts its number of poles with multiplicities, i.e.

$$h(f) := \sum_{\mathfrak{p} \in \mathcal{C}} \max\{0, -v_{\mathfrak{p}}(f)\}.$$

Let $f_0, \dots, f_m \in K$ not all zeros. Then $\mathbf{f} = [f_0 : \dots : f_m] \in \mathbb{P}^m(K)$ can be viewed as a morphism from \mathcal{C} to $\mathbb{P}^m(\kappa)$. The height of this morphism (or of the corresponding point in $\mathbb{P}^m(K)$) is defined by

$$h(\mathbf{f}) = h(f_0, \dots, f_m) := \sum_{\mathfrak{p} \in \mathcal{C}} -\min\{v_{\mathfrak{p}}(f_0), \dots, v_{\mathfrak{p}}(f_m)\}.$$

Let D be a Cartier divisor on a variety X over κ . Similarly to the number field case, the classical theory of heights (see e.g. [HS00, Part B] or [BG06, Chapter 2]) associates to every Cartier

divisor D on X a height function $h_D : X(K) \rightarrow \mathbb{R}$ and a Weil function (local height function) $\lambda_{D,\mathfrak{p}} : X(K) \setminus \text{Supp } D \rightarrow \mathbb{R}$, $\mathfrak{p} \in \mathcal{C}$, well-defined up to a bounded functions such that

$$(7.13) \quad \sum_{\mathfrak{p} \in \mathcal{C}} \lambda_{D,\mathfrak{p}}(P) = h_D(P) + O(1)$$

for all $P \in X(K) \setminus \text{Supp } D$. Let S be a finite set of points in \mathcal{C} . We denote by

$$m_{D,S}(P) = \sum_{\mathfrak{p} \in S} \lambda_{D,\mathfrak{p}}(P), \quad \text{and} \quad N_{D,S}(P) = \sum_{\mathfrak{p} \notin S} \lambda_{D,\mathfrak{p}}(P)$$

the proximity and counting functions as defined in [Voj87]. Let $a_0X_0 + a_1X_1 + \cdots + a_mX_m$ be a linear form with $a_0, \dots, a_m \in \kappa$ whose vanishing determine a hyperplane H in \mathbb{P}^m . Then for all $\mathfrak{p} \in \mathcal{C}$ and $P = [x_0 : \cdots : x_m] \in \mathbb{P}^m(K)$, the Weil function at \mathfrak{p} is given by

$$\lambda_{H,\mathfrak{p}}(P) = v_{\mathfrak{p}}(a_0x_0 + a_1x_1 + \cdots + a_mx_m) - \min\{v_{\mathfrak{p}}(x_0), \dots, v_{\mathfrak{p}}(x_m)\}.$$

We recall the following version of the second main theorem for function fields from [Wan04].

Theorem 7.5 ([Wan04, Theorem 1]). *In the above setting, let H_1, \dots, H_q be hyperplanes in $\mathbb{P}^m(K)$ defined by linear forms with coefficients in κ . If $P = [x_0 : \cdots : x_m] \in \mathbb{P}^m(K)$ is linearly nondegenerate over κ , then*

$$\sum_{\mathfrak{p} \in S} \max_{j \in J} \lambda_{H_j,\mathfrak{p}}(P) \leq (m+1)h(P) + \frac{m(m+1)}{2}(2g(\mathcal{C}) - 2 + \#S),$$

where the maximum is taken over all subset J of $\{1, \dots, q\}$ such that the linear forms defining H_j , $j \in J$, are linearly independent.

We will use Theorem 7.5 to obtain a function field analogue of Theorem 7.1 with an explicit dependence on the Euler characteristic of the curve \mathcal{C} .

Theorem 7.6. *Let $X \subset \mathbb{P}^m$ be a projective variety over κ of dimension n , let D_1, \dots, D_q be effective Cartier divisors intersecting properly on X , and let \mathcal{L} be a big line sheaf. Then for any $\epsilon > 0$, there exist constants c_1 and c_2 , independent of the curve \mathcal{C} and the set S , such that for any map $x = [x_0 : \dots : x_m] : \mathcal{C} \rightarrow X$, where $x_i \in K$, outside the augmented base locus of \mathcal{L} we have either*

$$\sum_{i=1}^q \beta_{\mathcal{L},D_i} m_{D_i,S}(x) \leq (1 + \epsilon)h_{\mathcal{L}}(x) + c_1 \max\{1, 2g(\mathcal{C}) - 2 + |S|\},$$

or the image of x is contained in a hypersurface (over κ) in \mathbb{P}^m of degree at most c_2 .

Proof. The proof is similar to the first part of the proof of Theorem 7.1. We will follow its argument and notation and only indicate the modification. Let $\epsilon > 0$ be given. Since \mathcal{L} is a big line sheaf, there is a constant c such that $\sum_{i=1}^q h_{D_i}(x) \leq ch_{\mathcal{L}}(x)$ for all $x \in X(K)$ outside the augmented base locus B of \mathcal{L} . This follows from [Voj11, Proposition 10.11]. By (7.13), together with the fact that $m_{D_i,S} \leq h_{D_i} + O(1)$, we can choose $\beta_i \in \mathbb{Q}$ for all i such that

$$\sum_{i=1}^q (\beta_{\mathcal{L},D_i} - \beta_i) m_{D_i,S}(x) \leq \frac{\epsilon}{2} h_{\mathcal{L}}(x)$$

for all $x \in X \setminus B(K)$. Therefore, we can assume that $\beta_{\mathcal{L}, D_i} = \beta_i \in \mathbb{Q}$ for all i . It's also clear that we may assume that $\beta_i \neq 0$ for each i . From now on we will assume that the point $x \in X(K)$ does *not* lie on B .

Choose positive integers N and b to satisfy (7.5). The same arguments in the proof of Theorem 7.1 give

$$(7.14) \quad \begin{aligned} & \frac{b}{b+n} \left(\min_{1 \leq i \leq q} \sum_{m \geq 1} \frac{h^0(\mathcal{L}^N(-mD_i))}{\beta_i} \right) \sum_{i=1}^q \beta_i \lambda_{D_i, \mathfrak{p}}(x) \\ & \leq \max_{1 \leq i \leq T_1} \lambda_{\mathcal{B}_i, \mathfrak{p}}(x) + O(1) = \max_{1 \leq i \leq T_1} \sum_{j \in J_i} \lambda_{s_j, \mathfrak{p}}(x) + O(1). \end{aligned}$$

Let $M = h^0(X, \mathcal{L}^N)$, let the set $\{\phi_1, \dots, \phi_M\}$ be a basis of the vector space $H^0(X, \mathcal{L}^N)$, and let

$$(7.15) \quad \Phi = [\phi_1, \dots, \phi_M] : X \dashrightarrow \mathbb{P}^{M-1}(\kappa)$$

be the corresponding rational map. By Theorem 7.5, either the map $\Phi \circ x$ is linearly degenerate, i.e. $\phi_1(x), \dots, \phi_M(x)$ are linearly dependent over κ , or

$$(7.16) \quad \sum_{\mathfrak{p} \in S} \max_J \sum_{j \in J} \lambda_{s_j, \mathfrak{p}}(x) \leq M h_{\mathcal{L}^N}(x) + \frac{M(M-1)}{2} (2g - 2 + |S|),$$

here the maximum is taken over all subsets J of $\{1, \dots, T_2\}$ for which the sections s_j , $j \in J$, are linearly independent. We first consider when ϕ_1, \dots, ϕ_M are linearly independent over κ . Combining (7.14) and (7.16) gives

$$\sum_{i=1}^q \beta_i m_{D_i, S}(x) \leq \left(1 + \frac{n}{b}\right) \max_{1 \leq i \leq q} \frac{\beta_i}{\sum_{m \geq 1} h^0(\mathcal{L}^N(-mD_i))} M h_{\mathcal{L}^N}(x) + c'_1 (2g - 2 + |S|) + O(1),$$

where $c'_1 = \frac{M(M-1)}{2}$. Using (7.5) and the fact that $h_{\mathcal{L}^N}(x) = N h_{\mathcal{L}}(x)$, we have

$$\sum_{i=1}^q \beta_i m_{D_i, S}(x) \leq (1 + \epsilon) h_{\mathcal{L}}(x) + c'_1 (2g - 2 + |S|) + O(1),$$

which implies the first case of the Theorem.

To conclude we note that, if $\phi_1(x), \dots, \phi_M(x)$ are linearly dependent over κ , there exist constants $a_1, \dots, a_M \in \kappa$, not all zero, such that $a_1 \phi_1(x) + \dots + a_M \phi_M(x) = 0$. Let H be the hyperplane in \mathbb{P}^{M-1} defined by $a_1 z_1 + \dots + a_M z_M = 0$; by assumption $\Phi(x(\mathcal{C}))$ is contained in H . On the other hand, since ϕ_1, \dots, ϕ_M is a basis of $H^0(X, \mathcal{L}^N)$, it follows that $\Phi(X)$ is not contained in H , hence $\Phi(x(\mathcal{C}))$ is contained in $\Phi(X) \cap H$ which is an hypersurface in $\Phi(X)$ whose degree is bounded independently of \mathcal{C} and x as wanted. \square

Corollary 7.7. *In the previous setting, if X has dimension 2, then for any $\epsilon > 0$, there exist constants c_1 and c_3 , independent of the curve \mathcal{C} and the set S , such that for any K -point $x = [x_0 : \dots : x_m] : \mathcal{C} \rightarrow X$, with $x_i \in K$, we have that, either $\deg x(\mathcal{C}) \leq c_3$, or*

$$\sum_{i=1}^q \beta(\mathcal{L}, D_i) m_{D_i, S}(x) \leq (1 + \epsilon) h_{\mathcal{L}}(x) + c_1 \max\{1, 2g(\mathcal{C}) - 2 + |S|\}.$$

Proof. Given Theorem 7.6, it is enough to observe that, since X has dimension 2, the hypersurface of degree bounded by c_2 intersects X in a union of finitely many curves whose degree is bounded independently of the point x . Moreover, the augmented base locus of \mathcal{L} has dimension at most 1, and its one dimensional locus is a union of finitely many curves, which are also independent of x . Therefore, it suffices to define c_3 to be the maximum of the degrees of all these finitely many curves. \square

REFERENCES

- [ADT18] Kenneth Ascher, Kristin DeVleming, and Amos Turchet. Hyperbolicity and uniformity of log general type varieties. *arXiv preprint arXiv:1807.05946*, 2018.
- [BG06] Enrico Bombieri and Walter Gubler. *Heights in Diophantine geometry*, volume 4 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2006.
- [Bog79] Fedor A. Bogomolov. Holomorphic tensors and vector bundles on projective varieties. *Math. USSR, Izv.*, 13:499–555, 1979.
- [BRT19] Federico Lo Bianco, Erwan Rousseau, and Frédéric Touzet. Symmetries of transversely projective foliations. *arXiv preprint arXiv:1901.05656*, 2019.
- [BT04] Fedor Bogomolov and Yuri Tschinkel. Special elliptic fibrations. In *The Fano Conference*, pages 223–234. Univ. Torino, Turin, 2004.
- [Cam04] Frédéric Campana. Orbifolds, special varieties and classification theory. *Ann. Inst. Fourier (Grenoble)*, 54(3):499–630, 2004.
- [Cam11] Frédéric Campana. Orbifolies géométriques spéciales et classification biméromorphe des variétés kählériennes compactes. *J. Inst. Math. Jussieu*, 10(4):809–934, 2011.
- [Che04] Xi Chen. On algebraic hyperbolicity of log varieties. *Commun. Contemp. Math.*, 6(4):513–559, 2004.
- [CP07] Frédéric Campana and Mihai Păun. Variétés faiblement spéciales à courbes entières dégénérées. *Compos. Math.*, 143(1):95–111, 2007.
- [CT19] Laura Capuano and Amos Turchet. Lang–Vojta conjecture over function fields for surfaces dominating \mathbb{G}_m^2 . *arXiv preprint arXiv:1911.07562*, 2019.
- [CZ04] Pietro Corvaja and Umberto Zannier. On integral points on surfaces. *Ann. of Math. (2)*, 160(2):705–726, 2004.
- [CZ08] Pietro Corvaja and Umberto Zannier. Some cases of Vojta’s Conjecture on integral points over function fields. *J. Algebraic Geometry*, 17:195–333, 2008.
- [CZ10] Pietro Corvaja and Umberto Zannier. Integral points, divisibility between values of polynomials and entire curves on surfaces. *Adv. Math.*, 225(2):1095–1118, 2010.
- [CZ13] Pietro Corvaja and Umberto Zannier. Algebraic hyperbolicity of ramified covers of \mathbb{G}_m^2 (and integral points on affine subsets of \mathbb{P}^2). *J. Differential Geom.*, 93(3):355–377, 2013.
- [Dem97] Jean-Pierre Demailly. Algebraic criteria for Kobayashi hyperbolic projective varieties and jet differentials. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 285–360. Amer. Math. Soc., Providence, RI, 1997.
- [FM94] Robert Friedman and John W. Morgan. *Smooth four-manifolds and complex surfaces*, volume 27 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1994.
- [GW19] Ji Guo and Julie Tzu-Yueh Wang. Asymptotic gcd and divisible sequences for entire functions. *Trans. Amer. Math. Soc.*, 371(9):6241–6256, 2019.
- [HR18] Saud Hussein and Min Ru. A general defect relation and height inequality for divisors in subgeneral position. *Asian J. Math.*, 22(3):477–491, 2018.
- [HS00] Marc Hindry and Joseph H. Silverman. *Diophantine geometry*, volume 201 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. An introduction.
- [HT00] Joe Harris and Yuri Tschinkel. Rational points on quartics. *Duke Math. J.*, 104(3):477–500, 2000.

- [JX19] Ariyan Javanpeykar and Junyi Xie. Finiteness properties of pseudo-hyperbolic varieties. *arXiv preprint arXiv:1909.12187*, 2019.
- [Lan86] Serge Lang. Hyperbolic and Diophantine analysis. *Bull. Amer. Math. Soc. (N.S.)*, 14(2):159–205, 1986.
- [Lan91] Serge Lang. *Number theory. III - Diophantine geometry*, volume 60 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1991.
- [Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 2004.
- [Lev09] Aaron Levin. Generalizations of Siegel’s and Picard’s theorems. *Ann. of Math. (2)*, 170(2):609–655, 2009.
- [McQ98] Michael McQuillan. Diophantine approximations and foliations. *Inst. Hautes Études Sci. Publ. Math.*, No. 87:121–174, 1998.
- [Nog82] Junjiro Noguchi. A higher-dimensional analogue of Mordell’s conjecture over function fields. *Math. Ann.*, 258(2):207–212, 1981/82.
- [Rou10] Erwan Rousseau. Hyperbolicity of geometric orbifolds. *Trans. Amer. Math. Soc.*, 362(7):3799–3826, 2010.
- [Ru97] Min Ru. On a general form of the second main theorem. *Trans. Amer. Math. Soc.*, 349(12):5093–5105, 1997.
- [RV] Min Ru and Paul Vojta. A birational nevanlinna constant and its consequences. *Amer. J. Math.*, to appear.
- [Tur17] Amos Turchet. Fibered threefolds and Lang-Vojta’s conjecture over function fields. *Trans. Amer. Math. Soc.*, 369(12):8537–8558, 2017.
- [vBJK19] Raymond van Bommel, Ariyan Javanpeykar, and Ljudmila Kamenova. Boundedness in families with applications to arithmetic hyperbolicity. *arXiv preprint arXiv:1907.11225*, 2019.
- [Voj87] Paul Vojta. *Diophantine Approximations and Value Distribution Theory*, volume 1239 of *Lecture Notes in Mathematics*. Springer Berlin Heidelberg, 1987.
- [Voj97] Paul Vojta. On Cartan’s theorem and Cartan’s conjecture. *Amer. J. Math.*, 119(1):1–17, 1997.
- [Voj11] Paul Vojta. Diophantine approximation and Nevanlinna theory. In *Arithmetic geometry*, volume 2009 of *Lecture Notes in Math.*, pages 111–224. Springer, Berlin, 2011.
- [Wan04] Julie Tzu-Yueh Wang. An effective Schmidt’s subspace theorem over function fields. *Math. Z.*, 246(4):811–844, 2004.
- [Yam15] Katsutoshi Yamanoi. Holomorphic curves in algebraic varieties of maximal Albanese dimension. *Internat. J. Math.*, 26(6):1541006, 45, 2015.

ERWAN ROUSSEAU

INSTITUT UNIVERSITAIRE DE FRANCE & AIX MARSEILLE UNIV, CNRS, CENTRALE MARSEILLE, I2M, MARSEILLE, FRANCE & FREIBURG INSTITUTE FOR ADVANCED STUDIES, UNIVERSITY OF FREIBURG, ALBERTSTR.19, 79104 FREIBURG, GERMANY

Email address: `erwan.rousseau@univ-amu.fr`

AMOS TURCHET

CENTRO DI RICERCA MATEMATICA ENNIO DE GIORGI, SCUOLA NORMALE SUPERIORE, PALAZZO PUTEANO, PIAZZA DEI CAVALIERI, 3, 56100 PISA, ITALY

Email address: `amos.turchet@sns.it`

JULIE TZU-YUEH WANG

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA NO. 1, SEC. 4, ROOSEVELT ROAD TAIPEI 10617, TAIWAN

Email address: `jwang@math.sinica.edu.tw`