

Asymptotic behavior of solutions of semilinear elliptic equations in unbounded domains: two approaches

Messoud Efendiev ^a and François Hamel ^{b*}

^a Helmholtz Zentrum München, Institut für Biomathematik und Biometrie

Ingolstädter Landstrasse 1, D-85764 Neuherberg, Germany

^b Aix-Marseille Université & Institut Universitaire de France

LATP, Avenue Escadrille Normandie-Niemen, F-13397 Marseille Cedex 20, France

Abstract

In this paper, we study the asymptotic behavior as $x_1 \rightarrow +\infty$ of solutions of semilinear elliptic equations in quarter- or half-spaces, for which the value at $x_1 = 0$ is given. We prove the uniqueness and characterize the one-dimensional or constant profile of the solutions at infinity. To do so, we use two different approaches. The first one is a pure PDE approach and it is based on the maximum principle, the sliding method and some new Liouville type results for elliptic equations in the half-space or in the whole space \mathbb{R}^N . The second one is based on the theory of dynamical systems.

Keywords. Semilinear elliptic equations; dynamical systems; asymptotic behavior; sliding method; nonlinear Liouville results.

1 Introduction

Let Ω be the domain of \mathbb{R}^N ($N \geq 2$) defined by

$$\begin{aligned}\Omega &= (0, +\infty) \times \mathbb{R}^{N-2} \times (0, +\infty) \\ &= \{x = (x_1, x', x_N) \in \mathbb{R}^N \mid x_1 > 0, x' = (x_2, \dots, x_{N-1}) \in \mathbb{R}^{N-2}, x_N > 0\}.\end{aligned}$$

This paper is devoted to the study of the large space behavior, that is as $x_1 \rightarrow +\infty$, of the nonnegative bounded classical solutions u of the equation

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \Omega, \\ u(x_1, x', 0) = 0 & \text{for all } x_1 > 0 \text{ and } x' \in \mathbb{R}^{N-2}, \\ u(0, x', x_N) = u_0(x', x_N) & \text{for all } x' \in \mathbb{R}^{N-2} \text{ and } x_N > 0, \end{cases} \quad (1.1)$$

*The second author is indebted to the Alexander von Humboldt Foundation for its support. He is also supported by the French ANR project PREFERED.

where the function

$$u_0 : \mathbb{R}^{N-2} \times (0, +\infty) \rightarrow \mathbb{R}_+ = [0, +\infty)$$

is given, continuous and bounded. The solutions u are understood to be bounded, of class $C^2(\Omega)$ and to be continuous on $\overline{\Omega} \setminus (\{0\} \times \mathbb{R}^{N-2} \times \{0\})$. From standard elliptic estimates, they are then automatically of class $C_b^{2,\beta}([\varepsilon, +\infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+)$ for all $\varepsilon > 0$ and $\beta \in [0, 1)$. Here and below, for any closed set $F \subset \mathbb{R}^N$ and $\beta \in [0, 1)$, we write

$$C_b^{2,\beta}(F) := \left\{ u : F \rightarrow \mathbb{R} \mid \|u\|_{C_b^{2,\beta}(F)} = \sup_{x \in F} \|u\|_{C^{2,\beta}(\overline{B(x,1)} \cap F)} < \infty \right\}, \quad (1.2)$$

where $B(x, 1)$ means the open Euclidean ball of radius 1 centered at x . Problems sets in the half-space

$$\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}$$

will also be considered in this paper, see (1.11) below.

The value of u at $x_1 = 0$ is then given and the goal is to describe the limiting profiles of u as $x_1 \rightarrow +\infty$. If the equation were parabolic in the variable x_1 , we would then be reduced to characterize the ω -limit set of the initial condition u_0 . However, problem (1.1) is an elliptic equation in all variables, including x_1 , and the ‘‘Cauchy’’ problem (1.1) with the ‘‘initial value’’ u_0 at $x_1 = 0$ is ill-posed. There might indeed be several solutions u with the same value u_0 at $x_1 = 0$. Nevertheless, under some assumptions on the nonlinearity f , we will see that the behavior as $x_1 \rightarrow +\infty$ of any solution u of (1.1) or of similar problems in the half-space $\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}$ is well-defined and unique (roughly speaking, no oscillation occur). In some cases, we will prove that all solutions u converge as $x_1 \rightarrow +\infty$ to the same limiting one-dimensional profile, irrespectively of u_0 . To do so, we will use two different approaches. The first one is a pure PDE approach based on comparisons with suitable sub-solutions and on Liouville type results. This paper indeed contains new Liouville type results of independent interest for the solutions of some elliptic equations in half-spaces $\mathbb{R}^{N-1} \times (0, +\infty)$ with homogeneous Dirichlet boundary conditions, or in the whole space \mathbb{R}^N (see Section 2 for more details). The second approach is a dynamical systems’ approach which says that x_1 can all the same be viewed as a time variable for a suitably defined dynamical system whose global attractor can be proved to exist and can be characterized.

Let us now describe more precisely the types of assumptions we make on the functions f , which are always assumed to be locally Lipschitz-continuous from \mathbb{R}_+ to \mathbb{R} . The first class of functions we consider corresponds to functions f such that

$$\left\{ \begin{array}{l} \exists \mu > 0, \quad f > 0 \text{ on } (0, \mu), \quad f \leq 0 \text{ on } [\mu, +\infty), \\ \exists 0 < \mu' < \mu, \quad f \text{ is nonincreasing on } [\mu', \mu], \\ \text{either } [f(0) > 0] \text{ or } [f(0) = 0 \text{ and } \liminf_{s \rightarrow 0^+} \frac{f(s)}{s} > 0]. \end{array} \right. \quad (1.3)$$

Under assumption (1.3) on f , it is immediate to see that there exists a unique solution $V \in C^2(\mathbb{R}_+)$ of the one-dimensional equation

$$\left\{ \begin{array}{l} V''(\xi) + f(V(\xi)) = 0 \text{ for all } \xi \geq 0, \\ V(0) = 0 < V(\xi) < \mu = V(+\infty) \text{ for all } \xi > 0. \end{array} \right. \quad (1.4)$$

Furthermore, $V'(\xi) > 0$ for all $\xi \geq 0$.

Under assumption (1.3), the behavior of the nontrivial solutions u of (1.1) as $x_1 \rightarrow +\infty$ is uniquely determined, as the following theorem shows.

Theorem 1.1 *Let N be any integer such that $N \geq 2$ and assume that f satisfies (1.3). Let u be any nonnegative and bounded solution of (1.1), where $u_0 : \mathbb{R}^{N-2} \times (0, +\infty) \rightarrow \mathbb{R}_+$ is any continuous and bounded function such that $u_0 \not\equiv 0$ in $\mathbb{R}^{N-2} \times (0, +\infty)$. Then*

$$\lim_{R \rightarrow +\infty} \left(\inf_{(R, +\infty) \times \mathbb{R}^{N-2} \times (R, +\infty)} u \right) \geq \mu \quad (1.5)$$

and

$$u(x_1 + h, x', x_N) \rightarrow V(x_N) \quad \text{as } h \rightarrow +\infty \text{ in } C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-2} \times [0, B]) \quad (1.6)$$

for all $A \in \mathbb{R}$, $B > 0$ and $\beta \in [0, 1)$, where $V \in C^2(\mathbb{R}_+)$ is the unique solution of (1.4).

Notice that property (1.5) means that the non-trivial nonnegative solutions u of (1.1) are separated from 0, irrespectively of u_0 , far away from the boundary $\partial\Omega$. If $u_0 \leq \mu$, then since $f \leq 0$ on $[\mu, +\infty)$ and $u = \mu$ on $\partial\Omega^+$, where $\Omega^+ = \Omega \cap \{u > \mu\}$, it follows that from the maximum principle applied in Ω^+ (see [5], since $\mathbb{R}^N \setminus \overline{\Omega^+}$ contains the closure of an infinite open connected cone), that actually $u \leq \mu$ in Ω^+ whence $\Omega^+ = \emptyset$ and $u \leq \mu$ in Ω . In this case, it also follows from Theorem 1.1 and standard elliptic estimates that the convergence (1.6) holds not only locally in x_N , but in $C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+)$ for all $A \in \mathbb{R}$ and $\beta \in [0, 1)$. However, without the assumption $u_0 \leq \mu$, it is not clear that this last convergence property holds globally with respect to x_N in general.

The second class of functions f we consider corresponds to the following assumption:

$$\begin{cases} f \geq 0 \text{ on } \mathbb{R}_+, \\ \forall z \in E, \liminf_{s \rightarrow z^+} \frac{f(s)}{s - z} > 0, \end{cases} \quad (1.7)$$

where

$$E = \{z \in \mathbb{R}_+; f(z) = 0\} \quad (1.8)$$

denotes the set of zeroes of f . A typical example of such a function f is $f(s) = |\sin s|$ for all $s \geq 0$, with $E = \pi\mathbb{N}$. More generally speaking, under the assumption (1.7), it follows immediately that the set E is at most countable. Furthermore, it is easy to check that, for each $z \in E \setminus \{0\}$, there exists a unique solution $V_z \in C^2(\mathbb{R}_+)$ of the one-dimensional equation

$$\begin{cases} V_z''(\xi) + f(V_z(\xi)) = 0 \text{ for all } \xi \geq 0, \\ V_z(0) = 0 < V_z(\xi) < z = V_z(+\infty) \text{ for all } \xi > 0. \end{cases} \quad (1.9)$$

Furthermore, $V_z'(\xi) > 0$ for all $\xi \geq 0$.

The following theorem states any solution of (1.1) is asymptotically one-dimensional as $x_1 \rightarrow +\infty$.

Theorem 1.2 *Let N be any integer such that $N \geq 2$ and assume that f satisfies (1.7). Let u be any nonnegative and bounded solution of (1.1), where $u_0 : \mathbb{R}^{N-2} \times (0, +\infty) \rightarrow \mathbb{R}_+$ is any continuous and bounded function such that $u_0 \not\equiv 0$ in $\mathbb{R}^{N-2} \times (0, +\infty)$. Then there exists $R > 0$ such that*

$$\inf_{(R, +\infty) \times \mathbb{R}^{N-2} \times (R, +\infty)} u > 0$$

and there exists $z \in E \setminus \{0\}$ such that

$$u(x_1 + h, x', x_N) \rightarrow V_z(x_N) \quad \text{as } h \rightarrow +\infty \text{ in } C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+) \quad (1.10)$$

for all $A \in \mathbb{R}$ and $\beta \in [0, 1)$, where $V_z \in C^2(\mathbb{R}_+)$ is the unique solution of (1.9) with the limit $V_z(+\infty) = z$.

This result shows that any non-trivial bounded solution u of (1.1) converges to a single one-dimensional profile as $x_1 \rightarrow +\infty$. More precisely, given u , the real number z defined by (1.10) is unique and, in the proof of Theorem 1.2, the explicit expression of z will be provided. Observe that the asymptotic profile may now depend on the solution u (unlike in Theorem 1.1) but Theorem 1.2 says that the oscillations in the x_1 variable are excluded at infinity, for any solution u .

The last two results are concerned with the analysis of the asymptotic behavior, as $x_1 \rightarrow +\infty$, of the nonnegative bounded classical solutions u of

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \Omega' = (0, +\infty) \times \mathbb{R}^{N-1}, \\ u(0, x_2, \dots, x_N) = u_0(x_2, \dots, x_N) & \text{for all } (x_2, \dots, x_N) \in \mathbb{R}^{N-1}, \end{cases} \quad (1.11)$$

in the half-space Ω' , where the function $u_0 : \mathbb{R}^{N-1} \rightarrow \mathbb{R}_+$ is given, continuous and bounded. The solutions u of (1.11) are understood to be bounded, of class $C^2(\Omega')$ and to be continuous on $\overline{\Omega'}$. They are then automatically of class $C_b^{2,\beta}([\varepsilon, +\infty) \times \mathbb{R}^{N-1})$ for all $\varepsilon > 0$ and $\beta \in [0, 1)$. Firstly, under the same assumptions (1.7) as in the previous theorem, the behavior as $x_1 \rightarrow +\infty$ of any non-trivial solution u of (1.11) is well-defined:

Theorem 1.3 *Let N be any integer such that $N \geq 2$ and assume that f satisfies (1.7). Let u be any nonnegative and bounded solution of (1.11), where $u_0 : \mathbb{R}^{N-1} \rightarrow \mathbb{R}_+$ is any continuous and bounded function such that $u_0 \not\equiv 0$ in \mathbb{R}^{N-1} . Then there exists $z \in E \setminus \{0\}$ such that*

$$u(x_1 + h, x_2, \dots, x_N) \rightarrow z \quad \text{as } h \rightarrow +\infty \text{ in } C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-1}) \quad (1.12)$$

for all $A \in \mathbb{R}$ and $\beta \in [0, 1)$.

Notice that the conclusion implies in particular that u is separated from 0 far away from the boundary $\{0\} \times \mathbb{R}^{N-1}$ of Ω' . Furthermore, as in Theorem 1.2, the real number z in (1.12) is uniquely determined by u and its explicit value will be given during the proof. In Theorem 1.3, if instead of (1.7) the function f now satisfies assumption (1.3), then u may not converge in general to a constant as $x_1 \rightarrow +\infty$. Furthermore, even if u does converge to a constant as $x_1 \rightarrow +\infty$, that constant may not be equal to the real number μ given

in (1.3). For instance, if there exists $\rho \in (\mu, +\infty)$ such that $f(\rho) = 0$, then the constant function $u = \rho$ solves (1.1) with $u_0 = \rho$. Therefore, under assumption (1.3), the asymptotic profile of a solution u of problem (1.11) in the half-space Ω' depends on u and is even not clearly well-defined in general. The situation is thus very different from Theorem 1.1 about the existence and uniqueness of the asymptotic behavior of the solutions of problem (1.1) in the quarter-space Ω .

However, under (1.3) and an additional appropriate assumption on f , the following result holds:

Theorem 1.4 *Let N be any integer such that $N \geq 2$ and assume that, in addition to (1.3), f is such that*

$$\liminf_{s \rightarrow z^-} \frac{f(s)}{s - z} > 0 \quad \text{for all } z > \mu \text{ such that } f(z) = 0. \quad (1.13)$$

Let u be any nonnegative and bounded solution of (1.11), where $u_0 : \mathbb{R}^{N-1} \rightarrow \mathbb{R}_+$ is any continuous and bounded function such that $u_0 \not\equiv 0$ in \mathbb{R}^{N-1} . Then there exists $z \geq \mu$ such that $f(z) = 0$ and

$$u(x_1 + h, x_2, \dots, x_N) \rightarrow z \quad \text{as } h \rightarrow +\infty \text{ in } C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-1})$$

for all $A \in \mathbb{R}$ and $\beta \in [0, 1)$.

Remark 1.5 It is worth noticing that all above results hold in any dimension $N \geq 2$.

2 The PDE approach

In this section, we use a pure PDE approach to prove the main results announced in Section 1. In Section 2.1, we deal with the case of problem (1.1) set in the quarter-space $\Omega = (0, +\infty) \times \mathbb{R}^{N-2} \times (0, +\infty)$, while Section 2.2 is concerned with problem (1.11) set in the half-space $\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}$.

2.1 Problem (1.1) in the quarter-space $\Omega = (0, +\infty) \times \mathbb{R}^{N-2} \times (0, +\infty)$

Let us first begin with the

Proof of Theorem 1.1. The proof is divided into three main steps: we first prove that u is bounded from below away from 0 when x_1 and x_N are large, uniformly with respect to $x' \in \mathbb{R}^{N-2}$. Then, we pass to the limit as $x_1 \rightarrow +\infty$ and use a classification result, which finally leads to the uniqueness and one-dimensional symmetry of the limiting profiles of u as $x_1 \rightarrow +\infty$.

Step 1. It consists in the following result, which we state as a lemma since it will be used several times in the paper. Notice that only the third line of (1.3) is needed.

Lemma 2.1 *Let N be any integer such that $N \geq 2$ and assume that either $f(0) > 0$, or $f(0) = 0$ and $\liminf_{s \rightarrow 0^+} f(s)/s > 0$. Let u be any nonnegative and bounded solution of (1.1), where $u_0 : \mathbb{R}^{N-2} \times (0, +\infty) \rightarrow \mathbb{R}_+$ is any continuous and bounded function such that $u_0 \not\equiv 0$ in $\mathbb{R}^{N-2} \times (0, +\infty)$. Then there exist $R > 0$ and $\varepsilon > 0$ such that*

$$u \geq \varepsilon \text{ in } [R, +\infty) \times \mathbb{R}^{N-2} \times [R, +\infty). \quad (2.14)$$

Proof. First of all, observe that, since $f(0) \geq 0$, the strong maximum principle implies that the function u is either positive in Ω , or identically equal to 0 in Ω . But since u is continuous up to $\{0\} \times \mathbb{R}^{N-2} \times (0, +\infty)$ and since $u_0 \not\equiv 0$, it follows that $u > 0$ in Ω .

In the sequel, for any $x \in \mathbb{R}^N$ and $R > 0$, denote $B(x, R)$ the open Euclidean ball of centre x and radius R . For each $R > 0$, let λ_R be the principal eigenvalue of the Laplace operator in $B(0, R)$ with Dirichlet boundary condition on $\partial B(0, R)$, and let φ_R be the normalized principal eigenfunction, that is

$$\begin{cases} \Delta \varphi_R + \lambda_R \varphi_R = 0 & \text{in } B(0, R), \\ \varphi_R > 0 & \text{in } B(0, R), \\ \|\varphi_R\|_{L^\infty(B(0, R))} = \varphi_R(0) = 1, & \\ \varphi_R = 0 & \text{on } \partial B(0, R). \end{cases} \quad (2.15)$$

Notice that $\lambda_R \rightarrow 0$ as $R \rightarrow +\infty$. If $f(0) = 0$, we can then choose $R > 0$ large enough so that

$$\lambda_R < \liminf_{s \rightarrow 0^+} \frac{f(s)}{s}.$$

If $f(0) > 0$, we simply choose $R = 1$. Then, fix a point $x_0 \in \Omega$ in such a way that $\overline{B(x_0, R)} \subset \Omega$. Since u is continuous and positive on $\overline{B(x_0, R)}$, there holds $\min_{\overline{B(x_0, R)}} u > 0$. Therefore, it follows from the choice of R that there exists $\varepsilon > 0$ small enough, such that the function

$$\underline{u}(x) = \varepsilon \varphi_R(x - x_0)$$

is a subsolution in $\overline{B(x_0, R)}$, that is

$$\Delta \underline{u} + f(\underline{u}) \geq 0 \text{ and } \underline{u} < u \text{ in } \overline{B(x_0, R)}. \quad (2.16)$$

Next, let \tilde{x}_0 be any point in Ω such that

$$\overline{B(\tilde{x}_0, R)} \subset \overline{\Omega},$$

that is $\tilde{x}_0 = (\tilde{x}_{0,1}, \tilde{x}'_0, \tilde{x}_{0,N})$ with $\tilde{x}_{0,1} \geq R$ and $\tilde{x}_{0,N} \geq R$. For all $t \in [0, 1]$, call

$$y_t = x_0 + t(\tilde{x}_0 - x_0)$$

and observe that $\overline{B(y_t, R)} \subset \Omega$ for all $t \in [0, 1]$. Define

$$\underline{u}_t(x) = \underline{u}(x - y_t + x_0) = \varepsilon \varphi_R(x - y_t)$$

for all $t \in [0, 1]$ and $x \in \overline{B(y_t, R)}$. By continuity and from (2.16), there holds $\underline{u}_t < u$ in $\overline{B(y_t, R)}$ for $t \in [0, t_0]$, where $t_0 > 0$ is small enough. On the other hand, for all $t \in [0, 1]$, the function \underline{u}_t is a subsolution of the equation satisfied by u , that is

$$\Delta \underline{u}_t + f(\underline{u}_t) \geq 0 \text{ in } \overline{B(y_t, R)}.$$

We shall now use a sliding method (see [5, 8]) to conclude that

$$\underline{u}_t < u \text{ in } \overline{B(y_t, R)} \text{ for all } t \in [0, 1).$$

Indeed, if this were not true, there would then exist a real number $t^* \in (0, 1)$ such that the inequality $\underline{u}_{t^*} \leq u$ holds in $\overline{B(y_{t^*}, R)}$ with equality at some point $x^* \in \overline{B(y_{t^*}, R)}$. Since $\overline{B(y_{t^*}, R)} \subset \Omega$, $u > 0$ in Ω and $\underline{u}_{t^*} = 0$ on $\partial B(y_{t^*}, R)$, one has $x^* \in B(y_{t^*}, R)$. But since \underline{u}_{t^*} is a subsolution of the equation satisfied by u , the strong maximum principle yields $\underline{u}_{t^*} = u$ in $B(y_{t^*}, R)$ and also on the boundary by continuity, which is impossible. One has then reached a contradiction. Hence, $\underline{u}_t < u$ in $\overline{B(y_t, R)}$ for all $t \in [0, 1)$. By continuity, one also gets that $\underline{u}_1 \leq u$ in $\overline{B(y_1, R)}$.

Therefore, for any $\tilde{x}_0 = (\tilde{x}_{0,1}, \tilde{x}'_0, \tilde{x}_{0,N})$ with $\tilde{x}_{0,1} \geq R$ and $\tilde{x}_{0,N} \geq R$, there holds

$$u(\tilde{x}_0) \geq \underline{u}_1(\tilde{x}_0) = \varepsilon \varphi_R(0) = \varepsilon.$$

In other words, (2.14) holds and the proof of Lemma 2.1 is complete. \square

Step 2. Let $(x_{1,n})_{n \in \mathbb{N}}$ be any sequence of positive numbers such that $x_{1,n} \rightarrow +\infty$ as $n \rightarrow +\infty$, and let $(x'_n)_{n \in \mathbb{N}}$ be any sequence in \mathbb{R}^{N-2} . From standard elliptic estimates, there exists a subsequence such that the functions

$$u_n(x) = u(x_1 + x_{1,n}, x' + x'_n, x_N)$$

converge in $C_{loc}^{2,\beta}(\mathbb{R}_+^N)$, for all $\beta \in [0, 1)$, to a bounded classical solution u_∞ of

$$\begin{cases} \Delta u_\infty + f(u_\infty) = 0 & \text{in } \mathbb{R}_+^N, \\ u_\infty = 0 & \text{on } \partial \mathbb{R}_+^N, \end{cases} \quad (2.17)$$

where $\mathbb{R}_+^N = \mathbb{R}^{N-1} \times [0, +\infty)$. Furthermore, $u_\infty \geq 0$ and $u_\infty \not\equiv 0$ in \mathbb{R}_+^N from Step 1, since

$$u_\infty \geq \varepsilon > 0 \text{ in } \mathbb{R}^{N-1} \times [R, +\infty)$$

from Lemma 2.1. Thus, $u_\infty > 0$ in $\mathbb{R}^{N-1} \times (0, +\infty)$ from the strong maximum principle. It follows from Theorems 1.1 and 1.2 of Berestycki, Caffarelli and Nirenberg [5]¹ (see also [1, 10]) that u_∞ is unique and has one-dimensional symmetry. By uniqueness of the problem (1.4), one gets that $u_\infty(x) = V(x_N)$ for all $x \in \mathbb{R}_+^N$, and the limit does not depend on the sequences $(x_{1,n})_{n \in \mathbb{N}}$ or $(x'_n)_{n \in \mathbb{N}}$. Property (1.6) of Theorem 1.1 then follows from

¹In [5], the function f was assumed to be globally Lipschitz-continuous. Here, f is just assumed to be locally Lipschitz-continuous. However, since u is bounded, it is always possible to find a Lipschitz-continuous function $\tilde{f} : \mathbb{R}^+ \rightarrow \mathbb{R}$ satisfying (1.3) and such that \tilde{f} and f coincide on the range of u .

the uniqueness of the limit.

Step 3. Let us now prove formula (1.5). One already knows from Lemma 2.1 that

$$m := \lim_{R \rightarrow +\infty} \left(\inf_{(R, +\infty) \times \mathbb{R}^{N-2} \times (R, +\infty)} u \right) \geq \varepsilon > 0.$$

Let $(x_n)_{n \in \mathbb{N}} = (x_{1,n}, x'_n, x_{N,n})_{n \in \mathbb{N}}$ be a sequence in Ω such that $(x_{1,n}, x_{N,n})_{n \in \mathbb{N}} \rightarrow (+\infty, +\infty)$ and $u(x_n) \rightarrow m$ as $n \rightarrow +\infty$. Up to extraction of a subsequence, the functions

$$v_n(x) = u(x + x_n)$$

converge in $C_{loc}^2(\mathbb{R}^N)$ to a classical bounded solution v_∞ of

$$\Delta v_\infty + f(v_\infty) = 0 \quad \text{in } \mathbb{R}^N$$

such that $v_\infty \geq m$ in \mathbb{R}^N and $v_\infty(0) = m > 0$. Thus, $f(m) \leq 0$, whence $m \geq \mu$ due to (1.3). The proof of Theorem 1.1 is thereby complete. \square

The proof of Theorem 1.2 is based on two Liouville type results for the bounded non-negative solutions u of the elliptic equation

$$\Delta u + f(u) = 0$$

in the whole space \mathbb{R}^N or in the half-space $\mathbb{R}^{N-1} \times \mathbb{R}_+$ with Dirichlet boundary condition on $\mathbb{R}^{N-1} \times \{0\}$.

Theorem 2.2 *Let N be any integer such that $N \geq 1$ and assume that the function f satisfies (1.7). Let u be a bounded nonnegative solution of*

$$\Delta u + f(u) = 0 \quad \text{in } \mathbb{R}^N. \tag{2.18}$$

Then u is constant.

The following result is concerned with the one-dimensional symmetry of nonnegative bounded solutions in a half-space with Dirichlet boundary conditions.

Theorem 2.3 *Let N be any integer such that $N \geq 1$ and assume that the function f satisfies (1.7). Let u be a bounded nonnegative solution of*

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \mathbb{R}_+^N = \mathbb{R}^{N-1} \times \mathbb{R}_+, \\ u = 0 & \text{on } \partial\mathbb{R}_+^N = \mathbb{R}^{N-1} \times \{0\}. \end{cases} \tag{2.19}$$

Then u is a function of x_N only. Furthermore, either $u = 0$ in $\mathbb{R}^{N-1} \times \mathbb{R}_+$ or there exists $z > 0$ such that $f(z) = 0$ and $u(x) = V_z(x_N)$ for all $x \in \mathbb{R}^{N-1} \times \mathbb{R}_+$, where the function V_z satisfies equation (1.9).

These results are of independent interest and will be proved in Section 3. Notice that one of the main points is that they hold in any dimensions $N \geq 1$ without any other assumption on u than its boundedness. In low dimensions $N \leq 4$, and under the additional assumption that u is stable, the conclusion of Theorem 2.2 holds for any nonnegative function f of class $C^1(\mathbb{R}_+)$, see Dupaigne and Farina [12]. Consequently, because of the monotonicity result in the direction x_N due to Berestycki, Caffarelli and Nirenberg [5] and Dancer [11] (since $f(0) \geq 0$), it follows that the conclusion of Theorem 2.3 holds for any nonnegative function f of class $C^1(\mathbb{R}_+)$, provided that $N \leq 5$, see Farina and Valdinoci [16]. However, observe that the nonnegativity and the C^1 character of f are incompatible with (1.7) for any positive zero z of f . Furthermore, assumption (1.7) is crucially used in the proof of Theorem 2.2 and 2.3. It is actually not true that these theorems stay valid in general when f is just assumed to be nonnegative and locally Lipschitz-continuous. For instance, non-constant solutions of (2.18), which are even stable, exist for power-like nonlinearities f in high dimensions (see [14] and the references therein).

With these results in hand, let us turn to the

Proof of Theorem 1.2. Observe that, from (1.7), either $f(0) > 0$, or $f(0) = 0$ and $\liminf_{s \rightarrow 0^+} f(s)/s > 0$. Therefore, Lemma 2.1 gives the existence of $R > 0$ and $\varepsilon > 0$ such that (2.14) holds. Set

$$M = \lim_{A \rightarrow +\infty} \left(\sup_{[A, +\infty) \times \mathbb{R}^{N-2} \times [0, +\infty)} u \right). \quad (2.20)$$

Our goal is to prove that the conclusion of Theorem 1.2 holds with $z = M$.

Since u is bounded and satisfies (2.14), M is such that $\varepsilon \leq M < +\infty$. Furthermore, there exists a sequence $(x_n)_{n \in \mathbb{N}} = (x_{1,n}, x'_n, x_{N,n})_{n \in \mathbb{N}}$ of points in $\bar{\Omega}$ such that $x_{1,n} \rightarrow +\infty$ and $u(x_n) \rightarrow M$ as $n \rightarrow +\infty$.

Assume first, up to extraction of a subsequence, that the sequence $(x_{N,n})_{n \in \mathbb{N}}$ converges to a nonnegative real number $x_{N,\infty}$ as $n \rightarrow +\infty$. From standard elliptic estimates, the functions

$$u_n(x) = u(x_1 + x_{1,n}, x' + x'_n, x_N)$$

converge in $C_{loc}^{2,\beta}(\mathbb{R}^{N-1} \times \mathbb{R}_+)$ for all $\beta \in [0, 1)$, up to extraction of another subsequence, to a bounded nonnegative solution u_∞ of the problem (2.19) in the half-space $\mathbb{R}^{N-1} \times [0, +\infty)$, such that

$$u_\infty(0, 0, x_{N,\infty}) = M = \sup_{\mathbb{R}^{N-1} \times [0, +\infty)} u_\infty > 0.$$

It follows from Theorem 2.3 that $u_\infty(x) = V_z(x_N)$ is a one-dimensional increasing solution of (1.9), whence $z = M$. But since V_M is (strictly) increasing, it cannot reach its maximum M at the finite point $x_{N,\infty}$. This case is then impossible.

Therefore, one can assume without loss of generality that $x_{N,n} \rightarrow +\infty$ as $n \rightarrow +\infty$. From standard elliptic estimates, the functions

$$u_n(x) = u(x + x_n)$$

converge in $C_{loc}^{2,\beta}(\mathbb{R}^N)$ for all $\beta \in [0, 1)$, up to extraction of another subsequence, to a bounded nonnegative solution u_∞ of the problem (2.18) in \mathbb{R}^N , such that

$$u_\infty(0) = M = \sup_{\mathbb{R}^N} u_\infty > 0.$$

Theorem 2.2 implies that $u_\infty = M$ in \mathbb{R}^N , whence $f(M) = 0$. Furthermore, since the limit M is unique, the convergence of the functions u_n to the constant M holds for the whole sequence.

Now, in order to complete the proof of Theorem 1.2, we shall make use of the following lemma of independent interest:

Lemma 2.4 *Let $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a locally Lipschitz-continuous nonnegative function. Then, for each $z > 0$ such that $g(z) = 0$ and for each $\varepsilon \in (0, z]$, there exist $R' = R'_{g,z,\varepsilon} > 0$ and a classical solution v of*

$$\left\{ \begin{array}{l} \Delta v + g(v) = 0 \text{ in } \overline{B(0, R')}, \\ 0 \leq v < z \text{ in } \overline{B(0, R')}, \\ v = 0 \text{ on } \partial B(0, R'), \\ v(0) = \max_{\overline{B(0, R')}} v \geq z - \varepsilon. \end{array} \right. \quad (2.21)$$

The proof of this lemma is postponed at the end of this section. Let us now finish the proof of Theorem 1.2. Fix an arbitrary ε in $(0, M]$. Let $R'(\varepsilon) = R'_{f,M,\varepsilon}$ be as in Lemma 2.4 and let v be a solution of (2.21) with $g = f$ and $z = M$. Since the functions $u_n(x) = u(x + x_n)$ converge locally uniformly in \mathbb{R}^N to the constant M and since

$$\max_{\overline{B(0, R'(\varepsilon))}} v < M,$$

there exists $n_0 \in \mathbb{N}$ large enough so that $\overline{B(x_{n_0}, R'(\varepsilon))} \subset \Omega$ and

$$v(x - x_{n_0}) < u(x) \text{ for all } x \in \overline{B(x_{n_0}, R'(\varepsilon))}. \quad (2.22)$$

But since v solves the same elliptic equation as u , the same sliding method as in Theorem 1.1 implies that

$$u \geq v(0) \geq M - \varepsilon \text{ in } [R'(\varepsilon), +\infty) \times \mathbb{R}^{N-2} \times [R'(\varepsilon), +\infty). \quad (2.23)$$

Lastly, choose any sequence $(\tilde{x}_{1,n})_{n \in \mathbb{N}}$ converging to $+\infty$, and any sequence $(\tilde{x}'_n)_{n \in \mathbb{N}}$ in \mathbb{R}^{N-2} . Up to extraction of a subsequence, the functions

$$\tilde{u}_n(x) = u(x_1 + \tilde{x}_{1,n}, x' + \tilde{x}'_n, x_N)$$

converge in $C_{loc}^{2,\beta}(\mathbb{R}^{N-1} \times \mathbb{R}_+)$ for all $\beta \in [0, 1)$ to a bounded nonnegative solution \tilde{u}_∞ of problem (2.19) in the half-space $\mathbb{R}^{N-1} \times \mathbb{R}_+$. The function \tilde{u}_∞ satisfies $\tilde{u}_\infty \leq M$ in $\mathbb{R}^{N-1} \times \mathbb{R}_+$ by definition of M , while

$$\lim_{A \rightarrow +\infty} \left(\inf_{\mathbb{R}^{N-1} \times [A, +\infty)} \tilde{u}_\infty \right) \geq M$$

because $\varepsilon > 0$ in (2.23) can be arbitrarily small. Theorem 2.3 and the above estimates imply that

$$\tilde{u}_\infty(x) = V_M(x_N) \quad \text{for all } x \in \mathbb{R}^{N-1} \times \mathbb{R}_+.$$

Since this limit does not depend on any subsequence, and due to (2.20), (2.23) and standard elliptic estimates, it follows in particular that

$$u(x_1 + h, x', x_N) \rightarrow V_M(x_N) \quad \text{in } C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+) \quad \text{as } h \rightarrow +\infty,$$

for all $A \in \mathbb{R}$ and $\beta \in [0, 1)$. The proof of Theorem 1.2 is thereby complete. \square

Remark 2.5 Instead of Theorem 2.2, if f is just assumed to be nonnegative and locally Lipschitz-continuous on \mathbb{R}_+ and if u_∞ is a solution of (2.18) which reaches its maximum and is such that $f(\max_{\mathbb{R}^N} u_\infty) = 0$ (as in some assumptions of [4, 6]), then u_∞ is constant, from the strong maximum principle. Therefore, it follows from similar arguments as in the above proof that if, instead of (1.7) in Theorem 1.2, the function f is just assumed to be nonnegative, locally Lipschitz-continuous and positive almost everywhere on \mathbb{R}_+ and if u is a bounded nonnegative solution of (1.1) such that $f(M) = 0$, where M is defined by (2.20), then either $M = 0$ and $u(x_1 + h, x', x_N) \rightarrow 0$ in $C^2([A, +\infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+)$ as $h \rightarrow +\infty$ for all $A \in \mathbb{R}$, or the conclusion (1.10) holds with $z = M$.

Proof of Lemma 2.4. The proof uses classical variational arguments, which we sketch here for the sake of completeness (see also e.g. [7, 13] for applications of this method). Let g and z be as in Lemma 2.4 and let \tilde{g} be the function defined in \mathbb{R} by

$$\tilde{g}(s) = \begin{cases} g(0) & \text{if } s < 0, \\ g(s) & \text{if } 0 \leq s \leq z, \\ 0 & \text{if } s > z. \end{cases}$$

The function \tilde{g} is nonnegative, bounded and Lipschitz-continuous on \mathbb{R} . Set

$$G(s) = \int_s^z \tilde{g}(\tau) d\tau \geq 0$$

for all $s \in \mathbb{R}$. The function G is nonnegative and Lipschitz-continuous on \mathbb{R} .

Let r be any positive real number. Define

$$I_r(v) = \frac{1}{2} \int_{B(0,r)} |\nabla v|^2 + \int_{B(0,r)} G(v)$$

for all $v \in H_0^1(B(0, r))$. The functional I_r is well-defined in $H_0^1(B(0, r))$ and it is coercive, from Poincaré's inequality and the nonnegativity of G . From Rellich's and Lebesgue's theorems, the functional I_r has a minimum v_r in $H_0^1(B(0, r))$. The function v_r is a weak and hence, from the elliptic regularity theory, a classical $C^2(\overline{B(0, r)})$ solution of the equation

$$\begin{cases} \Delta v_r + \tilde{g}(v_r) = 0 & \text{in } \overline{B(0, r)}, \\ v_r = 0 & \text{on } \partial B(0, r). \end{cases}$$

Since $\tilde{g} \geq 0$ on $(-\infty, 0]$, it follows from the strong maximum principle that $v_r \geq 0$ in $\overline{B(0, r)}$. Furthermore, either $v_r = 0$ in $\overline{B(0, r)}$, or $v_r > 0$ in $B(0, r)$. Similarly, since $\tilde{g} = 0$ on $[z, +\infty)$, one gets that $v_r < z$ in $\overline{B(0, r)}$. Consequently, $\tilde{g}(v_r) = g(v_r)$ in $\overline{B(0, r)}$. It also follows from the method of moving planes and Gidas, Ni and Nirenberg [17] that v_r is radially symmetric and decreasing with respect to $|x|$ (provided that $v_r \not\equiv 0$ in $B(0, r)$). In all cases, there holds

$$0 \leq v_r(0) = \max_{\overline{B(0, r)}} v_r < z.$$

In order to complete the proof of Lemma 2.4, it is sufficient to prove that, given ε in $(0, z]$, there exists $r > 0$ such that $v_r(0) \geq z - \varepsilon$. Let $\varepsilon \in (0, z]$ and assume that $\max_{\overline{B(0, r)}} v_r = v_r(0) < z - \varepsilon$ for all $r > 0$. Observe that the function G is non-increasing in \mathbb{R} , and actually decreasing and positive on the interval $[0, z]$, from (1.7). Therefore,

$$I_r(v_r) \geq \alpha_N r^N G(z - \varepsilon) \quad (2.24)$$

for all $r > 0$, where $\alpha_N > 0$ denotes the Lebesgue measure of the unit Euclidean ball in \mathbb{R}^N . For $r > 1$, let w_r be the test function defined in $\overline{B(0, r)}$ by

$$w_r(x) = \begin{cases} z & \text{if } |x| < r - 1, \\ z(r - |x|) & \text{if } r - 1 \leq |x| \leq r. \end{cases}$$

This function w_r belongs to $H_0^1(B(0, r))$ and $|\nabla w_r|^2$ and $G(w_r)$ are supported on the shell $\overline{B(0, r)} \setminus B(0, r - 1)$. Thus, there exists a constant C independent of r such that

$$I_r(w_r) \leq C(r^n - (r - 1)^n) \quad (2.25)$$

for all $r > 1$. But since $I_r(v_r) \leq I_r(w_r)$, by definition of v_r , and since $G(z - \varepsilon) > 0$, inequalities (2.24) and (2.25) lead to a contradiction as $r \rightarrow +\infty$. Therefore, there exists a radius $R' > 0$ such that $v_{R'}(0) \geq z - \varepsilon$, and $v_{R'}$ solves (2.21). The proof of Lemma 2.4 is now complete. \square

2.2 Problem (1.11) in the half-space $\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}$

Let us now turn to problem (1.11) set in the half-space $\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}$. This section is devoted to the proof of Theorems 1.3 and 1.4. A useful ingredient in the proof of these two theorems is the following result, which is the analogue of Lemma 2.1 in the case of half-spaces. Its proof is very similar to that of Lemma 2.1 and it is left to the reader.

Lemma 2.6 *Let N be any integer such that $N \geq 2$ and assume that either $f(0) > 0$, or $f(0) = 0$ and $\liminf_{s \rightarrow 0^+} f(s)/s > 0$. Let u be any nonnegative and bounded solution of (1.11), where $u_0 : \mathbb{R}^{N-1} \rightarrow \mathbb{R}_+$ is any continuous and bounded function such that $u_0 \not\equiv 0$ in \mathbb{R}^{N-1} . Then there exist $R > 0$ and $\varepsilon > 0$ such that*

$$u \geq \varepsilon \text{ in } [R, +\infty) \times \mathbb{R}^{N-1}.$$

Proof of Theorem 1.3. Assume here that f satisfies (1.7). First of all, it follows from Lemma 2.6 that there exists $R > 0$ such that $\inf_{[R, +\infty) \times \mathbb{R}^{N-1}} u > 0$. Call

$$M = \lim_{A \rightarrow +\infty} \left(\sup_{[A, +\infty) \times \mathbb{R}^{N-1}} u \right) > 0.$$

We shall now prove that the conclusion of Theorem 1.3 holds with $z = M$. Choose a sequence $(x_n)_{n \in \mathbb{N}} = (x_{1,n}, \dots, x_{N,n})_{n \in \mathbb{N}}$ in Ω' such that $x_{1,n} \rightarrow +\infty$ and $u(x_n) \rightarrow M$ as $n \rightarrow +\infty$. As in the proof of Theorem 1.2, it follows from Theorem 2.2 that the functions

$$u_n(x) = u(x + x_n)$$

converge in $C_{loc}^{2,\beta}(\mathbb{R}^N)$ for all $\beta \in [0, 1)$ to the constant M , whence $f(M) = 0$.

Then, for any $\varepsilon \in (0, M]$, let $R'(\varepsilon) = R'_{f,M,\varepsilon}$ be as in Lemma 2.4 and let v be a solution of (2.21) with $g = f$ and $z = M$. As in the proof of Theorem 1.2, there exists $n_0 \in \mathbb{N}$ large enough so that $\overline{B(x_{n_0}, R'(\varepsilon))} \subset \Omega'$ and (2.22) holds. The sliding method yields

$$u \geq v(0) \geq M - \varepsilon \quad \text{in } [R'(\varepsilon), +\infty) \times \mathbb{R}^{N-1}.$$

Since $\varepsilon > 0$ is arbitrarily small, the definition of M implies that, for all $A \in \mathbb{R}$,

$$u(x_1 + h, x_2, \dots, x_N) \rightarrow M \quad \text{as } h \rightarrow +\infty$$

uniformly with respect to $(x_1, x_2, \dots, x_N) \in [A, +\infty) \times \mathbb{R}^{N-1}$. The convergence also holds in $C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-1})$ for all $\beta \in [0, 1)$ from standard elliptic estimates. The proof of Theorem 1.3 is thereby complete. \square

In the case when f satisfies (1.3) and (1.13), the conclusion is similar to that of Theorem 1.3, as the following proof of Theorem 1.4 will show. As a matter of fact, it is also based on a Liouville type result for the bounded nonnegative solutions of (2.18), which is the counterpart of Theorem 2.2 under assumptions (1.3) and (1.13).

Theorem 2.7 *Let N be any integer such that $N \geq 1$ and assume that the function f satisfies (1.3) and (1.13). Then any bounded nonnegative solution u of (2.18) is constant.*

The proof is postponed in Section 3 and we now complete the

Proof of Theorem 1.4. First of all, it follows from Lemma 2.6 that there exist $\varepsilon > 0$ and $R > 0$ such that $u \geq \varepsilon$ in $[R, +\infty) \times \mathbb{R}^{N-1}$. Call now

$$m = \lim_{A \rightarrow +\infty} \left(\inf_{[A, +\infty) \times \mathbb{R}^{N-1}} u \right).$$

One has $m \in [\varepsilon, +\infty)$. Let $(x_n)_{n \in \mathbb{N}} = (x_{1,n}, \dots, x_{N,n})_{n \in \mathbb{N}}$ be a sequence in Ω' such that $x_{1,n} \rightarrow +\infty$ and $u(x_n) \rightarrow m$ as $n \rightarrow +\infty$. Up to extraction of a subsequence, the functions

$$u_n(x) = u(x + x_n)$$

converge in $C_{loc}^{2,\beta}(\mathbb{R}^N)$ for all $\beta \in [0, 1)$ to a classical bounded solution u_∞ of (2.18) such that $u_\infty \geq m > 0$ in \mathbb{R}^N and $u_\infty(0) = m$. Theorem 2.7 then implies that u_∞ is constant in \mathbb{R}^N , wence it is identically equal to m and $f(m) = 0$.

Call now

$$M' = \sup_{\Omega'} u$$

and let $g : [0, +\infty) \rightarrow \mathbb{R}$ be the function defined by

$$g(s) = \begin{cases} -f(M' + 1 - s) & \text{if } 0 \leq s \leq M' + 1 - m, \\ 0 & \text{if } s > M' + 1 - m. \end{cases}$$

The function g is Lipschitz-continuous and nonnegative. The real number

$$z = M' + 1 - m$$

is positive and fulfills $g(z) = f(m) = 0$. Choose any ε in $(0, z]$. From Lemma 2.4, there exist $R' > 0$ and a classical solution v of (2.21) in $\overline{B(0, R')}$, that is

$$\begin{cases} \Delta v + g(v) = 0 & \text{in } \overline{B(0, R')}, \\ 0 \leq v < z & \text{in } \overline{B(0, R')}, \\ v = 0 & \text{on } \partial B(0, R'), \\ v(0) = \max_{\overline{B(0, R')}} v \geq z - \varepsilon. \end{cases}$$

The function $V = M' + 1 - v$ then satisfies

$$\begin{cases} \Delta V + f(V) = 0 & \text{in } \overline{B(0, R')}, \\ m < V \leq M' + 1 & \text{in } \overline{B(0, R')}, \\ V = M' + 1 & \text{on } \partial B(0, R'), \\ V(0) = \min_{\overline{B(0, R')}} V \leq m + \varepsilon. \end{cases}$$

Since $u_n(x) = u(x + x_n) \rightarrow u_\infty(x) = m$ as $n \rightarrow +\infty$ in $C_{loc}^{2,\beta}(\mathbb{R}^N)$ for all $\beta \in [0, 1)$, it follows that there exists $n_0 \in \mathbb{N}$ large enough so that $\overline{B(x_{n_0}, R')} \subset \Omega'$ and

$$V(x - x_{n_0}) > u(x) \quad \text{for all } x \in \overline{B(x_{n_0}, R')}.$$

Since $V = \sup_{\Omega'} u + 1 > \sup_{\Omega'} u$ on $\partial B(0, R')$, it follows from the elliptic maximum principle and the sliding method that

$$u(x) \leq V(0) \leq m + \varepsilon \quad \text{for all } x \in [R', +\infty) \times \mathbb{R}^{N-1}.$$

Owing to the definition of m , one concludes that, for all $A \in \mathbb{R}$,

$$u(x_1 + h, x_2, \dots, x_N) \rightarrow m \quad \text{as } h \rightarrow +\infty$$

uniformly with respect to $(x_1, x_2, \dots, x_N) \in [A, +\infty) \times \mathbb{R}^{N-1}$ and the convergence holds in $C_b^{2,\beta}([A, +\infty) \times \mathbb{R}^{N-1})$ for all $\beta \in [0, 1)$ from standard elliptic estimates. The proof of Theorem 1.4 is thereby complete. \square

3 Classification results in the whole space \mathbb{R}^N or in the half-space $\mathbb{R}^{N-1} \times (0, +\infty)$ with Dirichlet boundary conditions

This section is devoted to the proof of the Liouville type results for the bounded nonnegative solutions u of problems (2.18) or (2.19). Theorems 2.3 and 2.7 are actually corollaries of Theorem 2.2. We then begin with the proof of the latter.

Proof of Theorem 2.2. Let u be a bounded nonnegative solution of (2.18) under assumption (1.7). Denote

$$m = \inf_{\mathbb{R}^N} u \geq 0.$$

Since $f(m) \geq 0$, the constant m is a subsolution for (2.18). It follows from the strong elliptic maximum principle that either $u = m$ in \mathbb{R}^N , or $u > m$ in \mathbb{R}^N .

Let us prove that the second case, that is $u > m$, is impossible. That will give the desired conclusion. Assume that $u > m$ in \mathbb{R}^N and let us get a contradiction. Let us first check that

$$f(m) = 0. \tag{3.26}$$

This could be done by considering a sequence along which u converges to its minimum; after changing the origin, the limiting function would be identically equal to m from the strong maximum principle, which would yield (3.26). Let us choose an alternate elementary parabolic argument. Assume that $f(m) > 0$ and let $\xi : [0, T) \rightarrow \mathbb{R}$ be the maximal solution of

$$\begin{cases} \xi'(t) = f(\xi(t)) & \text{for all } t \in [0, T), \\ \xi(0) = m. \end{cases}$$

The maximal existence time T satisfies $0 < T \leq +\infty$ (and $T = +\infty$ if f is globally Lipschitz-continuous). Since $\xi(0) \leq u$ in \mathbb{R}^N , it follows from the parabolic maximum principle for the equation $v_t = \Delta v + f(v)$, satisfied by both ξ and u in $[0, T) \times \mathbb{R}^N$, that

$$\xi(t) \leq u(x) \text{ for all } x \in \mathbb{R}^N \text{ and } t \in [0, T).$$

But $\xi'(0) = f(\xi(0)) = f(m) > 0$. Hence, there exists $\tau \in (0, T)$ such that $\xi(\tau) > m$, whence $u(x) \geq \xi(\tau) > m$ for all $x \in \mathbb{R}^N$, which contradicts the definition of m .

Therefore, (3.26) holds. Now, as in the proof of Lemma 2.1, because of property (1.7) at $z = m$, there exist $R > 0$ and $\varepsilon > 0$ such that

$$\Delta(m + \varepsilon \varphi_R) + f(m + \varepsilon \varphi_R) \geq 0 \text{ in } B(0, R)$$

and $m + \varepsilon \varphi_R < u$ in $\overline{B(0, R)}$, where φ_R solving (2.15) is the principal eigenfunction of the Dirichlet-Laplace operator in $B(0, R)$. Since $m + \varepsilon \varphi_R = m$ on $\partial B(0, R)$ and $u > m$ in \mathbb{R}^N , the same sliding method as in the proof of Lemma 2.1 implies that

$$m + \varepsilon \varphi_R(x - y) \leq u(x) \text{ for all } x \in \overline{B(y, R)}$$

and for all $y \in \mathbb{R}^N$. Therefore, $u \geq m + \varepsilon \varphi_R(0) > m$ in \mathbb{R}^N , which contradicts the definition of m .

As a conclusion, the assumption $u > m$ is impossible and, as already emphasized, the proof of Theorem 2.2 is thereby complete. \square

The proof of Theorem 2.3 also uses the sliding method and Theorem 2.2, combined with limiting arguments as $x_N \rightarrow +\infty$ and comparison with non-small subsolutions.

Proof of Theorem 2.3. Let u be a bounded nonnegative solution of (2.19) under assumption (1.7). Since $f(0) \geq 0$, it follows from the strong maximum principle that either $u = 0$ in $\mathbb{R}^{N-1} \times \mathbb{R}_+$, or $u > 0$ in $\mathbb{R}^{N-1} \times (0, +\infty)$. Let us then consider the second case. Since $f(0) \geq 0$, it follows from Corollary 1.3 of Berestycki, Caffarelli and Nirenberg [6] that u is increasing in x_N . Denote

$$M = \sup_{\mathbb{R}^{N-1} \times [0, +\infty)} u.$$

There exists a sequence $(x_n)_{n \in \mathbb{N}} = (x_{1,n}, \dots, x_{N,n})_{n \in \mathbb{N}}$ in $\mathbb{R}^{N-1} \times \mathbb{R}_+$ such that $x_{N,n} \rightarrow +\infty$ and $u(x_n) \rightarrow M$ as $n \rightarrow +\infty$. From standard elliptic estimates, the functions

$$u_n(x) = u(x + x_n),$$

which satisfy the same equation as u , converge in $C_{loc}^{2,\beta}(\mathbb{R}^N)$ for all $\beta \in [0, 1)$, up to extraction of a subsequence, to a solution u_∞ of (2.18) such that $u_\infty(0) = M$. Theorem 2.2 implies that

$$u_\infty = M \text{ in } \mathbb{R}^N,$$

whence $f(M) = 0$. It follows then from Berestycki, Caffarelli and Nirenberg [4] (see also Theorem 1.4 in [6]) that u depends on x_N only. In other words, the function $u(x)$ is equal to $V_M(x_N)$, which completes the proof of Theorem 2.3. \square

Let us complete this section with the

Proof of Theorem 2.7. Assume that the function f satisfies (1.3) and (1.13) and let u be a bounded nonnegative solution of (2.18). As already underlined, it first follows from the strong maximum principle that either $u \equiv 0$ in \mathbb{R}^N , or $u > 0$ in \mathbb{R}^N . Let us then consider the second case. Applying the sliding method and using the same notations as in the proof of Lemma 2.1, one gets the existence of $R > 0$ and $\varepsilon > 0$ such that $u(x) \geq \varepsilon \varphi_R(x - y)$ for all $x \in \overline{B(y, R)}$ and for all $y \in \mathbb{R}^N$, whence

$$m = \inf_{\mathbb{R}^N} u > 0.$$

Let $(x_n)_{n \in \mathbb{N}}$ be a sequence in \mathbb{R}^N such that $u(x_n) \rightarrow m$ as $n \rightarrow +\infty$. Up to extraction of a subsequence, the functions

$$u_n(x) = u(x + x_n)$$

converge in $C_{loc}^{2,\beta}(\mathbb{R}^N)$ for all $\beta \in [0, 1)$ to a classical bounded solution u_∞ of (2.18) in \mathbb{R}^N such that $u_\infty \geq m$ in \mathbb{R}^N and $u_\infty(0) = m$. Hence,

$$f(m) \leq 0,$$

whence $m \geq \mu$ from (1.3) and since $m > 0$.

Call now

$$M = \sup_{\mathbb{R}^N} u.$$

If $M = m$, then u is constant, which is the desired result. Assume now that $M > m$. The function

$$v = M - u$$

is a nonnegative bounded solution of

$$\Delta v + g(v) = 0 \quad \text{in } \mathbb{R}^N,$$

where the function $g : [0, +\infty) \rightarrow \mathbb{R}$ is defined by

$$g(s) = \begin{cases} -f(M - s) & \text{if } 0 \leq s \leq M - m, \\ -f(m) & \text{if } s > M - m. \end{cases}$$

Because of (1.3) and (1.13), the function g is Lipschitz-continuous and fulfills property (1.7). Theorem 2.2 applied to g and v implies that the function v is actually constant. Hence, u is also constant, which actually shows that the assumption $M > m$ is impossible. As a conclusion, $M = m$ and u is then constant. \square

4 The dynamical systems' approach

The goal of this section is to apply the dynamical systems' (shortly DS) approach to study the symmetrization and stabilization (as $x_1 \rightarrow +\infty$) properties of the nonnegative solutions (1.1) in

$$\Omega = (0, +\infty) \times \mathbb{R}^{N-2} \times (0, +\infty)$$

and (1.11) in

$$\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}.$$

To this end we apply as aforementioned the DS approach. One of the main difficulties which arises in the dynamical study of (1.1) in Ω or (1.11) in Ω' is the fact that the corresponding Cauchy problem is not well posed for (1.1) in Ω and for (1.11) in Ω' , and consequently the straightforward interpretation of (1.1) and (1.11) as an evolution equation leads to semigroups of multivalued maps even in the case of cylindrical domains, see [2]. The usage of multivalued maps can be overcome using the so-called trajectory dynamical approach (see [3, 9, 18] and the references therein). Under this approach, one fixes a signed direction \vec{l} in \mathbb{R}^N , which will play role of time. Then the space K^+ of all bounded nonnegative classical solutions of (1.1) in Ω or (1.11) in Ω' (in the sense described in Section 1) is considered

as a trajectory phase space for the semi-flow $(T_h^{\vec{l}})_{h \in \mathbb{R}_+}$ of translations along the direction \vec{l} defined via

$$(T_h^{\vec{l}}u)(x) = u(x + h\vec{l}), \quad h \in \mathbb{R}_+, \quad u \in K^+.$$

In order the trajectory dynamical system $(T_h^{\vec{l}}, K^+)$ to be well defined, one needs the domains Ω and Ω' to be invariant with respect to positive translations along the \vec{l} directions, that is

$$T_h^{\vec{l}}(\Omega) \subset \Omega \quad (\text{resp. } T_h^{\vec{l}}(\Omega') \subset \Omega'), \quad T_h^{\vec{l}}(x) := x + h\vec{l},$$

for all $h \geq 0$. In our case, the x_1 -axis will play the role of time, that is $\vec{l} = (1, 0, \dots, 0, 0)$. For the sake of simplicity of the notation, we then set

$$T_h^{\vec{l}} = T_h.$$

To apply the DS approach for our purposes, we apply the following Lemma 4.1 (see below), which also has an independent interest. For that purpose, let us introduce a few more notations. For any locally Lipschitz-continuous function f from \mathbb{R}_+ to \mathbb{R} , such that $f(0) \geq 0$, let Z_f be defined by

$$Z_f = \{z_0 \in \mathbb{R}_+ \mid f(z_0) = 0 \text{ and } F(z) < F(z_0) \text{ for all } z \in [0, z_0)\}, \quad (4.27)$$

where

$$F(z) = \int_0^z f(\sigma) d\sigma.$$

The set Z_f is then a subset of the set E of zeroes of f , defined in (1.8). Lastly, by R_f we denote the set of all bounded, nonnegative solutions $V \in C^2(\mathbb{R}_+)$ of

$$\begin{cases} V''(\xi) + f(V(\xi)) = 0 \text{ for all } \xi \geq 0, \\ V(0) = 0, \quad V \geq 0, \quad V \text{ is bounded.} \end{cases} \quad (4.28)$$

Lemma 4.1 *Let f be a locally Lipschitz-continuous function from \mathbb{R}_+ to \mathbb{R} , such that $f(0) \geq 0$. Then the set R_f is homeomorphic to Z_f and as a consequence is totally disconnected.*

Proof. Since $f(0) \geq 0$, it follows from elementary arguments that every nonnegative bounded solution of (4.28) is monotonically nondecreasing, that is $V(\xi_1) \geq V(\xi_2)$ if $\xi_1 \geq \xi_2$. Consequently, the following limit

$$z_0 = z_0(V) = \lim_{\xi \rightarrow +\infty} V(\xi) \quad (4.29)$$

exists and necessarily $f(z_0) = 0$ and $0 \leq V(z) \leq z_0$. Moreover, it is well known that either $V(\xi) = 0$ for all $\xi \geq 0$, or $V'(\xi) > 0$ for all $\xi \geq 0$.

Multiplying equation (4.28) by V' and integrating over $[0, \xi]$ we obtain the explicit expression for the derivative $V'(\xi)$:

$$V'(\xi)^2 = -2F(V(\xi)) + V'(0)^2 \quad \text{for all } \xi \geq 0. \quad (4.30)$$

This formula corresponds to the classical conservation of energy principle for the ODE (4.28) (we refer to [15] for much more general extensions of this principle and its use for classification results for PDEs). Passing to the limit $\xi \rightarrow +\infty$ in (4.30) and taking into account (4.29), it follows that $V'(0)^2 = 2F(z_0)$, whence $F(z_0) \geq 0$. Therefore we obtain the following equation for $V(\xi)$:

$$V'(\xi)^2 = 2(F(z_0) - F(V(\xi))) \quad \text{for all } \xi \geq 0. \quad (4.31)$$

Assume now that $F(z_0) > 0$, whence $z_0 > 0$ and then $V' > 0$ in \mathbb{R}_+ . Then the solution V_{z_0} of (4.28) that satisfies (4.29) and (4.31) exists if and only if $F(z_0) - F(z) > 0$ for every $z \in [0, z_0]$. Moreover, such a solution is unique because V_{z_0} satisfies (4.28) with the initial conditions

$$V_{z_0}(0) = 0, V'_{z_0}(0) = \sqrt{2F(z_0)}. \quad (4.32)$$

If $z_0 = 0$, then $V'(0) = 0$ and $V(\xi) = 0$ for all $\xi \in \mathbb{R}_+$, whence $f(0) = 0$. In this case, we define $V_0 = 0$ in \mathbb{R}_+ .

Next we show that Z_f defined by (4.27) is totally disconnected in \mathbb{R} . Indeed, otherwise it should contain a segment $[a, b]$ with $0 \leq a < b$. Then, $f(z_0) \equiv 0$ for $z_0 \in [a, b]$ and consequently $F(z_0) = F(b)$ for every $z_0 \in [a, b]$, which evidently leads to a contradiction.

To prove the disconnectedness of the set R_f it is then sufficient to show that there exists a homeomorphism

$$\tau : (Z_f, \mathbb{R}) \rightarrow (R_f, C_{loc}^2(\mathbb{R}_+)).$$

To do so, observe that (4.32) defines a homeomorphism between Z_f and the set

$$R_f(0) = \{(0, V'(0)) \mid V \in R_f\}$$

of values at $\xi = 0$ of the functions from R_f . Recall that R_f consists of the solutions of the second order ODE (4.28) and, consequently, thanks to the classical Cauchy-Lipschitz theorem on continuous dependence of solutions of ODE's, the set R_f is homeomorphic to $R_f(0)$ and this homeomorphism is given by $V \mapsto (0, V'(0))$. As a conclusion, the set R_f is totally disconnected. \square

Remark 4.2 Note that the homeomorphic sets Z_f and R_f may be uncountable for some very special choices of nonlinearities f . One of the simplest examples of such a function f is the following one:

$$f(z) := \text{dist}(z, \mathcal{C})$$

for all $z \in \mathbb{R}_+$, where \mathcal{C} is a standard Cantor set on $[0, 1]$ and $\text{dist}(z, \mathcal{C})$ means a distance from z to \mathcal{C} . Indeed, it is easy to verify that for this case $Z_f = \mathcal{C}$ and consequently R_f consists of a continuum of elements. One can also construct examples of C^k or even C^∞ functions f for which $Z_f = \mathcal{C}$: for instance, define $f(z) = e^{-1/(z-1)}$ for all $z > 1$, $f(z) = 0$ for all $z \in \mathcal{C}$, and $f(z) = e^{-1/((z-a)(b-z))}$ for all $z \in (a, b)$ and for all $0 < a < b < 1$ such that $a, b \in \mathcal{C}$ and $(a, b) \subset (0, 1) \setminus \mathcal{C}$.

Below we state the main results of this Section 4, that is Theorems 4.3-4.6, which are obtained by the dynamical systems' approach. To this end we define a class of functions K^+ to which the solutions of (1.1) as well as (1.11) belong to. Namely, a bounded nonnegative solution of (1.1) (resp. (1.11)) is understood to be a solution u of class $C^2(\Omega)$ (resp. $C^2(\Omega')$) and continuous on $\bar{\Omega} \setminus \{0\} \times \mathbb{R}^{N-2} \times \{0\}$ (resp. on $\bar{\Omega}'$). The set K^+ is endowed with the local topology according to the embedding of K^+ in $C_{loc}^{2,\beta}(\bar{\Omega} \setminus \{x_1 = 0\})$ (resp. $C_{loc}^{2,\beta}(\bar{\Omega}' \setminus \{x_1 = 0\})$) for all $\beta \in [0, 1)$. Actually, as already emphasized in Section 1, all solutions $u \in K^+$ are automatically in $C_b^{2,\beta}([\varepsilon, +\infty) \times \mathbb{R}^{N-2} \times [0, +\infty))$ (resp. $C_b^{2,\beta}([\varepsilon, +\infty) \times \mathbb{R}^{N-1})$) for all $\beta \in [0, 1)$ and for all $\varepsilon > 0$, where we refer to (1.2) for the definition of the sets $C_b^{2,\beta}(F)$.

The first two theorems are concerned with the case of functions f fulfilling the condition (1.3).

Theorem 4.3 *Let N be any integer such that $N \geq 2$ and let f be a locally Lipschitz-continuous function from \mathbb{R}_+ to \mathbb{R} , satisfying (1.3). Then the trajectory dynamical system (T_h, K^+) associated to (1.1) possesses a global attractor A_{tr} in K^+ which is bounded in $C_b^{2,\beta}(\bar{\Omega})$ and then compact in $C_{loc}^{2,\beta}(\bar{\Omega})$ for all $\beta \in [0, 1)$. Moreover A_{tr} has the following structure*

$$A_{tr} = \Pi_{\bar{\Omega}} K^+(\tilde{\Omega})$$

where $\tilde{\Omega} = \mathbb{R}_+^N$, $K^+(\tilde{\Omega})$ is the set of all bounded nonnegative solutions of (2.19) in $\tilde{\Omega} = \mathbb{R}_+^N$, and $\Pi_{\bar{\Omega}}$ denotes the restriction to $\bar{\Omega}$. Hence,

$$A_{tr} \subset \{x \mapsto 0, x \mapsto V(x_N)\}$$

and $A_{tr} = \{x \mapsto V(x_N)\}$ if $f(0) > 0$, where V is the unique solution of (1.4). Lastly, for any bounded nonnegative solution u of (1.1) in Ω , the functions $T_h u$ converge as $h \rightarrow +\infty$ in $C_{loc}^{2,\beta}(\bar{\Omega})$ for all $\beta \in [0, 1)$ either to 0 or to $x \mapsto V(x_N)$, and they do converge to the function $x \mapsto V(x_N)$ if $f(0) > 0$.

The next theorem deals with the analysis of the asymptotic behavior as $x_1 \rightarrow +\infty$ of the nonnegative bounded classical solutions $u \in K^+$ of equation (1.11) in the half-space $\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}$. For any $M \geq 0$, we define

$$K_M^+ = K^+ \cap \{0 \leq u \leq M\}.$$

Theorem 4.4 *Let N be any integer such that $N \geq 2$ and assume that, in addition to (1.3), the given locally Lipschitz-continuous function f from \mathbb{R}_+ to \mathbb{R} satisfies (1.13). Then, for every $M \geq \mu$, the trajectory dynamical system (T_h, K_M^+) associated to (1.11) possesses a global attractor A_{tr} , which is bounded in $C_b^{2,\beta}(\bar{\Omega}')$ and then compact in $C_{loc}^{2,\beta}(\bar{\Omega}')$ for all $\beta \in [0, 1)$, and satisfies*

$$A_{tr} = \Pi_{\bar{\Omega}'} K_M^+(\tilde{\Omega}') \tag{4.33}$$

where $\tilde{\Omega}' = \mathbb{R}^N$, $K_M^+(\tilde{\Omega}') = K^+(\tilde{\Omega}') \cap \{0 \leq u \leq M\}$, $K^+(\tilde{\Omega}')$ is the set of all bounded nonnegative solutions of (2.18) in $\tilde{\Omega}' = \mathbb{R}^N$, and $\Pi_{\bar{\Omega}'}$ denotes the restriction to $\bar{\Omega}'$. Hence,

$$A_{tr} = \{z \in [0, M] \mid f(z) = 0\} = E \cap [0, M].$$

Lastly, for any bounded nonnegative solution u of (1.11) in Ω' such that $0 \leq u \leq M$, the functions $T_h u$ converge as $h \rightarrow +\infty$ in $C_{loc}^{2,\beta}(\overline{\Omega'})$ for all $\beta \in [0, 1)$ to some $z \in E \cap [0, M]$ which is uniquely defined by u .

The next two theorems, which are concerned with the case of functions f fulfilling the condition (1.7), are based on the new Liouville type Theorems 2.2 and 2.3 which were already stated in Section 2.

Theorem 4.5 *Let N be any integer such that $N \geq 2$, let f be any locally Lipschitz-continuous function from \mathbb{R}_+ to \mathbb{R} satisfying (1.7) and assume that, for problem (1.1) in the quarter-space Ω , the set K^+ is not empty. Then, for every sufficiently large $M \geq 0$, the trajectory dynamical system (T_h, K_M^+) associated to (1.1) possesses a global attractor A_{tr} , which is bounded in $C_b^{2,\beta}(\overline{\Omega})$ and then compact in $C_{loc}^{2,\beta}(\overline{\Omega})$ for all $\beta \in [0, 1)$ and has the following structure*

$$A_{tr} = \Pi_{\overline{\Omega}} K_M^+(\tilde{\Omega}) \quad (4.34)$$

where $K_M^+(\tilde{\Omega}) = K^+(\tilde{\Omega}) \cap \{0 \leq u \leq M\}$. Hence,

$$A_{tr} = \{x \mapsto V_z(x_N) \mid z \in [0, M], f(z) = 0\}.$$

Lastly, for any bounded nonnegative solution u of (1.1) in Ω such that $0 \leq u \leq M$, the functions $T_h u$ converge as $h \rightarrow +\infty$ in $C_{loc}^{2,\beta}(\overline{\Omega})$ for all $\beta \in [0, 1)$ to some function $x \mapsto V_z(x_N)$, where $z \in E \cap [0, M]$ is uniquely defined by u and E denotes the set of zeroes of the function f .

Analogously to Theorem 4.4 we have the following Theorem 4.6 in the case of the half-space $\Omega' = (0, +\infty) \times \mathbb{R}^{N-1}$.

Theorem 4.6 *Let N be any integer such that $N \geq 2$, let f be any locally Lipschitz-continuous function from \mathbb{R}_+ to \mathbb{R} satisfying (1.7), and assume that, for problem (1.11) in the half-space Ω' , the set K^+ is not empty. Then, for every sufficiently large $M \geq 0$, the trajectory dynamical system (T_h, K_M^+) associated to (1.11) possesses a global attractor A_{tr} , which is bounded in $C_b^{2,\beta}(\overline{\Omega'})$ and then compact in $C_{loc}^{2,\beta}(\overline{\Omega'})$ for all $\beta \in [0, 1)$, and satisfies (4.33). Lastly, for any bounded nonnegative solution u of (1.11) in Ω' such that $0 \leq u \leq M$, the functions $T_h u$ converge as $h \rightarrow +\infty$ in $C_{loc}^{2,\beta}(\overline{\Omega'})$ for all $\beta \in [0, 1)$ to some $z \in E \cap [0, M]$ which is uniquely defined by u .*

In what follows we prove Theorem 4.3. A proof of Theorems 4.4-4.6 can be done in the same manner as in Theorem 4.3 with some minor modifications (see Remark 4.8).

Proof of Theorem 4.3. Let K^+ be the set of all bounded nonnegative solutions of (1.1) in Ω . Due to the assumptions (1.3), the set K^+ is not empty (the function $x \mapsto V(x_N)$, where V is the unique solution of (1.4), belongs to K^+) and due to the translation invariance of (1.1), it follows that $T_h : K^+ \rightarrow K^+$ is well defined for all $h \geq 0$, where $(T_h u)(x_1, x', x_N) := u(x_1 + h, x', x_N)$.

To show that (T_h, K^+) possesses a global attractor, it suffices to show (see [2] and the references therein) that

- for any fixed $h > 0$, T_h is a continuous map in K^+ (we recall that K^+ is endowed with local topology according to the embedding of K^+ in $C_{loc}^{2,\beta}(\overline{\Omega} \setminus \{x_1 = 0\})$) for all $\beta \in [0, 1)$;
- the semi-flow $(T_h)_{h \geq 0}$ possesses a compact attracting (absorbing) set in $C_{loc}^{2,\beta}(\overline{\Omega})$, which is even bounded in $C_b^{2,\beta}(\overline{\Omega})$, for all $\beta \in [0, 1)$.

Note that, the continuity of T_h in $C_{loc}^{2,\beta}(\overline{\Omega} \setminus \{x_1 = 0\})$ is obvious, because the shift operator is continuous in this topology, as well as its restriction to K^+ . As for the existence of compact attracting (absorbing) set for the semi-flow $(T_h)_{h \geq 0}$, it follows from the fact that the set of all bounded nonnegative solutions of (2.19) in $\cup_{h \geq 0} T_{-h}(\overline{\Omega}) = \tilde{\Omega} = \mathbb{R}_+^N$ under the assumption (1.3) on f is uniformly bounded. Indeed, as already recalled in Section 2.1 and according to a result of [5], under the assumption (1.3), any bounded solution of (2.19) in \mathbb{R}_+^N which is positive in $\mathbb{R}^{N-1} \times (0, +\infty)$ has one-dimensional symmetry, that is

$$u(x_1, x', x_N) = V(x_N)$$

where $0 \leq V < \mu$ is the unique solution of (1.4). On the other hand, since $f(0) \geq 0$, any bounded nonnegative solution of (2.19) is either positive in $\mathbb{R}^{N-1} \times (0, +\infty)$, or identically 0 in \mathbb{R}_+^N , and it cannot be 0 if $f(0) > 0$. Thus, the set $K^+(\tilde{\Omega})$ of all bounded nonnegative solutions of (2.19) in $\tilde{\Omega} = \mathbb{R}_+^N$ is bounded in $L^\infty(\tilde{\Omega})$, namely

$$\sup_{u \in K^+(\tilde{\Omega})} \|u\|_\infty \leq \mu.$$

Then the existence of a compact absorbing set for $(T_h)_{h \geq 0}$ in $C_{loc}^{2,\beta}(\overline{\Omega})$, which is even bounded in $C_b^{2,\beta}(\overline{\Omega})$, for all $\beta \in [0, 1)$ is a consequence of the uniform boundedness of all solutions of (2.19) in $\tilde{\Omega} = \mathbb{R}_+^N$ and of standard elliptic estimates. Hence the semigroup (T_h, K^+) possesses a global attractor A_{tr} in K^+ which is bounded in $C_b^{2,\beta}(\overline{\Omega})$ and compact in $C_{loc}^{2,\beta}(\overline{\Omega})$ for all $\beta \in [0, 1)$.

To prove the convergence part of Theorem 4.3, as we will see below, it is sufficient to show that $A_{tr} = \Pi_{\overline{\Omega}} K^+(\tilde{\Omega})$. Assuming for a moment that this representation is true, we obtain from the previous considerations that

$$A_{tr} \subset \{x \mapsto 0, x \mapsto V(x_N)\}, \quad (4.35)$$

and A_{tr} is then equal to the singleton $\{x \mapsto V(x_N)\}$ if $f(0) > 0$. Hence, for any bounded nonnegative solution u of (1.1), since $\{T_h u, h \geq 1\}$ is bounded in $C_b^{2,\beta}(\overline{\Omega})$ and compact in $C_{loc}^{2,\beta}(\overline{\Omega})$ for all $\beta \in [0, 1)$, the ω -limit set $\omega(u)$ of u is not empty and it is an invariant and connected subset of A_{tr} . Since A_{tr} is totally disconnected,² it follows that either $\omega(u) = \{x \mapsto 0\}$ or $\omega(u) = \{x \mapsto V(x_N)\}$, the latter being necessarily true if $f(0) > 0$.

To complete the proof of Theorem 4.3 it remains to show that $A_{tr} = \Pi_{\overline{\Omega}} K^+(\tilde{\Omega})$. First we prove that $\Pi_{\overline{\Omega}} \hat{u} \in A_{tr}$ for any bounded nonnegative solution \hat{u} of (2.19) in $\tilde{\Omega} = \mathbb{R}_+^N$,

²This property is obvious here due to (4.35). See Remark 4.8 for a comment about the other situations, corresponding to Theorems 4.4, 4.5 and 4.6.

that is $\hat{u} \in K^+(\tilde{\Omega})$. Indeed, for such a \hat{u} , the family $(\Pi_{\tilde{\Omega}}(T_{-h}\hat{u}))_{h \geq 0}$ is uniformly bounded in $C_b^{2,\beta}(\tilde{\Omega})$ and compact in $C_{loc}^{2,\beta}(\tilde{\Omega})$ for all $\beta \in [0, 1)$ and, according to definition of the attractor, there holds

$$T_h \Pi_{\tilde{\Omega}}(T_{-h}\hat{u}) \longrightarrow A_{tr} \text{ in } C_{loc}^{2,\beta}(\tilde{\Omega}) \text{ as } h \rightarrow +\infty,$$

for all $\beta \in [0, 1)$. On the other hand, $T_h \Pi_{\tilde{\Omega}}(T_{-h}\hat{u}) = \Pi_{\tilde{\Omega}}\hat{u}$. Hence $\Pi_{\tilde{\Omega}}\hat{u} \in A_{tr}$.

Next we prove the reverse inclusion. To this end, let us recall that (T_h, K^+) possesses an absorbing set which is bounded in $C_b^{2,\beta}(\tilde{\Omega})$ and then compact in $C_{loc}^{2,\beta}(\tilde{\Omega})$ for all $\beta \in [0, 1)$, say $\mathbb{B}_* \subset K^+$, and, as a consequence,

$$A_{tr} = \omega(\mathbb{B}_*) = \bigcap_{h \geq 0} \left[\bigcup_{s \geq h} T_s \mathbb{B}_* \right],$$

where $[\]$ means the closure in $C_{loc}^{2,\beta}(\tilde{\Omega})$ (see [2, 9] and the references therein). Let now $u \in A_{tr}$. The property $A_{tr} = \omega(\mathbb{B}_*)$ implies that there exist an increasing sequence $(h_k)_{k \in \mathbb{N}} \rightarrow +\infty$ and a sequence of solutions $(u_k)_{k \in \mathbb{N}}$ in \mathbb{B}_* , such that

$$u = \lim_{k \rightarrow +\infty} T_{h_k} u_k \tag{4.36}$$

in $C_{loc}^{2,\beta}(\tilde{\Omega})$ for all $\beta \in [0, 1)$. Note that the solution $T_{h_k} u_k$ is defined not only in Ω , but also in the domain $(-h_k, +\infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+$, and that

$$\sup_{k \in \mathbb{N}} \|T_{h_k} u_k\|_{C_b^{2,\beta}([-h_k + \varepsilon, \infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+)} < +\infty \tag{4.37}$$

for all $\varepsilon > 0$ and $\beta \in [0, 1)$, from standard elliptic estimates. Consequently, for every $k_0 \in \mathbb{N}$ and $\beta \in [0, 1)$, the sequence $(T_{h_k} u_k)_{k > k_0}$ is precompact in $C_{loc}^{2,\beta}([-h_{k_0}, \infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+)$. Taking a subsequence, if necessary, and using Cantor's diagonal procedure and the fact $h_k \rightarrow \infty$, we can say that this sequence converges to $\hat{u} \in C_{loc}^{2,\beta}(\mathbb{R}_+^N)$ in the spaces $C_{loc}^{2,\beta}([-h_{k_0}, \infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+)$ for every $k_0 \in \mathbb{N}$ and for every $\beta \in [0, 1)$. Then (4.37) implies that $\hat{u} \in C_b^{2,\beta}(\mathbb{R}_+^N)$ for every $\beta \in [0, 1)$. Lastly, the functions $T_{h_k} u_k$ are nonnegative solutions of (1.1) in $(-h_k, +\infty) \times \mathbb{R}^{N-2} \times \mathbb{R}_+$ and by letting $k \rightarrow +\infty$, we easily obtain that \hat{u} is a bounded nonnegative solution of (2.19) in $\tilde{\Omega} = \mathbb{R}_+^N$. Finally, formula (4.36) implies that

$$\Pi_{\tilde{\Omega}}\hat{u} = u.$$

Thus $u \in \Pi_{\tilde{\Omega}}K^+(\tilde{\Omega})$ and the representation formula $A_{tr} = \Pi_{\tilde{\Omega}}K^+(\tilde{\Omega})$ is proved. The proof of Theorem 4.3 is thereby complete. \square

Remark 4.7 As far as Theorems 1.1 and 1.2 on the one hand, and Theorems 4.3 and 4.5 on the other hand, are concerned, we especially emphasize that the DS approach simplified in a very elegant way most of the computations regarding the asymptotic behavior of the solutions of (1.1) as $x_1 \rightarrow +\infty$. However, the DS approach and Theorem 4.3 (resp. Theorem 4.5) do not provide as in PDE approach the fact that only the limiting profile $V(x_N)$ (resp. $V_z(x_N)$ for some $z \in E \setminus \{0\}$) is selected, even if $f(0) = 0$, as soon as $u_0 \not\equiv 0$

on $\{0\} \times \mathbb{R}^{N-2} \times (0, +\infty)$. In the PDE proof of Theorems 1.1 and 1.2, it is indeed shown that the condition $u_0 \not\equiv 0$ implies that u is separated from 0 for large enough x_1 and x_N . This property is not shown in the DS proof of Theorems 4.3 and 4.5. Similar comments also hold for Theorems 1.3, 1.4, 4.4 and 4.6, where the PDE proof provides the convergence to a *non-zero* zero of f , what the DS proof does not. Lastly, in some of the results obtained through the PDE approach, the convergence of the solutions as $x_1 \rightarrow +\infty$ is proved to be uniform with respect to the variables (x', x_N) , while the DS approach only provides local convergence, due to the necessity of using the local topology to get the existence of a global attractor.

Remark 4.8 In Theorem 4.5 (resp. Theorem 4.6) under assumption (1.7), the phase space K_M^+ , which is invariant under the semigroup T_h , is not empty for any sufficiently large M , because K^+ is assumed to be not empty (one can take M as any nonnegative real number such that $M \geq \|U\|_\infty$, where U is any fixed element in K^+). Then, in the same manner as in the proof of Theorem 4.3, using both the representation formula $A_{tr} = \Pi_{\overline{\Omega}} K_M^+(\overline{\Omega})$ (resp. $A_{tr} = \Pi_{\overline{\Omega'}} K_M^+(\mathbb{R}^N)$), the Liouville theorems of Section 2 and Lemma 4.1 one obtains the desired conclusions. In particular, for Theorem 4.5 (resp. Theorem 4.6) about problem (1.1) in Ω (resp. (1.11) in Ω'), the total disconnectedness of A_{tr} follows from the representation formula (4.34) (resp. (4.33)), from Theorem 2.3 (resp. Theorem 2.2) and from Lemma 4.1 with, here, $Z_f = E$ (resp. condition (1.7) again). Note that for Theorem 4.4 in the case of assumptions (1.3) and (1.13), the total disconnectedness of A_{tr} follows from Theorem 2.7 and assumption (1.13) again.

Remark 4.9 Note that neither the concrete choice of the domain Ω (or Ω') nor the concrete choice of the “time” direction x_1 are essential for the use of the trajectory dynamical system’ approach. Indeed, let us replace the “time” direction x_1 by any fixed direction $\vec{l} \in \mathbb{R}^N$ and correspondingly $T_h^{\vec{l}} u = u(\cdot + h\vec{l})$ for $h \in \mathbb{R}_+$ and $u \in K^+$. Then the above construction seems to be applicable if the domain Ω satisfies the following assumptions:

- $T_h \Omega \subset \Omega$ (this is necessary in order to define the restriction of T_h to the trajectory phase space K^+ or K_M^+).
- $\cup_{h \geq 0} T_{-h} \overline{\Omega} = \mathbb{R}_+^N$, or \mathbb{R}^N (this is required in order to obtain representation formulas of the type $A_{tr} = \Pi_{\overline{\Omega}} K^+(\mathbb{R}_+^N)$ or $A_{tr} = \Pi_{\overline{\Omega}} K^+(\mathbb{R}^N)$, with possibly K_M^+ instead of K^+).

References

- [1] S.B. Angenent, *Uniqueness of the solution of a semilinear boundary value problem*, Math. Ann. **272** (1985), 129-138.
- [2] A. Babin, *Attractor of the generalized semigroup generated by an elliptic equation in a cylindrical domain*, Russian Acad. Sci. Izv. Math. **44** (1995), 207-233.
- [3] J. Ball, *Continuity properties and attractors of generalized semiflows and the Navier-Stokes equations*, Nonlinear Science **4** (1997), 475-502.

- [4] H. Berestycki, L. Caffarelli, L. Nirenberg, *Symmetry for elliptic equations in a half space*, In: Boundary Value Problems for Partial Differential Equations and Applications, J.-L. Lions et al. (eds.), RMA Res. Notes Appl. Math., Masson, Paris, 1993, 27-42.
- [5] H. Berestycki, L. Caffarelli, L. Nirenberg, *Monotonicity for elliptic equations in unbounded Lipschitz domains*, Comm. Pure Appl. Math. **50** (1997), 1089-1111.
- [6] H. Berestycki, L. Caffarelli, L. Nirenberg, *Further qualitative properties for elliptic equations in unbounded domains*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **5** (1997), 69-94.
- [7] H. Berestycki, F. Hamel, H. Matano, *Bistable travelling waves around an obstacle*, Comm. Pure Appl. Math. **62** (2009), 729-788.
- [8] H. Berestycki, L. Nirenberg, *On the method of moving planes and the sliding method*, Bol. Soc. Bras. Mat. **22** (1991), 1-37.
- [9] V. Chepyzhov, M. Vishik, *Attractors for equations of mathematical physics*, American Mathematical Society, Providence, RI, 2002.
- [10] Ph. Clément, G. Sweers, *Existence and multiplicity results for a semilinear eigenvalue problem*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **14** (1987), 97-121.
- [11] E.N. Dancer, *Some notes on the method of moving planes*, Bull. Aust. Math. Soc. **46** (1992), 425-434.
- [12] L. Dupaigne, A. Farina, *Liouville Theorems for stable solutions of semilinear elliptic equations with convex nonlinearities*, Nonlinear Anal. **70** (2009), 2882-2888.
- [13] M.A. Efendiev, H. Ibrahim, R. Monneau, *Different profiles for a semi-linear elliptic equation in unbounded domain*, preprint.
- [14] A. Farina, *On the classification of solutions of the Lane-Emden equation on unbounded domains of \mathbb{R}^N* , J. Math. Pures Appl. **87** (2007), 537-561.
- [15] A. Farina, B. Sciunzi, E. Valdinoci, *Bernstein and De Giorgi type problems: new results via a geometric approach*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (5) **7** (2008), 741-791.
- [16] A. Farina, E. Valdinoci, *Flattening results for elliptic PDEs in unbounded domains with applications to overdetermined problems*, Arch. Ration. Mech. Anal. **195** (2010), 1025-1058.
- [17] B. Gidas, W.N. Ni, L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), 209-243.
- [18] G. Sell, *Global attractors for the three dimensional Navier-Stokes equations*, J. Dynam. Diff. Equations **8** (1996), 1-33.