

Geometrical web models

Jesper L. Jacobsen ^{1,2,3}

¹Ecole Normale Supérieure, Paris (Laboratoire de Physique)

²Sorbonne Université, Paris (Faculté des Sciences et Ingénierie)

³Commissariat à l'Energie Atomique, Saclay (Institut de Physique Théorique)

Les Roches Rouges, Agay

Probability and Conformal Field Theory

21 September 2022


Collaborators: Augustin Lafay, Azat Gainutdinov

Loop models: what, why, how?

- Self-avoiding (open or closed) simple curves in two dimensions
- Polymers, level lines, domain walls, electron gases
- Lattice: Integrability, knot theory, cellular algebras, category theory
- Continuum limit: CFT, CLE, SLE

Definition and features

- Fix lattice of nodes and links
- Place bonds on some links so as to form set of loops
- Weight x per bond (+ maybe further local weights) and N per loop
- For $|N| \leq 2$, dense and dilute critical points x_c^\pm
- Continuum limit of compactified free bosonic field (Coulomb gas)
[Nienhuis, Di Francesco-Saleur-Zuber, Duplantier, Cardy . . .]







Generalisation to webs

- Allow for branchings and bifurcations (with weights)
- Topological rules give weight to each connected web component
- Properties and possible critical behaviour?

Motivations for webs

- Domain walls in spin systems [[Dubail-JJ-Saleur](#), [Picco-Santachiara](#)]
- Network models for topological phases [[Kitaev](#), [Levin-Wen](#), [Fendley](#)]
- Spiders in invariance theory [[Kuperberg](#), [Kim](#), [Cautis-Kamnitzer-Morrison](#)]

Thin and thick domain walls (Q = 3 Potts model)




Questions (physics)

- How to define a “good” model of webs on the lattice?
- Fractal dimension of such domain walls (bulk / boundary)?
- Fractal dimension of an entire web component?
- Topological weight of web versus chromatic polynomial in $Q = 3$?
- Web model away from this special point?

Questions (mathematics)

- Algebraic construction accounting for bifurcations?
- Loop model has $U_{-q}(\mathfrak{sl}_2)$ symmetry, can we get $U_{-q}(\mathfrak{sl}_n)$?




Web model from Kuperberg A_2 spider ($U_{-q}(\mathfrak{sl}_3)$ case)

Lattice considerations

- Hexagonal (honeycomb) lattice \mathbb{H} with nodes and links
- Configuration c by drawing bonds on some links, with constraints:
 - Nodes have valence 0, 2 or 3: closed web with 3-valent vertices
 - Each bond is oriented. Orientations conserved at 2-valent nodes
 - Vertices are sources or sinks (all bonds point in or out)

Each configuration can be seen as an abstract graph (vertices/edges). It is **closed**, **planar**, **trivalent**, **bipartite**. Fix an orientation (= 'up').



Rules for 'reducing' a configuration [Kuperberg]

$$\text{circle with arrow} = [3]_q \quad (1)$$

$$\text{vertical line with two loops} = [2]_q \text{ vertical line} \quad (2)$$

$$\text{two vertical lines with crossings} = \text{two vertical lines} + \text{two arcs} \quad (3)$$

- Rotated and arrow-reversed diagrams not shown.
- A web component always has ≥ 1 polygon of degree 0, 2 or 4.
- The three rules thus evaluate any web to a number (its weight)

Define q -deformed numbers: $[k]_q = \frac{q^k - q^{-k}}{q - q^{-1}}$

Defining the web model

- Sum over configurations $c \in K$ on \mathbb{H}
- Local weights: x_1 (up bond), x_2 (down bond), y (sink), z (source)
- Partition function:

$$Z_K = \sum_{c \in K} x_1^{N_1} x_2^{N_2} (yz)^{N_V} w_K(c)$$

with N_1 up-bonds, N_2 down-bonds, and N_V vertex pairs

Definition

- Spins $\sigma_i \in \mathbb{Z}_3 := \{0, 1, 2\}$ defined on triangular lattice $\mathbb{T} = \mathbb{H}^*$.
- Weight of link $(ij) \in \mathbb{T}$ defined as $x_{\sigma_j - \sigma_i}$, with j to the right of i .
- Normalise $x_0 = 1$. Weight x_1 or x_2 for a piece of domain wall.

Note: vertex is a sink (source) if spins follow cyclically $0 \rightarrow 1 \rightarrow 2 \rightarrow 0$ upon turning anticlockwise (clockwise).

Partition function

$$Z_{\text{spin}} = 3 \sum_{c \in K} x_1^{N_1} x_2^{N_2}$$

- Equivalent to web model if $w'_K(c) := (yz)^{N_V} w_K(c) = 1$ for any c .

Equivalence at a special point:

$$q = e^{j\frac{\pi}{4}},$$

$$yz = 2^{-\frac{1}{2}}.$$


Proof: Absorb y and z into the vertices. Use $[3]_q = 1$ and $[2]_q = \sqrt{2}$. Then the rules become probabilistic:

$$\begin{array}{l}
 \text{Circle with arrow} = 1 \\
 \text{Line with loop} = \text{Line} \\
 \text{Line with crossings} = \frac{1}{2} \text{Parallel lines} + \frac{1}{2} \text{Crossing}
 \end{array}$$

Generalisation to $U_{-q}(\mathfrak{sl}_n)$ symmetry


Based on spider defined by [Cautis-Kamnitzer-Morrison]

- Webs are still closed, oriented, planar, trivalent graphs. But not always bipartite as before.
- Edges carry an integer flow $i \in \llbracket 1, n-1 \rrbracket$.
- Generators conserve flow, or change by n due to ‘tags’:



- Flow labels fundamental representations of $U_{-q}(\mathfrak{sl}_n)$. Orientation distinguishes between dual or not.

Rules (mirrored and the arrow-reversed versions omitted):



$$\begin{array}{c} k \\ \downarrow \\ k-1 \\ \downarrow \\ k \end{array} \begin{array}{c} \leftarrow 1 \\ \downarrow \\ \leftarrow 1 \\ \downarrow \\ \leftarrow 1 \end{array} \begin{array}{c} l \\ \downarrow \\ l+1 \\ \downarrow \\ l \end{array} = \begin{array}{c} k \\ \downarrow \\ k+1 \\ \downarrow \\ k \end{array} \begin{array}{c} \leftarrow 1 \\ \downarrow \\ \leftarrow 1 \\ \downarrow \\ \leftarrow 1 \end{array} \begin{array}{c} l \\ \downarrow \\ l-1 \\ \downarrow \\ l \end{array} + [k-l]_q \begin{array}{c} \downarrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \downarrow \end{array}$$

$$\begin{array}{c} k \\ \downarrow \\ n-k \\ \downarrow \\ k \end{array} = \begin{array}{c} \downarrow \\ \downarrow \end{array}$$

$$\begin{array}{c} n-k-l \\ \downarrow \\ k+l \\ \swarrow \quad \searrow \\ k \quad l \end{array} = \begin{array}{c} n-k-l \\ \downarrow \\ n-l \\ \swarrow \quad \searrow \\ k \quad l \end{array}$$

$$\begin{array}{c} n-l \\ \downarrow \\ l \\ \swarrow \quad \searrow \\ k+l \quad k \end{array} = \begin{array}{c} n-l \\ \downarrow \\ n-k-l \\ \swarrow \quad \searrow \\ k+l \quad k \end{array}$$

$$\begin{array}{c} n-k \\ \downarrow \\ \downarrow \\ \downarrow \\ k \end{array} = (-1)^{k(n-k)} \begin{array}{c} n-k \\ \downarrow \\ \downarrow \\ \downarrow \\ k \end{array}$$

Short summary of results

- Case $n = 3$ gives back the Kuperberg web model.
- Case $n = 2$ gives the well-known Nienhuis loop model.
- Special point $q = e^{j\frac{\pi}{n+1}}$ equivalent to \mathbb{Z}_n spin model.

Outlook this far

- \mathbb{Z}_n spin models known to be critical (with appropriate weights) [Fateev-Zamolodchikov]
- Therefore expect the special point to be critical for any n .
- Web models likely have larger critical manifold (vary q and x, y, z).


To investigate criticality we wish a local formulation

- Analogous to vertex models for Potts and $O(N)$ models.
- The locality enables us to define a transfer matrix.
 - Good for numerical study and makes contact with integrability.
 - Non-local TM also possible for loops, but seems difficult for webs.
- Vertex model defines equivalent ($n - 1$ component) height model.
 - Starting point for Coulomb gas construction and CFT identification.

Local reformulation for $U_{-q}(\mathfrak{sl}_3)$ web model

Basic idea

- Decorate bonds by extra degrees of freedom ($n = 3$ colours).
- They allow to redistribute the web weight locally.
- Summing over colours gives back the undecorated model.
- Each link can now be in 7 different states.



Reminder for $n = 2$ loop case

- Write $N = q + q^{-1} = [2]_q$.
- Orient each loop in two ways (clockwise, anticlockwise).
- Give $q^{-\frac{\theta}{2\pi}}$ to a left-turn through angle θ .


$\begin{array}{c} \uparrow \\ | \\ \swarrow \nearrow \\ \nearrow \swarrow \\ \searrow \end{array} = xq^{-\frac{1}{6}}, \quad \begin{array}{c} \downarrow \\ | \\ \swarrow \nearrow \\ \nearrow \swarrow \\ \searrow \end{array} = xq^{\frac{1}{6}}, \quad \begin{array}{c} \text{---} \\ | \\ \swarrow \nearrow \\ \nearrow \swarrow \\ \text{---} \end{array} = 1$

Remark

Better to think of these two 'orientations' as colourings. The analogue for $n = 3$ is the three colours. The orientations distinguish (for $n \geq 3$) fundamental and dual fundamental, but for $n = 2$ the two coincide!


Basic idea for $n = 3$

- Three colours **RGB**.
- Weight $q^2 + 1 + q^{-2} = [3]_q$ for sum over (say) clockwise loop. Opposite phases for an anticlockwise loop (same sum). Set $x_1 = x_2$ for convenience.




A red vertex with three edges. The top edge is solid with an upward arrow. The bottom-left edge is solid with a leftward arrow. The bottom-right edge is dashed with a rightward arrow. This configuration represents a clockwise loop.

$$= xq^{-\frac{1}{3}},$$




A blue vertex with three edges. The top edge is solid with an upward arrow. The bottom-left edge is solid with a leftward arrow. The bottom-right edge is dashed with a rightward arrow. This configuration represents a clockwise loop.

$$= x,$$




A green vertex with three edges. The top edge is solid with an upward arrow. The bottom-left edge is solid with a leftward arrow. The bottom-right edge is dashed with a rightward arrow. This configuration represents a clockwise loop.

$$= xq^{\frac{1}{3}}$$




A red vertex with three edges. The top edge is solid with an upward arrow. The bottom-left edge is dashed with a leftward arrow. The bottom-right edge is solid with a rightward arrow. This configuration represents an anticlockwise loop.

$$= xq^{\frac{1}{3}},$$



A blue vertex with three edges. The top edge is solid with an upward arrow. The bottom-left edge is dashed with a leftward arrow. The bottom-right edge is solid with a rightward arrow. This configuration represents an anticlockwise loop.


$$= x,$$





A green vertex with three edges. The top edge is solid with an upward arrow. The bottom-left edge is dashed with a leftward arrow. The bottom-right edge is solid with a rightward arrow. This configuration represents an anticlockwise loop.


$$= xq^{-\frac{1}{3}}$$

The 'tricky' part involving vertices


$$= zx^{\frac{3}{2}}q^{-\frac{1}{6}},$$


$$= zx^{\frac{3}{2}}q^{\frac{1}{6}}$$


$$= yx^{\frac{3}{2}}q^{\frac{1}{6}},$$


$$= yx^{\frac{3}{2}}q^{-\frac{1}{6}}$$

Proof for the 'digon' rule (2)


$$\text{Digon (blue/green)} + \text{Digon (green/blue)} = q^{\frac{1}{6} \times 2 + \frac{2}{3}} + q^{-\frac{1}{6} \times 2 - \frac{2}{3}} = [2]_q$$

Proof for the 'square' rule (3)

$$\text{Square (blue/red)} + \text{Square (red/blue)} = \text{Two green arcs} + \text{Green arc with crossing}$$

Other colours / arrangements of external legs work similarly.

Defining the transfer matrix



Built of pieces $t_{(1)} : \mathcal{H} \otimes \mathcal{H} \rightarrow \mathcal{H}$ and $t_{(2)} : \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}$, so that

$$T = \left(\prod_{k=0}^{L-1} t_{2k+1} \right) \left(\prod_{k=1}^{L-1} t_{2k} \right)$$

with $t = t_{(2)}t_{(1)}$. Write t_i , with i specifying the position.

Technically T is an intertwiner of the quantum group action.


- Let $\{v_1, v_2, v_3\}$ be a basis of the first fundamental V_1 of $U_{-q}(sl_3)$.
- Let $\{w_1, w_2, w_3\}$ be a basis of the dual V_1^* , so that $w_i(v_j) = \delta_{ij}$.
- Relate $\{v_1, v_2, v_3, w_1, w_2, w_3, 1\}$ to the basis $\{|\uparrow\rangle, |\uparrow\rangle, |\uparrow\rangle, |\downarrow\rangle, |\downarrow\rangle, |\downarrow\rangle, | \rangle\}$ of coloured arrows.
Amounts to drawing each link vertically and providing the corresponding powers of q .
- Draw the diagrams of all transitions in $t_{(1)}$ and $t_{(2)}$. For instance:

$$\begin{aligned}
 t_{(1)} = & z x_1 x_2^{\frac{1}{2}} \begin{array}{c} \downarrow \\ \swarrow \quad \searrow \end{array} + y x_1^{\frac{1}{2}} x_2 \begin{array}{c} \uparrow \\ \swarrow \quad \searrow \end{array} + x_1 \begin{array}{c} \uparrow \\ \swarrow \quad \text{---} \end{array} + x_1 \begin{array}{c} \uparrow \\ \text{---} \quad \searrow \end{array} + x_2 \begin{array}{c} \downarrow \\ \swarrow \quad \text{---} \end{array} \\
 & + x_2 \begin{array}{c} \downarrow \\ \text{---} \quad \searrow \end{array} + x_1^{\frac{1}{2}} x_2^{\frac{1}{2}} \begin{array}{c} \text{---} \\ \swarrow \quad \searrow \end{array} + x_1^{\frac{1}{2}} x_2^{\frac{1}{2}} \begin{array}{c} \text{---} \\ \swarrow \quad \searrow \end{array} + \begin{array}{c} \text{---} \\ \text{---} \end{array}
 \end{aligned}$$

- Let us have a look at just the first term!




- Express each diagram in terms of the elementary blocks (maps)



Their expressions follow from quantum group considerations.


- The first term is the composition of $coev$ and w :





- In the bases $\{|\uparrow\uparrow\rangle, |\uparrow\downarrow\rangle, |\uparrow\downarrow\rangle, |\uparrow\uparrow\rangle, |\uparrow\downarrow\rangle, |\uparrow\downarrow\rangle, |\uparrow\uparrow\rangle, |\uparrow\downarrow\rangle, |\uparrow\uparrow\rangle\}$ of $V_1 \otimes V_1$ and $\{|\downarrow\rangle, |\downarrow\rangle, |\downarrow\rangle\}$ of V_1^* , we finally get


$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & q^{\frac{1}{6}} & 0 & q^{-\frac{1}{6}} & 0 \\ 0 & 0 & q^{\frac{1}{6}} & 0 & 0 & 0 & q^{\frac{1}{6}} & 0 & 0 \\ 0 & q^{\frac{1}{6}} & 0 & q^{-\frac{1}{6}} & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

- Looks familiar?
- Hint:


 $= zx^{\frac{3}{2}} q^{-\frac{1}{6}},$


 $= zx^{\frac{3}{2}} q^{\frac{1}{6}}$


 $= yx^{\frac{3}{2}} q^{\frac{1}{6}},$



 $= yx^{\frac{3}{2}} q^{-\frac{1}{6}}$

Summary of this technical part

- The diagrams are intertwiners of $U_{-q}(\mathfrak{sl}_3)$.
- We can compute all elements of T in this way.
- We are now ready to diagonalise T numerically.

Phase diagram of the web model

- More efficient to use the geometry




- Connection to the (effective) central charge of CFT:

$$f_L = -\frac{2}{\sqrt{3}L} \log(\Lambda_{\max}),$$


$$f_L = f_\infty - \frac{\pi c_{\text{eff}}}{6L^2} + o\left(\frac{1}{L^2}\right).$$

c_{eff} for $q = e^{i\pi/5}$ in the (\sqrt{x}, y) plane



- Based on sizes $L = 5$ and $L = 6$.
- Coulomb gas prediction: dilute $c = \frac{4}{5}$ and dense $c = \frac{6}{5}$ phases.

Zoom of the interesting region



Coulomb gas predictions

Set $q = e^{i\gamma}$ with $\gamma \in [0, \pi]$.

CG of two bosons compactified on the root lattice of $s/3$

Coupling constant $g = 1 \pm \frac{\gamma}{\pi}$ in dilute (+) or dense (-) phase.

Central charge $c = 2 - 24 \frac{(g-1)^2}{g}$.

Example I: $\gamma = \frac{\pi}{5}$ as in numerical figures

Coupling constant $g = \frac{6}{5}$ (dilute) or $g = \frac{4}{5}$ (dense).

Central charge $c = \frac{6}{5}$ (dilute) or $c = \frac{4}{5}$ (dense).

Example II: $\gamma = \frac{\pi}{4}$ as at special point

Coupling constant $g = \frac{5}{4}$ (dilute) or $g = \frac{3}{4}$ (dense).

Central charge $c = \frac{4}{5}$ (dilute) or $c = 0$ (dense).

Corresponds to $Q = 3$ Potts model at $T = T_c$ or $T = \infty$.

What about integrability?

- The $n = 2$ model (Nienhuis loops) is integrable in both the dilute and dense phases [Baxter 1986-87]
- What about $n = 3$?
 - In the fully-packed case (a bond on every link = all fundamental reps) it is integrable [Reshethikhin]
 - We need now 7-dimensional reps (fundamental + dual + trivial)
- There are other rank-2 spiders (G_2 and B_2) related to 3-state Potts interfaces
 - Exact mappings (to appear)
 - In the G_2 case, we can relate to an integrable model coming from $U_q(D_4^{(3)})$ (in progress)

Summary

- Web models generalise the $U_{-q}(\mathfrak{sl}_2)$ loop model to $U_{-q}(\mathfrak{sl}_n)$.
- Geometrical content with applications to \mathbb{Z}_n spin interfaces.
- Dense and dilute critical points for $q = e^{i\gamma}$ and $\gamma \in [0, \pi]$.

In the pipeline

- Coulomb gas description and fractal dimension of defects
- Statistical models for other spiders
- Detailed representation theoretical study
- More relations to and input from integrable models

Further possibilities

- SLE-like description of branching curves?