

# Maximal regularity for non-autonomous Robin boundary conditions

Wolfgang Arendt<sup>\*</sup>

Sylvie Monniaux<sup>†‡</sup>

## Abstract

We consider a non-autonomous Cauchy problem

$$\dot{u}(t) + \mathcal{A}(t)u(t) = f(t), \quad u(0) = u_0$$

where  $\mathcal{A}(t)$  is associated with the form  $\mathfrak{a}(t; \cdot, \cdot) : V \times V \rightarrow \mathbb{C}$ , where  $V$  and  $H$  are Hilbert spaces such that  $V$  is continuously and densely embedded in  $H$ . We prove  $H$ -maximal regularity, i.e., the weak solution  $u$  is actually in  $H^1(0, T; H)$  (if  $u_0 \in V$  and  $f \in L^2(0, T; H)$ ) under a new regularity condition on the form  $a$  with respect to time; namely Hölder continuity with values in an interpolation space. This result is best suited to treat Robin boundary conditions. The maximal regularity allows one to use fixed point arguments to some non linear parabolic problems with Robin boundary conditions.

## 1 Introduction

In the background of this article is a longstanding problem by J.-L. Lions on non-autonomous forms. We give a solution of the problem in a special case which is most suitable for treating non-autonomous Robin boundary conditions. To be more specific we consider a non-autonomous form

$$\mathfrak{a} : [0, T] \times V \times V \rightarrow \mathbb{C}$$

where  $V$  is a Hilbert space continuously and densely embedded into another Hilbert space  $H$ . We assume that

$$|\mathfrak{a}(t; u, v)| \leq M \|u\|_V \|v\|_V \quad (t \in [0, T], u, v \in V) \quad (1.1)$$

$$\Re \mathfrak{a}(t; u, u) \geq \delta \|u\|_V^2 \quad (t \in [0, T], u \in V) \quad (1.2)$$

for some constants  $M, \delta > 0$ , and that  $\mathfrak{a}(\cdot; u, v)$  is measurable for all  $u, v \in V$ . Denote by  $\mathcal{A}(t) : V \rightarrow V'$  the operator given by

$$\langle \mathcal{A}(t)u, v \rangle = \mathfrak{a}(t; u, v), \quad v \in V.$$

The space

$$MR(V, V') := H^1(0, T; V') \cap L^2(0, T; V)$$

---

<sup>\*</sup>Universität Ulm, Germany

<sup>†</sup>Aix-Marseille Université, CNRS, Centrale Marseille, I2M, UMR 7373 – 13453 Marseille, France

<sup>‡</sup>partially supported by the ANR project HAB, ANR-12-BS01-0013-03, the Labex Archimède (ANR-11-LABX-0033) and the A\*MIDEX project (ANR-11-IDEX-0001-02)

is contained in  $\mathcal{C}([0, T]; H)$  and one has the following well-posedness result for weak solutions.

**Theorem** (Lions). *For all  $f \in L^2(0, T; V')$ ,  $u_0 \in H$ , there exists a unique  $u \in MR(V, V')$  solution of*

$$\dot{u}(t) + \mathcal{A}(t)u(t) = f(t), \quad u(0) = u_0$$

The letters  $MR$  are used to refer to “maximal regularity”; and indeed one has maximal regularity in  $V'$  in the sense that all three terms  $\dot{u}$ ,  $\mathcal{A}(\cdot)u(\cdot)$  and  $f$  occurring in the equation belong to  $L^2(0, T; V')$ . However, considering boundary valued problems one is interested in *strong solutions*, i.e., solutions  $u \in H^1(0, T; H)$  and not only in  $H^1(0, T; V')$  (note that  $H \hookrightarrow V'$  by the natural embedding).

**Problem.** Given  $f \in L^2(0, T; H)$ ,  $u_0 \in H$  good enough, under which regularity assumptions on the form  $\mathfrak{a}$  is  $u$  in  $H^1(0, T; H)$ ?

This problem is explicitly formulated by Lions [14, p. 98] if  $\mathfrak{a}(t; v, w) = \overline{\mathfrak{a}(t; w, v)}$  for all  $v, w \in V$ . In general, even for  $u_0 = 0$ , (1.1) and (1.2) are not sufficient for having  $u \in H^1(0, T; V)$ . This has been shown recently by Dier [9]. But several positive answers are given by Lions [14]. More recently it has been shown that actually  $u \in H^1(0, T; H)$  for any  $u_0 \in V$  provided  $\mathfrak{a}(\cdot; v, w)$  is Lipschitz continuous and symmetric (see [4] where also a multiplicative perturbation is admitted) or if  $\mathfrak{a}(\cdot; v, w)$  is symmetric and of bounded variations (see Dier [9]). Moreover, for  $u_0 = 0$ , one has  $u \in H^1(0, T; H)$  if  $\mathfrak{a}(\cdot; v, w)$  is Hölder continuous of order  $\alpha > \frac{1}{2}$  for all  $u, v \in V$ , see Ouhabaz-Spina [20]. This has been improved by Haak-Ouhabaz [10] where the authors remove the symmetry condition and allow non-zero initial conditions. The purpose of this article is to establish a different case. We consider  $0 < \gamma < 1$  and the complex interpolation space  $V_\gamma := [H, V]_\gamma$ . We assume that  $\mathfrak{a}$  satisfies (1.1), (1.2) and

$$|\mathfrak{a}(t, v, w) - \mathfrak{a}(s, v, w)| \leq c|t - s|^\alpha \|v\|_V \|w\|_{V_\gamma}$$

for all  $v, w \in V$ ,  $t, s \in [0, T]$ , where  $\alpha > \frac{\gamma}{2}$ . Then we show that the solution  $u$  from Lions' theorem is actually in  $H^1(0, T; H)$  whenever  $u_0 \in V$ . In other words, for all  $f \in L^2(0, T, H)$ ,  $u_0 \in V$  there is a unique

$$u \in MR_{\mathfrak{a}}(V, H) := \{u \in H^1(0, T; H) \cap L^2(0, T; V) : \mathcal{A}(\cdot)u(\cdot) \in L^2(0, T; H)\}$$

satisfying

$$\dot{u}(t) + \mathcal{A}(t)u(t) = f(t) \quad t - a.e.$$

Thus we have maximal regularity in  $H$  in the sense that all three terms  $\dot{u}$ ,  $\mathcal{A}(\cdot)u(\cdot)$  and  $f$  are in  $L^2(0, T; H)$ . Similar results can be found in a preprint by Ouhabaz [19] which was put on the arXiv website a few weeks after the present work.

Moreover, we show that  $MR_{\mathfrak{a}}(V, H) \subset \mathcal{C}([0, T]; V)$ . Our result can be applied to Robin boundary conditions. If  $\Omega$  is a bounded Lipschitz domain and

$$\mathcal{B} : [0, T] \rightarrow \mathcal{L}(L^2(\partial\Omega))$$

is Hölder continuous of order  $\alpha > \frac{1}{4}$  then given  $u_0 \in H^1(\Omega)$ ,  $f \in L^2(0, T; L^2(\Omega))$  there exists a unique  $u \in H^1(0, T; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega))$  such that  $\Delta u \in L^2(0, T; L^2(\Omega))$  and

$$\dot{u}(t) - \Delta u(t) = f(t)$$

$$\partial_\nu u(t) + B(t)u(t)|_{\partial\Omega} = 0$$

$$u(0) = u_0.$$

This in turn can be used to establish solutions of a non linear problem with non-autonomous boundary conditions, see Section 5.

## 2 Forms, interpolation and square root property

Throughout this paper we consider separable complex Hilbert spaces  $V$  and  $H$  with the property  $V \xhookrightarrow[d]{} H$ , i.e.,  $V$  is densely and continuously embedded in  $H$ . Then, as usual, we have

$$H \xhookrightarrow[d]{} V'$$

by associating to  $u \in H$  the linear mapping  $v \mapsto (v|u)$  where  $(\cdot|\cdot)$  is the scalar product in  $H$  and  $V'$  the antidual of  $V$ . For  $\ell \in [0, 1]$ , we denote by  $V_\ell$  the complex interpolation space  $[H, V]_\ell$ . Thus

$$V \hookrightarrow V_\ell \hookrightarrow H.$$

Moreover  $V_0 = H$ ,  $V_1 = V$ . Then  $V'_\ell := (V_\ell)' = [V', H]_{1-\ell}$ . In particular

$$H \hookrightarrow V'_\ell \hookrightarrow V'$$

and  $V'_0 = H$ ,  $V'_1 = V'$ .

Let  $\mathfrak{a} : V \times V \rightarrow \mathbb{C}$  be a sesquilinear form which is *continuous* (satisfying (1.1)) and *coercive* (satisfying (1.2)). Then  $\langle \mathcal{A}u, v \rangle := \mathfrak{a}(u, v)$  defines an invertible operator  $\mathcal{A} \in \mathcal{L}(V, V')$ . We denote by  $A$  the part of  $\mathcal{A}$  in  $H$ , i.e.,

$$D(A) := \{u \in V : \mathcal{A}u \in H\}, \quad Au := \mathcal{A}u.$$

The operators  $\mathcal{A}$  and  $A$  are sectorial. More precisely there exists a sector

$$\Sigma_\theta := \{re^{i\varphi} : r > 0, |\varphi| < \theta\}$$

with  $0 \leq \theta < \frac{\pi}{2}$  such that  $\sigma(A) \subset \Sigma_\theta$ ,  $\sigma(\mathcal{A}) \subset \Sigma_\theta$  and for all  $\ell \in [0, 1]$  and all  $\lambda \notin \Sigma_\theta$ ,

$$\|(\lambda \text{Id} - \mathcal{A})^{-1}\|_{\mathcal{L}(V'_\ell)} \leq \frac{c}{1 + |\lambda|}, \quad (2.1)$$

$$\|(\lambda \text{Id} - A)^{-1}\|_{\mathcal{L}(V_\ell)} \leq \frac{c}{1 + |\lambda|}, \quad (2.2)$$

$$\|(\lambda \text{Id} - \mathcal{A})^{-1}\|_{\mathcal{L}(V'_\ell, V)} \leq \frac{c}{(1 + |\lambda|)^{\frac{1-\ell}{2}}}, \quad (2.3)$$

$$\|(\lambda \text{Id} - \mathcal{A})^{-1}\|_{\mathcal{L}(V'_\ell, H)} \leq \frac{c}{(1 + |\lambda|)^{1-\frac{\ell}{2}}}, \quad (2.4)$$

$$\|(\lambda \text{Id} - A)^{-1}\|_{\mathcal{L}(H, V_\ell)} \leq \frac{c}{(1 + |\lambda|)^{1-\frac{\ell}{2}}}, \quad (2.5)$$

$$\|(\lambda \text{Id} - \mathcal{A})^{-1}\|_{\mathcal{L}(V'_\ell, V_\ell)} \leq \frac{c}{(1 + |\lambda|)^{1-\ell}}. \quad (2.6)$$

The angle  $\theta$  and the constant  $c$  merely depend on  $\delta$ ,  $M$ ,  $\ell$  and the embedding constant  $c_H$ ,

$$\|v\|_H \leq c_H \|v\|_V, \quad v \in V. \quad (2.7)$$

For the proof of the estimates above, we refer to Tanabe [22, Chapter 2], or Ouhabaz [18, Theorem 1.52 and Theorem 1.55] (see also Arendt, [2, Theorem 7.1.4 and Theorem 7.1.5]) We fix an angle  $\theta < \vartheta < \frac{\pi}{2}$  and denote by  $\Gamma$  the contour  $\Gamma := \{re^{\pm i\vartheta}, r \geq 0\}$  oriented downwards. The operator  $-A$  generates a holomorphic  $C_0$ -semigroup  $(e^{-tA})_{t \geq 0}$  on  $H$  given by

$$e^{-tA} = \frac{1}{2\pi i} \int_{\Gamma} e^{-\lambda t} (\lambda \text{Id} - A)^{-1} d\lambda, \quad (2.8)$$

and there exists a constant  $c > 0$  such that

$$\|tAe^{-tA}\|_{\mathcal{L}(H)} \leq c \quad \text{for all } t > 0. \quad (2.9)$$

Moreover a theorem by De Simon [8, Lemma 3.1] shows  $L^2$  maximal regularity in Hilbert spaces, i.e., for holomorphic semigroups on Hilbert spaces there exists a constant  $c > 0$  such that for all  $f \in L^2(0, \infty; H)$ ,

$$\begin{aligned} t \mapsto \int_0^t Ae^{(t-s)A} f(s) ds &\in L^2(0, \infty; H) \\ \text{and } \left\| t \mapsto \int_0^t Ae^{(t-s)A} f(s) \right\|_{L^2(0, \infty; H)} &\leq c \|f\|_{L^2(0, \infty; H)}. \end{aligned} \quad (2.10)$$

Moreover

$$\|e^{-tA}\|_{\mathcal{L}(H)} \leq ce^{-\varepsilon t}, \quad t \geq 0,$$

for some  $\varepsilon > 0, c > 0$ . Also, the operator  $-A$  generates an exponentially stable holomorphic  $C_0$ -semigroup  $(e^{-tA})_{t \geq 0}$  on  $V'$ . By  $\mathfrak{a}^*(u, v) := \mathfrak{a}(v, u)$  ( $u, v \in V$ ) we define the form  $\mathfrak{a}^*$  which is adjoint to  $\mathfrak{a}$ . Then the operator associated with  $\mathfrak{a}^*$  on  $H$  coincides with the adjoint  $A^*$  of  $A$ . We define the operator  $A^{-\frac{1}{2}} \in \mathcal{L}(H)$  by

$$A^{-\frac{1}{2}}u = \frac{1}{\sqrt{\pi}} \int_0^\infty t^{-\frac{1}{2}} e^{-tA} u dt, \quad (2.11)$$

and we let  $D(A^{\frac{1}{2}}) := A^{-\frac{1}{2}}H$ . One has  $D((\mu \text{Id} + A)^{\frac{1}{2}}) = D(A^{\frac{1}{2}})$  for all  $\mu \geq 0$  (see [3, Proposition 3.8.2, p. 165]). The domain of  $A^{\frac{1}{2}}$  is of importance since it describes the initial values  $u_0$  for which the Cauchy problem

$$\dot{u} + Au = 0, \quad u(0) = u_0$$

has an  $H^1$ -solution. In fact,  $u(t) := e^{-tA}u_0$  is the mild solution of this problem which is defined for all  $u_0 \in H$ . Lions and Magenes (see [16, Théorème 10.1]) showed the equivalence

$$u \in H^1(0, T; H) \quad \text{if and only if} \quad u_0 \in D(A^{\frac{1}{2}}). \quad (2.12)$$

The space  $V$  is in general known, it is typically a Sobolev space as  $H^1(\Omega)$  or  $H_0^1(\Omega)$ . However, the right space  $D(A^{\frac{1}{2}})$  for the admissible initial values is in general different from  $V$ . We introduce a name to describe the important property that both spaces coincide.

**Definition 2.1.** The form  $\mathfrak{a}$  has the *square root property* if  $D(A^{\frac{1}{2}}) = V$ .

We give an abstract criterion for a particular case where the square root property holds.

**Example 2.2.** Assume that  $\mathbf{a}$  can be written in the form  $\mathbf{a} = \mathbf{a}_1 + \mathbf{a}_2$  where  $\mathbf{a}_1 : V \times V \rightarrow \mathbb{C}$  is bounded and symmetric and  $\mathbf{a}_2 : V \times H \rightarrow \mathbb{C}$  is bounded. Then  $\mathbf{a}$  has the square root property. See McIntosh [17].

Not each form has the square root property. The famous solution of the Kato square root problem says that elliptic forms describing a second order differential operator with measurable coefficients on bounded open sets of  $\mathbb{R}^N$  with Dirichlet or Neumann boundary conditions have the square root property (see [5] for the case of  $\Omega = \mathbb{R}^N$  and [6] for the case of strongly Lipschitz domains). We will need the following result by J.-L. Lions [15, Théorème 5.1].

**Lemma 2.3.** *The form  $\mathbf{a}$  has the square root property if and only if  $D(A^{\frac{1}{2}}) \subset V$  and  $D(A^{*\frac{1}{2}}) \subset V$ .*

Note that  $e|_H^{-tA} = e^{-tA}$  for all  $t \geq 0$ . We obtain from (2.1)–(2.6) and (2.9) the following estimates for the semigroup.

**Proposition 2.4.** *There exists  $c > 0$  such that for all  $\ell \in [0, 1]$  and all  $t > 0$ ,*

$$\|e^{-tA}\|_{\mathcal{L}(V'_\ell, V)} \leq \frac{c}{t^{\frac{1+\ell}{2}}} \quad (2.13)$$

$$\|e^{-tA}\|_{\mathcal{L}(V_\ell, V)} \leq \frac{c}{t^{\frac{1-\ell}{2}}} \quad (2.14)$$

$$\|e^{-tA}\|_{\mathcal{L}(V'_\ell, H)} \leq \frac{c}{t^{\frac{\ell}{2}}} \quad (2.15)$$

We will consider form perturbations which are continuous on  $V \times V_\gamma$  and on  $V_\gamma \times V$ . They preserve the square root property.

**Proposition 2.5.** *Let  $\mathbf{a}_1, \mathbf{a}_2 : V \times V \rightarrow \mathbb{C}$  be two bounded, coercive forms. Assume that there exists a constant  $c > 0$  such that*

$$|\mathbf{a}_1(u, v) - \mathbf{a}_2(u, v)| \leq c \|u\|_V \|v\|_{V_\gamma} \quad u, v \in V,$$

where  $0 \leq \gamma < 1$ . Then,  $\mathbf{a}_1$  has the square root property if, and only if,  $\mathbf{a}_2$  has it.

In the following proof the constant  $c > 0$  will vary from one line to the other but does not depend on the variables to be estimated. We keep this convention throughout the paper.

*Proof.* By hypothesis we have  $A_1^{-1/2}H \subset V$ . We show that  $(A_1^{-1/2} - A_2^{-1/2})H \subset V$ . Let  $u \in H$ . Then

$$\begin{aligned} A_1^{-1/2}u - A_2^{-1/2}u &= \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{1}{\sqrt{\sigma}} (e^{-\sigma A_1}u - e^{-\sigma A_2}u) d\sigma \\ &= \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{1}{\sqrt{\sigma}} \frac{1}{2\pi i} \int_\Gamma e^{-\sigma \lambda} ((\lambda \text{Id} - A_1)^{-1}u - (\lambda \text{Id} - A_2)^{-1}u) d\lambda d\sigma \\ &= \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{1}{\sqrt{\sigma}} \frac{1}{2\pi i} \int_\Gamma e^{-\sigma \lambda} (\lambda \text{Id} - \mathcal{A}_1)^{-1} (\mathcal{A}_1 - \mathcal{A}_2) (\lambda \text{Id} - A_2)^{-1}u d\lambda d\sigma. \end{aligned}$$

Since  $\mathcal{A}_1 - \mathcal{A}_2 \in \mathcal{L}(V, V'_\gamma)$  and (2.5) for  $\ell = 1$  and (2.3) for  $\ell = \gamma$

$$\begin{aligned} \|(\lambda \text{Id} - A_2)^{-1}\|_{\mathcal{L}(H, V)} &\leq \frac{c}{(1+|\lambda|)^{1/2}} \\ \text{and} & \qquad \qquad \qquad \text{for all } \lambda \in \Gamma \\ \|(\lambda \text{Id} - \mathcal{A}_1)^{-1}\|_{\mathcal{L}(V'_\gamma, V)} &\leq \frac{c}{(1+|\lambda|)^{\frac{1-\gamma}{2}}} \end{aligned}$$

we see that  $(\lambda \text{Id} - \mathcal{A}_1)^{-1}(\mathcal{A}_1 - \mathcal{A}_2)(\lambda \text{Id} - A_2)^{-1}u \in V$  and the integral converges in  $V$ . In fact,

$$\|(\lambda \text{Id} - \mathcal{A}_1)^{-1}(\mathcal{A}_1 - \mathcal{A}_2)(\lambda \text{Id} - A_2)^{-1}u\|_V \leq \frac{c}{(1+|\lambda|)^{1-\gamma/2}} \|u\|_H \quad (2.16)$$

and therefore

$$\begin{aligned} \|A_1^{-1/2}u - A_2^{-1/2}u\|_V &\leq c \int_0^\infty \frac{1}{\sqrt{\sigma}} \left| \int_\Gamma e^{-\sigma \Re e \lambda} \frac{1}{(1+|\lambda|)^{\frac{1-\gamma}{2}}} \frac{1}{(1+|\lambda|)^{1/2}} \|u\|_H d|\lambda| \right| d\sigma \\ &\leq c \left( \int_0^\infty \frac{1}{\sqrt{\sigma}} \int_0^\infty e^{-\sigma r \cos \vartheta} \frac{1}{(1+r)^{1-\gamma/2}} dr d\sigma \right) \|u\|_H \\ &\leq c \|u\|_H \int_0^\infty \left( \int_0^\infty \frac{1}{\sqrt{\sigma}} e^{-\sigma r \cos \vartheta} d\sigma \right) \frac{1}{(1+r)^{1-\gamma/2}} dr \\ &\leq c \|u\|_H \int_0^\infty \left( \int_0^\infty \frac{\sqrt{r}}{\sqrt{s}} e^{-s \cos \vartheta} \frac{1}{r} ds \right) \frac{1}{(1+r)^{1-\gamma/2}} dr \\ &\leq c \|u\|_H \int_0^\infty \frac{1}{\sqrt{r}} \frac{1}{(1+r)^{1-\gamma/2}} dr \leq c \|u\|_H. \end{aligned}$$

Thus the claim is proved and  $D(A_2^{\frac{1}{2}}) \subset V$ . Applying this result to  $(\mathbf{a}_1^*, \mathbf{a}_2^*)$  instead of  $(\mathbf{a}_1, \mathbf{a}_2)$  we find that  $D(A_2^{\frac{1}{2}}) \subset V$ . Indeed we have  $\mathcal{A}_1^* - \mathcal{A}_2^* \in \mathcal{L}(V_\gamma, V')$ . Thanks to (2.3) for  $\ell = 1$  and (2.5) for  $\ell = \gamma$ , (2.16) becomes

$$\|(\lambda \text{Id} - \mathcal{A}_1^*)^{-1}(\mathcal{A}_1^* - \mathcal{A}_2^*)(\lambda \text{Id} - A_2^*)^{-1}u\|_V \leq \frac{c}{(1+|\lambda|)^{1-\gamma/2}} \|u\|_H$$

It follows from Lemma 2.3 that  $\mathbf{a}_2$  has the square root property.  $\square$

### 3 Non-autonomous forms

In this section, we consider a time-dependent form  $\mathbf{a}$ . Let  $V, H$  be separable complex Hilbert spaces. Let  $T > 0$  and let

$$\begin{aligned} \mathbf{a}(t; \cdot, \cdot) : \times V \times V &\rightarrow \mathbb{C} \quad \text{be a sesquilinear form for all } t \in [0, T] \text{ satisfying} \\ (1.1) \text{ (boundedness)} \text{ and } (1.2) \text{ (coercivity)} & \end{aligned} \quad (3.1)$$

$$\mathbf{a}(\cdot, \cdot; u, v) : [0, T] \rightarrow \mathbb{C} \text{ is measurable for all } u, v \in V. \quad (3.2)$$

Then for each  $t \in [0, T]$  we consider the operator  $\mathcal{A}(t)$  on  $V'$  which is associated with  $\mathbf{a}(t; \cdot, \cdot)$  and we denote by  $A(t)$  the part of  $\mathcal{A}(t)$  in  $H$ . A classical theorem of Lions (see [7,

Théorème 1 p.619, Théorème 2 p.620, Chap.XVIII §3], [21, Proposition 2.3, Chap. III.2]) establishes well-posedness and maximal regularity in  $V'$  of the Cauchy problem

$$\begin{cases} \dot{u}(t) + \mathcal{A}(t)u(t) &= f(t) \\ u(0) &= u_0. \end{cases} \quad (3.3)$$

More precisely, we let

$$MR(V, V') := H^1(0, T; V') \cap L^2(0, T; V).$$

Then  $MR(V, V') \subset \mathcal{C}([0, T]; H)$  and the following holds.

**Theorem 3.1** (Lions). *Let  $f \in L^2(0, T; V')$ ,  $u_0 \in H$ . Then there exists a unique solution  $u \in MR(V, V')$  of (3.3).*

The operator  $\mathcal{A}(t)$  is not the real object of interest if one considers boundary value problems (see Section 5), it is rather its part in  $H$  which realizes the boundary conditions. So the following question is of great importance.

**Question 3.2.** Assume that  $f \in L^2(0, T; H)$  and  $u_0 \in V$ . Does it follow that the solution  $u \in MR(V, V')$  of (3.3) is actually in  $H^1(0, T; H)$ ?

In that case, since  $u$  is a solution,  $u(t) \in D(A(t))$  a.e. and  $\dot{u}(t) + A(t)u(t) = f(t)$ . We have seen that we have to impose at least that  $\mathfrak{a}(0; \cdot, \cdot)$  has the square root property (since otherwise, even for  $f \equiv 0$  and even for  $A(t) \equiv A(0)$  there exists a counterexample). Our aim is to give a positive answer to the question if  $\mathfrak{a}$  satisfies some further regularity in time.

**Definition 3.3.** The form  $\mathfrak{a}$  (or Problem (3.3)) satisfies *maximal regularity in  $H$*  if for each  $f \in L^2(0, T; H)$  and for each  $u_0 \in V$ , the solution  $u \in MR(V, V')$  of (3.3) is actually in  $H^1(0, T; H)$ .

The problem (3.3) is invariant under shifting the operator by a scalar operator as the following proposition shows.

**Proposition 3.4.** *Let  $\mu \in \mathbb{R}$ .*

1. *For each  $f \in L^2(0, T; V')$  and  $u_0 \in H$  there is a unique  $u \in MR(V, V')$  such that*

$$\begin{cases} \dot{u}(t) + \mathcal{A}(t)u(t) + \mu u(t) &= f(t) \quad a.e., \\ u(0) &= u_0. \end{cases} \quad (3.4)$$

2. *If problem (3.3) has maximal regularity in  $H$  and if  $u_0 \in V$  and  $f \in L^2(0, T; H)$ , then this solution  $u$  of (3.4) belongs to  $H^1(0, T; H)$ .*

*Proof.* 1. Let  $v$  be the unique solution in  $MR(V, V')$  of

$$\begin{cases} \dot{v}(t) + \mathcal{A}(t)v(t) &= e^{\mu t} f(t) \quad a.e. \\ v(0) &= u_0 \end{cases}$$

and let  $u$  be defined by  $u(t) = e^{-\mu t}v(t)$ ,  $t \in [0, T]$ . It is immediate that  $u \in MR(V, V')$ ,  $u(0) = u_0$  and

$$\begin{aligned}\dot{u}(t) &= -\mu e^{-\mu t}v(t) + e^{-\mu t}\dot{v}(t) \\ &= -\mu u(t) + e^{-\mu t}(-\mathcal{A}(t)v(t) + f(t)e^{\mu t}) \\ &= -\mu u(t) - \mathcal{A}(t)u(t) + f(t), \quad a.e. \ t \in [0, T]\end{aligned}$$

which proves that  $u$  satisfies (3.4). Assume now that (3.4) admits two solutions in  $MR(V, V')$   $u_1$  and  $u_2$ . Then  $v_1(t) = e^{\mu t}u_1(t)$  and  $v_2(t) = e^{\mu t}u_2(t)$  define two solutions of (3.3) in  $MR(V, V')$  with initial value  $u_0$  and  $e^{\mu \cdot}f(\cdot)$  instead of  $f$ . Therefore they coincide by Lions' Theorem 3.1.

2. Assume now that  $u$  is a solution in  $MR(V, V')$  of (3.4). Let  $v : t \mapsto e^{\mu t}u(t)$ ;  $v$  is the unique solution of (3.3) in  $MR(V, V')$  with  $g = e^{\mu \cdot}f(\cdot)$  instead of  $f$ . Since problem (3.3) has the maximal regularity property and  $g \in L^2(0, T; H)$ ,  $v(0) = u_0 \in V$ , the solution  $v$  belongs to  $H^1(0, T; H)$  and  $t \mapsto \mathcal{A}(t)v(t) \in L^2(0, T; H)$ . This proves that  $u : t \mapsto e^{-\mu t}v(t)$  also belongs to  $H^1(0, T; H)$  (since  $\dot{u}(t) = \mu e^{-\mu t}v(t) + e^{-\mu t}\dot{v}(t)$ ) and  $t \mapsto \mathcal{A}(t)u(t) = e^{-\mu t}\mathcal{A}(t)v(t) \in L^2(0, T; H)$ .

□

Finally, we establish a representation formula of the solution  $u \in MR(V, V')$  of (3.3).

**Proposition 3.5.** *1. Let  $f \in L^2(0, T; V')$ ,  $u_0 \in H$ . Let  $u \in MR(V, V')$  be the solution of (3.3). Then*

$$\begin{aligned}u(t) &= e^{-t\mathcal{A}(t)}u_0 + \int_0^t e^{-(t-s)\mathcal{A}(t)}f(s) \, ds + \int_0^t e^{-(t-s)\mathcal{A}(t)}(\mathcal{A}(t) - \mathcal{A}(s))u(s) \, ds, \\ &\text{for all } t \in [0, T].\end{aligned}\tag{3.5}$$

2. Moreover, there is only one  $u \in MR(V, V')$  satisfying this identity if we assume in addition that  $t \mapsto \mathcal{A}(t) \in \mathcal{L}(V, V')$  is Dini-continuous, i.e., admits a modulus of continuity  $\omega$  in the operator norm with the property that  $t \mapsto \frac{1}{t}\omega(t) \in L^1(0, T)$ .

*Proof.* 1. This formula already appears in [1, formula (1.18), p.57] for operators with different properties; see also [10, Lemma 2.4]. Let  $0 < t \leq T$ . Consider the function  $v : [0, t] \ni s \mapsto e^{-(t-s)\mathcal{A}(t)}u(s)$ . Then  $v \in \mathcal{C}([0, t]; H) \cap H^1(0, t; V')$  and

$$\begin{aligned}\dot{v}(s) &= \mathcal{A}(t)e^{-(t-s)\mathcal{A}(t)}u(s) + e^{-(t-s)\mathcal{A}(t)}\dot{u}(s) \\ &= e^{-(t-s)\mathcal{A}(t)}\left(\mathcal{A}(t)u(s) + (-\mathcal{A}(s)u(s) + f(s))\right) \\ &= e^{-(t-s)\mathcal{A}(t)}(\mathcal{A}(t) - \mathcal{A}(s))u(s) + e^{-(t-s)\mathcal{A}(t)}f(s).\end{aligned}\tag{3.6}$$

Thus integrating between 0 and  $t$  gives  $v(t) = \int_0^t \dot{v}(s) \, ds + v(0)$  which is the claim.



2. In order to prove uniqueness, assume that there are two functions  $u_1$  and  $u_2$  in the space  $MR(V, V')$  satisfying (3.5) and denote by  $w$  the difference  $u_1 - u_2$ . Then  $w \in MR(V, V')$  and satisfies

$$w(t) = \int_0^t e^{-(t-s)\mathcal{A}(t)} (\mathcal{A}(t) - \mathcal{A}(s)) w(s) ds, \quad (3.7)$$

for all  $t \in [0, T]$ . Let  $\tau := \sup\{t \in [0, T]; w(s) = 0 \text{ on } [0, t]\}$ . Assume that  $\tau < T$ . Using (3.7), the continuity properties of  $\mathcal{A}(\cdot)$  and the estimate (2.13) for  $\ell = 1$ , we obtain for all  $\tau \leq t \leq T$

$$\|w(t)\|_V \leq \int_\tau^t \frac{c}{t-s} \omega(t-s) \|w(s)\|_V ds$$

and therefore by Young's inequality for convolution

$$\|w\|_{L^2(\tau, t; V)} \leq c \left( \int_0^{t-\tau} \frac{\omega(s)}{s} ds \right) \|w\|_{L^2(\tau, t; V)}.$$

Choosing  $\varepsilon$  small enough so that  $\tau + \varepsilon \leq T$  and  $c \left( \int_0^\varepsilon \frac{\omega(s)}{s} ds \right) < 1$ , we proved that  $w(t) = 0$  almost everywhere on  $[\tau, \tau + \varepsilon]$ . Since  $MR(V, V') \subset \mathcal{C}([0, T]; H)$ , we have then  $w(t) = 0$  everywhere on  $[\tau, \tau + \varepsilon]$ , which contradict the definition of  $\tau$ . This proves that  $\tau = T$  and ultimately  $w(t) = 0$  on  $[0, T]$ .  $\square$

## 4 Maximal regularity in $H$

Let  $V$  and  $H$  be two separable complex Hilbert spaces such that  $V \hookrightarrow_d H$ . Let  $\mathfrak{a} : [0, T] \times V \times V \rightarrow \mathbb{C}$  be a non-autonomous form satisfying (3.1) and (3.2). We denote by  $\mathcal{A}(t)$  the operator on  $V'$  associated with  $\mathfrak{a}(t; \cdot, \cdot)$  and by  $A(t)$  its part in  $H$ . The essential further condition concerns continuity in time. We assume that there exist  $0 \leq \gamma < 1$  and a modulus of continuity  $\omega$  such that

$$|\mathfrak{a}(t; u, v) - \mathfrak{a}(s; u, v)| \leq \omega(|t - s|) \|u\|_V \|v\|_{V_\gamma} \quad (4.1)$$

for all  $t, s \in [0, T]$ ,  $u, v \in V_\gamma$ . We suppose that  $\omega : [0, T] \rightarrow [0, +\infty)$  is continuous and satisfies

$$\sup_{t \in [0, T]} \frac{\omega(t)}{t^{\gamma/2}} < \infty \quad (4.2)$$

$$\text{and } \int_0^T \frac{\omega(t)}{t^{1+\gamma/2}} dt < \infty. \quad (4.3)$$

The main example of such a continuity modulus is the function  $\omega(t) = t^\alpha$  with  $\alpha > \frac{\gamma}{2}$ . We remark that conditions (4.2), (4.3) imply that

$$\int_0^T \frac{\omega(t)^2}{t^{1+\gamma}} dt < \infty. \quad (4.4)$$

Finally, we impose that  $\mathfrak{a}(0; \cdot, \cdot)$  has the square root property. By Proposition 2.5 this implies that  $\mathfrak{a}(t; \cdot, \cdot)$  has the square root property for all  $t \in [0, T]$ . Under the preceding conditions we have the following result on *maximal regularity in  $H$* .

**Theorem 4.1.** Assume that  $\mathfrak{a}(0; \cdot, \cdot)$  has the square root property. Let  $u_0 \in V$ ,  $f \in L^2(0, T; H)$ . Then there exists a unique  $u \in H^1(0, T; H) \cap L^2(0, T; V)$  such that  $u(t) \in D(A(t))$  a.e. and

$$\begin{cases} \dot{u}(t) + A(t)u(t) = f(t) & \text{a.e.} \\ u(0) = u_0. \end{cases} \quad (4.5)$$

Thus the solution  $u$  is in the space

$$MR_{\mathfrak{a}} := \{u \in H^1(0, T; H) \cap L^2(0, T; V) : u(0) \in V, \mathcal{A}(\cdot)u(\cdot) \in L^2(0, T; H)\}.$$

We will see below that  $MR_{\mathfrak{a}} \subset \mathcal{C}([0, T]; V)$ .

**Remark 4.2.** (a) The space  $MR_{\mathfrak{a}}$  endowed with the norm

$$\|u\|_{MR_{\mathfrak{a}}} = \|\dot{u}\|_{L^2(0, T; H)} + \|A(\cdot)u(\cdot)\|_{L^2(0, T; H)} + \|u(0)\|_V$$

is a Banach space.

(b) It follows from the Closed Graph Theorem that there exists a constant  $c > 0$  such that

$$\|u\|_{H^1(0, T; H)} + \|\mathcal{A}(\cdot)u(\cdot)\|_{L^2(0, T; H)} \leq c(\|u_0\|_V + \|f\|_{L^2(0, T; H)})$$

for each  $u_0 \in V$  and  $f \in L^2(0, T; H)$ , where  $u$  is the solution of (4.5).

*Proof of Theorem 4.1.* By Lions' Theorem 3.1 there exists a unique solution  $u \in MR(V, V')$  of the problem. We have to show that  $\mathcal{A}(\cdot)u(\cdot) \in L^2(0, T; H)$ . For that we use the decomposition (3.5) and show that  $\mathcal{A}(\cdot)u_j(\cdot) \in L^2(0, T; H)$  for  $j = 1, 2, 3$  where

$$u_1(t) = e^{-tA(t)}u_0, \quad u_2(t) = \int_0^t e^{-(t-s)A(s)}f(s) \, ds,$$

$$\text{and } u_3(t) = \int_0^t e^{-(t-s)\mathcal{A}(t)}(\mathcal{A}(t) - \mathcal{A}(s))u(s) \, ds.$$

Remark that (4.1) implies for all  $t, s \in [0, T]$

$$\mathcal{A}(t) - \mathcal{A}(s) \in \mathcal{L}(V, V'_\gamma) \quad \text{and} \quad \|\mathcal{A}(t) - \mathcal{A}(s)\|_{\mathcal{L}(V, V'_\gamma)} \leq c\omega(|t - s|). \quad (4.6)$$

We divide the proof into three steps.

**Step 1:** We adapt the proof of [10, Lemma 2.7] to our situation. Since  $\mathfrak{a}(0; \cdot, \cdot)$  has the square root property, thanks to (2.12),  $t \mapsto A(0)e^{-tA(0)}u_0 \in L^2(0, T; H)$ . Thus it suffices to show that

$$\phi : t \mapsto A(t)e^{-tA(t)}u_0 - A(0)e^{-tA(0)}u_0 \in L^2(0, T; H).$$

Using

$$(\lambda \text{Id} - A(t))^{-1} - (\lambda \text{Id} - A(0))^{-1} = (\lambda \text{Id} - \mathcal{A}(t))^{-1}(\mathcal{A}(t) - \mathcal{A}(0))(\lambda \text{Id} - A(0))^{-1} \quad (4.7)$$

we see that

$$\phi(t) = \frac{1}{2\pi i} \int_{\Gamma} \lambda e^{-\lambda t} (\lambda \text{Id} - A(t))^{-1} (\mathcal{A}(t) - \mathcal{A}(0)) (\lambda \text{Id} - A(0))^{-1} u_0 \, d\lambda.$$

Using (2.5) for  $\ell = \gamma$  and (2.2) for  $\ell = 1$  we estimate

$$\begin{aligned}
\|\phi(t)\|_H &\leq c \int_{\Gamma} |\lambda| e^{-t\Re \lambda} \frac{1}{|\lambda|^{1-\gamma/2}} \omega(t) \|(\lambda \text{Id} - A(0))^{-1} u_0\|_V d|\lambda| \\
&\leq c \omega(t) \int_{\Gamma} |\lambda| e^{-t\Re \lambda} \frac{1}{|\lambda|^{1-\gamma/2}} \frac{1}{|\lambda|} \|u_0\|_V d|\lambda| \\
&\leq c \omega(t) \|u_0\|_V \int_0^\infty e^{-tr \cos \vartheta} \frac{1}{r^{1-\gamma/2}} dr \\
&\leq c \omega(t) \|u_0\|_V \int_0^\infty e^{-\rho \cos \vartheta} \frac{t^{1-\gamma/2}}{\rho^{1-\gamma/2}} \frac{1}{t} d\rho \leq c \frac{\omega(t)}{t^{\gamma/2}} \|u_0\|_V.
\end{aligned}$$

It follows from (4.2) that  $\phi \in L^2(0, T; H)$ .

**Step 2:** We show that  $t \mapsto \int_0^t A(t) e^{-(t-s)A(t)} f(s) ds \in L^2(0, T; H)$ . The proof here is much more elementary than the one of [10, Lemma 2.5] thanks to our stronger condition (4.1) on the form  $\mathfrak{a}$ . By (2.10)  $A(0)$  satisfies maximal regularity and

$$t \mapsto \int_0^t A(0) e^{-(t-s)A(0)} f(s) ds \in L^2(0, T; H).$$

Thus it suffices to show that

$$\phi : t \mapsto \int_0^t A(t) e^{-(t-s)A(t)} f(s) ds - \int_0^t A(0) e^{-(t-s)A(0)} f(s) ds \in L^2(0, T; H).$$

As before we have

$$\phi(t) = \int_0^t \frac{1}{2\pi i} \int_{\Gamma} \lambda e^{-(t-s)\lambda} (\lambda \text{Id} - \mathcal{A}(t))^{-1} (\mathcal{A}(t) - \mathcal{A}(0)) (\lambda \text{Id} - A(0))^{-1} f(s) d\lambda ds.$$

Using (2.4) for  $\ell = \gamma$  and (2.5) for  $\ell = 1$  we obtain

$$\begin{aligned}
\|\phi(t)\|_H &\leq c \int_0^t \int_0^\infty r e^{-(t-s)r \cos \vartheta} \frac{1}{r^{1-\gamma/2}} \omega(t) \frac{1}{r^{1/2}} \|f(s)\|_H dr ds \\
&= c \omega(t) \int_0^t \left( \int_0^\infty e^{-\rho \cos \vartheta} \rho^{\frac{\gamma-1}{2}} d\rho \right) (t-s)^{-\frac{1+\gamma}{2}} \|f(s)\|_H ds \\
&= c \omega(t) (h * \|f(\cdot)\|_H)(t) \quad \text{for all } 0 \leq t \leq T
\end{aligned}$$

where  $h$  is defined by  $h(t) = \begin{cases} 0 & \text{for } t \leq 0 \\ t^{-\frac{1+\gamma}{2}} & \text{for } 0 < t \leq T \\ 0 & \text{for } t \geq T \end{cases}$  is in  $L^1(\mathbb{R})$  since  $\frac{1+\gamma}{2} < 1$ . It follows that

$$\int_0^T \|\phi(t)\|_H^2 dt < \infty.$$

**Step 3:** In order to show that  $A(\cdot)u_3(\cdot) \in L^2(0, T; H)$ , we define for  $g \in L^2(0, T; H)$ ,

$$\begin{aligned}
(Qg)(t) &:= \int_0^t A(t) e^{-(t-s)A(t)} (\mathcal{A}(t) - \mathcal{A}(s)) A(s)^{-1} g(s) ds \\
&= \int_0^t A(t) e^{-\frac{t-s}{2}A(t)} e^{-\frac{t-s}{2}\mathcal{A}(t)} (\mathcal{A}(t) - \mathcal{A}(s)) A(s)^{-1} g(s) ds.
\end{aligned}$$

The proof here is very much inspired by [1, end of §1] (see also [11, beginning of §3], [10, Proof of Theorem 1.2]). Let  $\varepsilon > 0$ . Replacing  $A(s)$  by  $A(s) + \mu \text{Id}$  (see Proposition 3.4) we may assume that  $\|A(s)^{-1}\|_{\mathcal{L}(H,V)} \leq \varepsilon$  (see (2.5) for  $\ell = 1$ ) for all  $s \geq 0$ . Thus by (2.15) for  $\ell = \gamma$  and (2.9) we have the following estimates

$$\begin{aligned} \|Qg(t)\|_H &\leq \int_0^t \|A(t)e^{-\frac{t-s}{2}A(t)}\|_{\mathcal{L}(H)} \|e^{-\frac{t-s}{2}\mathcal{A}(t)}\|_{\mathcal{L}(V'_\gamma, H)} \omega(t-s) \varepsilon \|g(s)\|_H ds \\ &\leq c\varepsilon \int_0^t \frac{1}{t-s} \frac{1}{(t-s)^{\gamma/2}} \omega(t-s) \|g(s)\|_H ds. \end{aligned}$$

Since  $k(t) = \frac{1}{t^{1+\gamma/2}} \omega(t)$  defines a function  $k \in L^1(0, T)$  thanks to (4.3), it follows that  $Qg \in L^2(0, T; H)$  and

$$\|Qg\|_{L^2(0, T; H)} \leq c\varepsilon \|g\|_{L^2(0, T; H)}.$$

Choosing  $\varepsilon > 0$  small enough we can arrange that  $\|Q\|_{\mathcal{L}(L^2(0, T; H))} \leq \frac{1}{2}$ . Thus  $\text{Id} - Q$  is invertible. By Step 1 and Step 2 we know that

$$h := A(\cdot)(u_1(\cdot) + u_2(\cdot)) \in L^2(0, T; H).$$

Let  $w = A(\cdot)^{-1}(\text{Id} - Q)^{-1}h$ . Then  $A(\cdot)w(\cdot) \in L^2(0, T; H)$ . Since  $h = A(\cdot)w(\cdot) - Q(A(\cdot)w(\cdot))$  one has

$$\begin{aligned} A(t)e^{-tA(t)}u_0 + A(t) \int_0^t e^{-(t-s)A(t)} f(s) ds + A(t) \int_0^t e^{-(t-s)A(t)} (\mathcal{A}(t) - \mathcal{A}(s))w(s) ds \\ = A(t)w(t). \end{aligned}$$

Applying  $A(t)^{-1}$  on both sides we see from Proposition 3.5 that  $w = u$ . Hence  $A(\cdot)u(\cdot) = A(\cdot)w(\cdot) \in L^2(0, T; H)$ .  $\square$

**Remark 4.3.** If we do not suppose the square root property, then the proof of Theorem 4.1 shows that for  $u_0 \in D(A(0)^{\frac{1}{2}})$ ,  $f \in L^2(0, T; H)$ , the solution  $u$  given by Lions' Theorem is in  $H^1(0, T; H)$ , i.e., we have the same conclusion as in Theorem 4.1 if we choose the right trace space.

As announced above, we now show that

**Theorem 4.4.** *The space  $MR_{\mathfrak{a}}$  is continuously embedded into  $\mathcal{C}([0, T], V)$ .*

To prove this theorem, we need the following lemma.

**Lemma 4.5.** *Let  $\mathfrak{a}$  be a non-autonomous form satisfying (3.1), (3.2) and (4.1). Denote by  $\mathcal{A}(t) : V \rightarrow V'$  the operator associated with  $\mathfrak{a}(t; \cdot, \cdot)$  and by  $A(t)$  its part in  $H$ . Then for all  $\lambda \notin \Sigma_\theta$ ,  $s \in (0, T]$  and  $\sigma > 0$  the following mappings*

$$t \mapsto (\lambda \text{Id} - tA(0))^{-1} \in \mathcal{L}(V), \quad (4.8)$$

$$t \mapsto \mathcal{A}(t) - \mathcal{A}(s) \in \mathcal{L}(V_\gamma, V'_\gamma), \quad (4.9)$$

$$t \mapsto (\lambda \text{Id} - \mathcal{A}(t))^{-1} \in \mathcal{L}(V'_\gamma, V), \quad (4.10)$$

$$t \mapsto (\lambda \text{Id} - tA(t))^{-1} \in \mathcal{L}(V'_\gamma, V), \quad (4.11)$$

$$t \mapsto e^{-\sigma \mathcal{A}(t)} \in \mathcal{L}(V'_\gamma, V) \quad (4.12)$$

are continuous on  $(0, T]$ .

*Proof.* To prove (4.8), we write for  $t, s \in (0, T]$  and  $\lambda \notin \Sigma_\theta$

$$(\lambda \text{Id} - tA(0))^{-1} - (\lambda \text{Id} - sA(0))^{-1} = \left(\frac{1}{s} - \frac{1}{t}\right) tA(0)(\lambda \text{Id} - tA(0))^{-1} \left(\frac{\lambda}{s} \text{Id} - A(0)\right)^{-1}.$$

Thanks to (2.2) for  $\ell = 1$  the operators  $tA(0)(\lambda \text{Id} - tA(0))^{-1}$  and  $\left(\frac{\lambda}{s} \text{Id} - A(0)\right)^{-1}$  are uniformly (w.r.t.  $t, s$  and  $\lambda$ ) bounded in  $V$ . Therefore

$$(\lambda \text{Id} - tA(0))^{-1} - (\lambda \text{Id} - sA(0))^{-1} \longrightarrow 0 \quad \text{in } \mathcal{L}(V) \text{ as } s \rightarrow t.$$

The claim (4.9) follows immediately from (4.1) since the latter implies

$$\|\mathcal{A}(t) - \mathcal{A}(s)\|_{\mathcal{L}(V, V'_\gamma)} \leq \omega(|t - s|) \longrightarrow 0 \quad \text{as } s \rightarrow t. \quad (4.13)$$

We now prove (4.10) as follows. Let  $t, s \in (0, T]$  and  $\lambda \notin \Sigma_\theta$ . We have

$$(\lambda \text{Id} - \mathcal{A}(t))^{-1} - (\lambda \text{Id} - \mathcal{A}(s))^{-1} = (\lambda \text{Id} - \mathcal{A}(t))^{-1} (\mathcal{A}(t) - \mathcal{A}(s)) (\lambda \text{Id} - \mathcal{A}(s))^{-1}. \quad (4.14)$$

Therefore, by (2.3) for  $\ell = \gamma$ , using (4.13) we have

$$\|(\lambda \text{Id} - \mathcal{A}(t))^{-1} - (\lambda \text{Id} - \mathcal{A}(s))^{-1}\|_{\mathcal{L}(V'_\gamma, V)} \leq \frac{c\omega(|t - s|)}{(1 + |\lambda|)^{1-\gamma}} \longrightarrow 0 \quad \text{as } s \rightarrow t,$$

which proves (4.10). The proof of (4.11) combines the ideas of the proofs of (4.8) and (4.10). We write

$$\begin{aligned} (\lambda \text{Id} - t\mathcal{A}(t))^{-1} - (\lambda \text{Id} - s\mathcal{A}(s))^{-1} &= \left(\frac{1}{s} - \frac{1}{t}\right) A(t) \left(\frac{\lambda}{t} \text{Id} - A(t)\right)^{-1} \left(\frac{\lambda}{s} \text{Id} - \mathcal{A}(s)\right)^{-1} \\ &\quad + \frac{1}{t} \left(\frac{\lambda}{t} \text{Id} - \mathcal{A}(t)\right)^{-1} (\mathcal{A}(t) - \mathcal{A}(s)) \left(\frac{\lambda}{s} \text{Id} - \mathcal{A}(s)\right)^{-1} \end{aligned}$$

which implies the following estimate thanks to (2.2) for  $\ell = 1$ , (2.3) for  $\ell = \gamma$  and (4.13)

$$\begin{aligned} &\|(\lambda \text{Id} - t\mathcal{A}(t))^{-1} - (\lambda \text{Id} - s\mathcal{A}(s))^{-1}\|_{\mathcal{L}(V'_\gamma, V)} \\ &\leq \frac{c}{(1 + \frac{|\lambda|}{s})^{\frac{1-\gamma}{2}}} \left( (c+1) \left| \frac{1}{s} - \frac{1}{t} \right| + \frac{1}{t} \frac{c\omega(|t - s|)}{(1 + \frac{|\lambda|}{t})^{\frac{1-\gamma}{2}}} \right) \longrightarrow 0 \quad \text{as } s \rightarrow t. \end{aligned}$$

Finally, we show (4.12) using the representation (2.8) for the semigroup and (4.14):

$$e^{-\sigma \mathcal{A}(t)} - e^{-\sigma \mathcal{A}(s)} = \frac{1}{2\pi i} \int_{\Gamma} e^{-\lambda \sigma} (\lambda \text{Id} - \mathcal{A}(t))^{-1} (\mathcal{A}(t) - \mathcal{A}(s)) (\lambda \text{Id} - \mathcal{A}(s))^{-1} d\lambda$$

we obtain, using (2.3) for  $\ell = \gamma$  and (4.13)

$$\begin{aligned} \|e^{-\sigma \mathcal{A}(t)} - e^{-\sigma \mathcal{A}(s)}\|_{\mathcal{L}(V'_\gamma, V)} &\leq c\omega(|t - s|) \int_0^\infty e^{-\sigma r \cos \vartheta} \frac{1}{(1 + r)^{1-\gamma}} dr \\ &\longrightarrow 0 \quad \text{as } s \rightarrow t \quad \text{for all } \sigma > 0, \end{aligned}$$

which proves the claim.  $\square$

*Proof of Theorem 4.4.* Assume that  $u \in MR_a \subset MR(V, V') \subset \mathcal{C}([0, T]; H)$ . Let  $f = \dot{u}(\cdot) + A(\cdot)u(\cdot)$  and  $u_0 = u(0)$ :  $f \in L^2(0, T; H)$ ,  $u_0 \in V$  and  $u$  satisfies (3.3). By Proposition 3.5, we have  $u = u_1 + u_2 + u_3$  where

$$u_1(t) = e^{-tA(t)}u_0, \quad u_2(t) = \int_0^t e^{-(t-s)A(s)}f(s) \, ds,$$

$$\text{and } u_3(t) = \int_0^t e^{-(t-s)A(s)}(\mathcal{A}(t) - \mathcal{A}(s))u(s) \, ds.$$

We will show that each term  $u_j$ ,  $j = 1, 2, 3$ , belongs to  $\mathcal{C}([0, T]; V)$ .

**Step 1:** We claim that  $u_1 \in \mathcal{C}([0, T]; V)$ . Indeed,  $u_0 \in V$  and since  $(e^{-tA(0)})_{t \geq 0}$  defines a  $C_0$ -semigroup one has  $t \mapsto e^{-tA(0)}u_0 \in \mathcal{C}([0, T]; V)$ . Let us first consider the case where  $t > 0$ . We have

$$\begin{aligned} e^{-tA(t)}u_0 - e^{-tA(0)}u_0 &= \frac{1}{2\pi i} \int_{\Gamma} e^{-t\lambda}(\lambda \text{Id} - \mathcal{A}(t))^{-1}(\mathcal{A}(t) - \mathcal{A}(0))(\lambda \text{Id} - A(0))^{-1}u_0 \, d\lambda \\ &= \frac{1}{2\pi i} \int_{\Gamma} e^{-\eta} \left( \frac{\eta}{t} \text{Id} - \mathcal{A}(t) \right)^{-1} (\mathcal{A}(t) - \mathcal{A}(0)) \left( \frac{\eta}{t} \text{Id} - A(0) \right)^{-1} u_0 \frac{d\eta}{t}. \end{aligned} \quad (4.15)$$

Estimates (2.2) for  $\ell = 1$  and (2.3) for  $\ell = \gamma$  imply

$$\begin{aligned} &\left\| \frac{1}{t} e^{-\eta} \left( \frac{\eta}{t} \text{Id} - \mathcal{A}(t) \right)^{-1} (\mathcal{A}(t) - \mathcal{A}(0)) \left( \frac{\eta}{t} \text{Id} - A(0) \right)^{-1} u_0 \right\|_V \\ &\leq \frac{c}{t} e^{-\Re \eta} \frac{1}{(1 + \frac{\eta}{t})^{\frac{1-\gamma}{2}}} \omega(t) \frac{1}{1 + \frac{\eta}{t}} \|u_0\|_V \\ &\leq \frac{c}{t} e^{-|\eta| \cos \vartheta} \frac{1}{(1 + \frac{\eta}{t})^{\frac{1-\gamma}{2}}} \omega(t) \frac{1}{(1 + \frac{\eta}{t})^{1/2 + \gamma/4}} \|u_0\|_V \\ &\leq c e^{-|\eta| \cos \vartheta} t^{-\gamma/4} \omega(t) \frac{1}{|\eta|^{1-\gamma/4}} \|u_0\|_V \\ &\leq c \frac{e^{-|\eta| \cos \vartheta}}{|\eta|^{1-\gamma/4}} (t^{-\gamma/2} \omega(t)) t^{\gamma/4} \|u_0\|_V. \end{aligned}$$

Since  $r \mapsto \frac{e^{-r \cos \vartheta}}{r^{1-\gamma/4}}$  is integrable on  $(0, \infty)$  and

$$t \mapsto e^{-\eta t}(\eta \text{Id} - tA(t))^{-1}(\mathcal{A}(t) - A(0))(\eta \text{Id} - tA(0))^{-1}u_0 \in V$$

is continuous on  $(0, T]$  we may apply Lebesgue's dominated convergence theorem. Therefore we obtain the continuity of  $t \mapsto e^{-tA(t)}u_0 - e^{-tA(0)}u_0 \in V$  on  $(0, T]$ . It remains to prove the continuity at 0. Using the formula (4.15), thanks to (2.2) for  $\ell = 1$  and (2.3) for  $\ell = \gamma$ , we obtain the following estimate

$$\|e^{-tA(t)}u_0 - e^{-tA(0)}u_0\|_V \leq C \omega(t) \int_0^\infty \frac{1}{(1+r)^{\frac{3-\gamma}{2}}} \, dr$$

where we have used that  $|e^{-t\lambda}| \leq 1$  for all  $\lambda \in \Gamma$ . Since  $\omega(t) \rightarrow 0$  as  $t \rightarrow 0$ , this proves that  $t \mapsto e^{-tA(t)}u_0 - e^{-tA(0)}u_0 \in V$  is continuous on  $[0, T]$ , and ultimately that  $u_1$  is continuous on  $[0, T]$ .

**Step 2:** We claim that  $u_2 \in \mathcal{C}([0, T]; V)$ . The embedding

$$H^1(0, T; H) \cap L^2(0, T; D(A(0))) \hookrightarrow \mathcal{C}([0, T]; V)$$

(recall that  $\mathbf{a}(0; \cdot, \cdot)$  has the square root property so that  $V = D(A(0)^{\frac{1}{2}})$ ), together with the maximal regularity property in the autonomous case (2.10) imply that

$$t \mapsto \int_0^t e^{-(t-s)A(0)} f(s) \, ds \in \mathcal{C}([0, T]; V).$$

It suffices to prove now that

$$\phi : t \mapsto \int_0^t e^{-(t-s)A(t)} f(s) \, ds - \int_0^t e^{-(t-s)A(0)} f(s) \, ds \in \mathcal{C}([0, T]; V).$$

For every  $t \in [0, T]$  we have

$$\phi(t) = \int_0^t \frac{1}{2\pi i} \int_{\Gamma} e^{-(t-s)\lambda} (\lambda \text{Id} - \mathcal{A}(t))^{-1} (\mathcal{A}(t) - \mathcal{A}(0)) (\lambda \text{Id} - A(0))^{-1} f(s) \, d\lambda \, ds.$$

This integral is convergent in  $V$ . Indeed, by (2.5) for  $\ell = 1$  and (2.3) for  $\ell = \gamma$ ,

$$\begin{aligned} & \|e^{-(t-s)\lambda} (\lambda \text{Id} - \mathcal{A}(t))^{-1} (\mathcal{A}(t) - \mathcal{A}(0)) (\lambda \text{Id} - A(0))^{-1} f(s)\|_V \\ & \leq c e^{-(t-s)|\lambda| \cos \vartheta} \frac{1}{(1 + |\lambda|)^{\frac{1-\gamma}{2}}} \omega(t) \frac{1}{(1 + |\lambda|)^{1/2}} \|f(s)\|_H \\ & \leq c e^{-(t-s)|\lambda| \cos \vartheta} \frac{\omega(t)}{(1 + |\lambda|)^{1-\gamma/2}} \|f(s)\|_H \end{aligned}$$

and the function  $(s, r) \mapsto \frac{e^{-(t-s)r \cos \vartheta}}{(1+r)^{1-\gamma/2}} \|f(s)\|_H$  is integrable on  $[0, t] \times [0, +\infty)$ . We can then apply Fubini's theorem and obtain the following representation for  $\phi$

$$\phi(t) = \frac{1}{2\pi i} \int_{\Gamma} (\lambda \text{Id} - A(t))^{-1} (A(t) - A(0)) (\lambda \text{Id} - A(0))^{-1} \left( \int_0^t e^{-(t-s)\lambda} f(s) \, ds \right) d\lambda.$$

The following two facts are the keys to prove the continuity of  $\phi$ :

- $t \mapsto \int_0^t e^{-(t-s)\lambda} f(s) \, ds \in \mathcal{C}([0, T]; H)$  and for all  $\lambda \in \Gamma \setminus \{0\}$ ,

$$\left\| \int_0^t e^{-(t-s)\lambda} f(s) \, ds \right\|_H \leq \frac{1}{\sqrt{2|\lambda| \cos \vartheta}} \|f\|_{L^2(0, T; H)};$$

- $t \mapsto (\lambda \text{Id} - A(t))^{-1} (A(t) - A(0)) (\lambda \text{Id} - A(0))^{-1} \in \mathcal{C}([0, T]; \mathcal{L}(H, V))$  thanks to (4.9) and (4.10) and for all  $\lambda \in \Gamma$ ,

$$\|(\lambda \text{Id} - A(t))^{-1} (A(t) - A(0)) (\lambda \text{Id} - A(0))^{-1}\|_{\mathcal{L}(H, V)} \leq \frac{c \omega(t)}{(1 + |\lambda|)^{1-\gamma/2}}.$$

Since  $r \mapsto \frac{1}{\sqrt{r}(1+r)^{1-\gamma/2}} \in L^1(0, \infty)$  we can apply Lebesgue's dominated convergence theorem and we obtain that  $\phi \in \mathcal{C}([0, T]; V)$  and

$$\|\phi(t)\|_V \leq c\omega(t) \left( \int_0^\infty \frac{1}{\sqrt{r}(1+r)^{1-\gamma/2}} dr \right) \|f\|_{L^2(0, T; H)},$$

which proves the claim.

**Step 3:** We conclude in a similar way as in Step 3 of the proof of Theorem 4.1. We define for  $h \in \mathcal{C}([0, T]; V)$

$$\begin{aligned} (Ph)(t) &:= \int_0^t e^{-(t-s)A(t)} (\mathcal{A}(t) - \mathcal{A}(s)) h(s) ds \\ &= A(t)^{-1/2} \int_0^t A(t)^{1/2} e^{-(t-s)A(t)} (\mathcal{A}(t) - \mathcal{A}(s)) h(s) ds. \end{aligned}$$

Thanks to (4.9) and (4.12),  $Ph$  is continuous on  $[0, T]$  with values in  $V$  by Lebesgue's dominated convergence theorem. For all  $t, s \in [0, T]$ ,  $t > s$ , the estimate (2.13) for  $\ell = \gamma$ , the fact that  $(e^{-\sigma A(t)})_{\sigma \geq 0}$  is a holomorphic semigroup in  $V$  and the property (4.1) give

$$\|A(t)^{1/2} e^{-(t-s)A(t)} (\mathcal{A}(t) - \mathcal{A}(s)) h(s)\|_V \leq \frac{c\omega(t-s)}{(t-s)^{1+\gamma/2}} \|h(s)\|_V.$$

Let  $\varepsilon > 0$ . As in Step 3 of the proof of Theorem 4.1, Replacing  $A(t)$  by  $A(t) + \mu \text{Id}$  we may assume that  $\|A(t)^{-1/2}\|_{\mathcal{L}(V)} \leq \varepsilon$ . Therefore, we have

$$\sup_{t \in [0, T]} \|(Ph)(t)\|_V \leq c\varepsilon \left( \int_0^T \frac{\omega(\sigma)}{\sigma^{1+\gamma/2}} d\sigma \right) \sup_{t \in [0, T]} \|h(t)\|_V.$$

Now, thanks to (4.3), choosing  $\varepsilon$  small enough we can arrange that  $\|P\|_{\mathcal{L}(\mathcal{C}([0, T]; V))} \leq \frac{1}{2}$ . Thus  $\text{Id} - P$  is invertible in  $\mathcal{L}(\mathcal{C}([0, T]; V))$ . We have, by definition of  $u_1$  and  $u_2$ , that  $u - Pu = u_1 + u_2$ . Since we have proved in Step 1 and Step 2 that  $u_1 + u_2 \in \mathcal{C}([0, T]; V)$ , it shows that  $u = (\text{Id} - P)^{-1}(u_1 + u_2) \in \mathcal{C}([0, T]; V)$ .  $\square$

## 5 Non-autonomous Robin boundary conditions

Let  $\Omega \subset \mathbb{R}^N$  be a bounded open set with Lipschitz boundary. We denote by  $\partial\Omega$  the boundary of  $\Omega$  and take  $L^2(\partial\Omega)$  with respect to the  $(N-1)$ -dimensional Hausdorff measure. There exists a unique bounded operator  $\text{Tr} : H^1(\Omega) \rightarrow L^2(\partial\Omega)$  such that  $\text{Tr}(u) = u|_{\partial\Omega}$  if  $u \in H^1(\Omega) \cap \mathcal{C}(\overline{\Omega})$ . We call  $\text{Tr}(u)$  the *trace* of  $u$  and also use the notation  $u|_{\partial\Omega}$  for  $u \in H^1(\Omega)$ . Let  $\alpha > \frac{1}{4}$  and  $B : [0, T] \rightarrow \mathcal{L}(L^2(\partial\Omega))$  be a mapping such that

$$\|B(t) - B(s)\|_{\mathcal{L}(L^2(\partial\Omega))} \leq c|t - s|^\alpha \quad (5.1)$$

for all  $t, s \in [0, T]$  and some  $c \geq 0$ . We need some further definitions. If  $u \in H^1(\Omega)$  such that  $\Delta u \in L^2(\Omega)$  and if  $b \in L^2(\partial\Omega)$  then we write

$$\partial_\nu u = b \quad \text{if} \quad \int_\Omega \Delta u \bar{v} + \int_\Omega \nabla u \cdot \overline{\nabla v} = \int_{\partial\Omega} b \bar{v}, \quad \text{for all } v \in H^1(\Omega).$$

This means that we define the *normal derivative*  $\partial_\nu u$  of  $u$  by the validity of Green's formula. Now we can formulate our main result on the heat equation with non-autonomous Robin boundary conditions.



**Theorem 5.1.** *Let  $H = L^2(\Omega)$ ,  $f \in L^2(0, T; H)$ ,  $u_0 \in H^1(\Omega)$ . Then there exists a unique function  $u \in H^1(0, T; H) \cap L^2(0, T; H^1(\Omega))$  such that  $\Delta u \in L^2(0, T; H)$  and*

$$\begin{cases} \dot{u}(t) - \Delta u(t) &= f(t) & t - a.e. \\ \partial_\nu u(t) + B(t)u(t)|_{\partial\Omega} &= 0 & t - a.e. \\ u(0) &= u_0. \end{cases}$$

*Proof.* Given is  $\alpha > \frac{1}{4}$ . Central for the proof is a result by Jerison and Kenig (see [12, p. 165]; see also [13, Theorem 1, Ch. V.1, §1.1, p. 103]) which says that for  $0 < s < 1$  there is a unique bounded linear operator

$$\text{Tr}_s : H^{s+1/2}(\Omega) \rightarrow H^s(\partial\Omega)$$

such that  $\text{Tr}_s(u) = u|_{\partial\Omega}$  for all  $u \in H^{s+1/2}(\Omega) \cap \mathcal{C}(\bar{\Omega})$ . In particular,  $\text{Tr}_{1/2} = \text{Tr}$ . Moreover  $H^{1/2}(\partial\Omega) = \text{Tr}(H^1(\Omega))$ . Now choose  $0 < s < \frac{1}{2}$  such that  $\gamma := s + \frac{1}{2} < 2\alpha$ . Then  $\gamma < 1$  and  $\alpha > \frac{\gamma}{2}$  as needed in Section 4 for  $\omega(t) = ct^\alpha$ . Moreover, for  $u \in H^1(\Omega)$  we have

$$\begin{aligned} \|B(t)u|_{\partial\Omega} - B(s)u|_{\partial\Omega}\|_{L^2(\partial\Omega)} &\leq c|t - s|^\alpha \|u|_{\partial\Omega}\|_{L^2(\partial\Omega)} \\ &\leq c|t - s|^\alpha \|u|_{\partial\Omega}\|_{H^s(\partial\Omega)} \\ &\leq c|t - s|^\alpha \|u\|_{H^{s+1/2}(\Omega)}. \end{aligned}$$

Thus the form

$$\mathfrak{a}(t; u, v) := \int_{\Omega} \nabla u \cdot \overline{\nabla v} + \int_{\partial\Omega} B(t)u|_{\partial\Omega} \bar{v}|_{\partial\Omega}$$

defined on  $[0, T] \times H^1(\Omega) \times H^1(\Omega)$  satisfies condition (4.1).

We now choose  $\mu > \|B(\cdot)\|_{L^\infty(0, T; \mathcal{L}(L^2(\partial\Omega)))}$  so that the form

$$[0, T] \times H^1(\Omega) \times H^1(\Omega) \ni (t, u, v) \mapsto \mathfrak{a}(t; u, v) + \mu \int_{\Omega} u \bar{v}$$

satisfies (3.1), (3.2) in addition to (4.1). By Theorem 4.1 this perturbed form has maximal regularity in  $H$ . It follows from Proposition 3.4 that also the form  $\mathfrak{a}$  has maximal regularity in  $H$ . Denote by  $A(t)$  the operator associated with  $\mathfrak{a}(t; \cdot, \cdot)$  in  $H = L^2(\Omega)$ . Then

$$\begin{aligned} D(A(t)) &= \{u \in H^1(\Omega) : \Delta u \in L^2(\Omega), \partial_\nu u + B(t)u|_{\partial\Omega} = 0\} \\ A(t)u &= -\Delta u, \end{aligned}$$

as is easy to see using the definition of  $\partial_\nu u$  by Green's formula. Thus maximal regularity in  $H$  is exactly the statement of Theorem 5.1.  $\square$

Next we consider a non-linear problem. Keeping the assumptions and settings of this section we consider bounded continuous functions  $\beta_j : \mathbb{R} \rightarrow \mathbb{R}$ ,  $j = 0, 1, \dots, N$ .

**Theorem 5.2.** *Let  $u_0 \in H^1(\Omega)$ ,  $f \in L^2(0, T; L^2(\Omega))$ . Then there exists  $u \in H^1(0, T; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega))$  such that  $\Delta u \in L^2(0, T; L^2(\Omega))$  and*

$$\begin{cases} \dot{u}(t) - \Delta u(t) + \sum_{j=1}^N \beta_j(u(t)) \partial_j u(t) + \beta_0(u(t))u(t) &= f(t) & a.e. \text{ on } \Omega \\ \partial_\nu u(t) + B(t)u(t)|_{\partial\Omega} &= 0 & a.e. \text{ on } \partial\Omega \\ u(0) &= u_0. \end{cases} \quad (5.2)$$

*Proof.* We let  $\mathfrak{a}(t; \cdot, \cdot) : H^1(\Omega) \times H^1(\Omega) \rightarrow \mathbb{C}$  be defined as before. Given  $w \in L^2(0, T; L^2(\Omega))$  we define the form  $\mathfrak{a}_2^w : [0, T] \times H^1(\Omega) \times H^1(\Omega) \rightarrow \mathbb{C}$  by

$$\mathfrak{a}_2^w(t; u, v) = \int_{\Omega} \sum_{j=1}^N \beta_j(w(t)) \partial_j u v + \int_{\Omega} \beta_0(w(t)) u v, \quad u \in H^1(\Omega), v \in L^2(\Omega).$$

Then  $\mathfrak{a} + \mathfrak{a}_2^w$  satisfies (4.1) with  $\gamma = 2\alpha$  and with a constant  $c$  which does not depend on  $w \in L^2(0, T; L^2(\Omega))$ . Thus by Theorem 4.1, there exists a unique solution  $u$  belonging to the space

$$E := H^1(0, T; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega))$$

with  $\Delta u \in L^2(0, T; L^2(\Omega))$  of the problem

$$\begin{cases} \dot{u}(t) - \Delta u(t) + \sum_{j=1}^N \beta_j(w(t)) \partial_j u(t) + \beta_0(w(t)) u(t) &= f(t) \quad \text{a.e. on } \Omega \\ \partial_{\nu} u(t) + B(t) u(t)|_{\partial\Omega} &= 0 \quad \text{a.e. on } \partial\Omega \\ u(0) &= u_0. \end{cases}$$

We define  $Tw := u$ . Then  $T : L^2(0, T; L^2(\Omega)) \rightarrow L^2(0, T; L^2(\Omega))$  is a continuous mapping (as is easy to see). Moreover,  $TL^2(0, T; L^2(\Omega))$  is a bounded subset of  $E$ . This follows from Theorem 4.1. Since the embedding of  $H^1(\Omega)$  into  $L^2(\Omega)$  is compact (recall that  $\Omega$  is bounded), it follows from the Lemma of Aubin-Lions that the embedding of  $E$  into  $L^2(0, T; L^2(\Omega))$  is compact as well. It follows from Schauder's Fixed Point Theorem that  $T$  has a fixed point  $u$ . This function  $u$  solves the problem.  $\square$

## Acknowledgements

We are most grateful to El Maati Ouhabaz and Dominik Dier for fruitful and enjoyable discussions. We would like to thank the anonymous referee who helped to improve Section 4.

## References

- [1] P. Acquistapace and B. Terreni, *A unified approach to abstract linear nonautonomous parabolic equations*, Rend. Sem. Mat. Univ. Padova **78** (1987), 47–107. [1](#), [4](#)
- [2] W. Arendt, *Heat kernels. ISEM 2005/06*, available at [http://www.uni-ulm.de/fileadmin/website\\_uni\\_ulm/mawi.inst.020/arendt/downloads/internetseminar.pdf](http://www.uni-ulm.de/fileadmin/website_uni_ulm/mawi.inst.020/arendt/downloads/internetseminar.pdf), Ulm, 2006. [2](#)
- [3] W. Arendt, C.J.K. Batty, M. Hieber, and F. Neubrander, *Vector-valued Laplace transforms and Cauchy problems*, second ed., Monographs in Mathematics, vol. 96, Birkhäuser/Springer Basel AG, Basel, 2011. [2](#)
- [4] W. Arendt, D. Dier, H. Laasri, and E.M. Ouhabaz, *Maximal regularity for evolution equations governed by non autonomous forms*, Adv. Diff. Eq. **19** (2014), no. 11-12, 1043–1066. [1](#)
- [5] P. Auscher, S. Hofmann, M. Lacey, A. McIntosh, and Ph. Tchamitchian, *The solution of the Kato square root problem for second order elliptic operators on  $\mathbb{R}^n$* , Ann. of Math. (2) **156** (2002), no. 2, 633–654. [2](#)
- [6] P. Auscher and Ph. Tchamitchian, *Square roots of elliptic second order divergence operators on strongly Lipschitz domains:  $L^2$  theory*, J. Anal. Math. **90** (2003), 1–12. [2](#)

- [7] R. Dautray and J.-L. Lions, *Analyse mathématique et calcul numérique pour les sciences et les techniques. Vol. 8*, INSTN: Collection Enseignement. [INSTN: Teaching Collection], Masson, Paris, 1988, Évolution: semi-groupe, variationnel. [Evolution: semigroups, variational methods], Reprint of the 1985 edition. [3](#)
- [8] L. de Simon, *Un'applicazione della teoria degli integrali singolari allo studio delle equazioni differenziali lineari astratte del primo ordine*, Rend. Sem. Mat. Univ. Padova **34** (1964), 205–223. [2](#)
- [9] D. Dier, *Non-autonomous evolutionary problems governed by forms: maximal regularity and invariance*, PhD-Thesis, Ulm, 2014. [1](#)
- [10] H. Haak, B. and E.M. Ouhabaz, *Maximal regularity for non autonomous evolution equations*, available at <http://fr.arxiv.org/pdf/1402.1136>, to appear in Math. Ann., 2014. [1](#), [1](#), [4](#), [4](#)
- [11] M. Hieber and S. Monniaux, *Pseudo-differential operators and maximal regularity results for non-autonomous parabolic equations*, Proc. Amer. Math. Soc. **128** (2000), no. 4, 1047–1053. [4](#)
- [12] D. Jerison and C.E. Kenig, *The inhomogeneous Dirichlet problem in Lipschitz domains*, J. Funct. Anal. **130** (1995), no. 1, 161–219. [5](#)
- [13] A. Jonsson and H. Wallin, *Function spaces on subsets of  $\mathbb{R}^n$* , Math. Rep. **2** (1984), no. 1, xiv+221. [5](#)
- [14] J.-L. Lions, *Équations différentielles opérationnelles et problèmes aux limites*, Die Grundlehren der mathematischen Wissenschaften, Bd. 111, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1961. [1](#)
- [15] J.-L. Lions, *Espaces d'interpolation et domaines de puissances fractionnaires d'opérateurs*, J. Math. Soc. Japan **14** (1962), 233–241. [2](#)
- [16] J.-L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications. Vol. 1*, Travaux et Recherches Mathématiques, No. 17, Dunod, Paris, 1968. [2](#)
- [17] A. McIntosh, *On the comparability of  $A^{1/2}$  and  $A^{*1/2}$* , Proc. Amer. Math. Soc. **32** (1972), 430–434. [2.2](#)
- [18] E.M. Ouhabaz, *Analysis of heat equations on domains*, London Mathematical Society Monographs Series, vol. 31, Princeton University Press, Princeton, NJ, 2005. [2](#)
- [19] E.M. Ouhabaz, *Maximal regularity for non-autonomous evolution equations governed by forms having less regularity*, available at <http://fr.arxiv.org/pdf/1411.0139v1>, 2014. [1](#)
- [20] E.M. Ouhabaz and C. Spina, *Maximal regularity for non-autonomous Schrödinger type equations*, J. Differential Equations **248** (2010), no. 7, 1668–1683. [1](#)
- [21] R. E. Showalter, *Monotone operators in Banach space and nonlinear partial differential equations*, Mathematical Surveys and Monographs, vol. 49, American Mathematical Society, Providence, RI, 1997. [3](#)
- [22] H. Tanabe, *Equations of evolution*, Monographs and Studies in Mathematics, vol. 6, Pitman (Advanced Publishing Program), Boston, Mass., 1979, Translated from the Japanese by N. Mugibayashi and H. Haneda. [2](#)